

**$\rho(1450)$** 

$$I^G(J^{PC}) = 1^+(1^{--})$$

See our mini-review under the  $\rho(1700)$ . **$\rho(1450)$  MASS**VALUE (MeV)DOCUMENT ID**1465 ± 25 OUR ESTIMATE** This is only an educated guess; the error given is larger than the error on the average of the published values. **$\eta\rho^0$  MODE**VALUE (MeV)DOCUMENT IDTECNCOMMENT

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

1497 ± 14	<sup>1</sup> AKHMETSHIN 01B	CMD2	$e^+e^- \rightarrow \eta\gamma$
1421 ± 15	<sup>2</sup> AKHMETSHIN 00D	CMD2	$e^+e^- \rightarrow \eta\pi^+\pi^-$
1470 ± 20	ANTONELLI 88	DM2	$e^+e^- \rightarrow \eta\pi^+\pi^-$
1446 ± 10	FUKUI 88	SPEC	$8.95 \pi^- p \rightarrow \eta\pi^+\pi^- n$

<sup>1</sup> Using the data of AKHMETSHIN 01B on  $e^+e^- \rightarrow \eta\gamma$ , AKHMETSHIN 00D and ANTONELLI 88 on  $e^+e^- \rightarrow \eta\pi^+\pi^-$ .<sup>2</sup> Using the data of ANTONELLI 88, DOLINSKY 91, and AKHMETSHIN 00D. The energy-independent width of the  $\rho(1450)$  and  $\rho(1700)$  mesons assumed. **$\omega\pi$  MODE**VALUE (MeV)EVTSDOCUMENT IDTECNCOMMENT

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

1510 ± 7	10.2k	<sup>1</sup> ACHASOV 16D	SND	1.05–2.00 $e^+e^- \rightarrow \pi^0\pi^0\gamma$
1544 ± 22 <sup>+11</sup> <sub>-46</sub>	821	<sup>2</sup> MATVIENKO 15	BELL	$\bar{B}^0 \rightarrow D^{*+}\omega\pi^-$
1491 ± 19	7815	<sup>3</sup> ACHASOV 13	SND	1.05–2.00 $e^+e^- \rightarrow \pi^0\pi^0\gamma$
1582 ± 17 ± 25	2382	<sup>4</sup> AKHMETSHIN 03B	CMD2	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
1349 ± 25 <sup>+10</sup> <sub>-5</sub>	341	<sup>5</sup> ALEXANDER 01B	CLE2	$B \rightarrow D^{(*)}\omega\pi^-$
1523 ± 10		<sup>6</sup> EDWARDS 00A	CLE2	$\tau^- \rightarrow \omega\pi^- \nu_\tau$
1463 ± 25		<sup>7</sup> CLEGG 94	RVUE	
1250		<sup>8</sup> ASTON 80C	OMEG	20–70 $\gamma p \rightarrow \omega\pi^0 p$
1290 ± 40		<sup>8</sup> BARBER 80C	SPEC	3–5 $\gamma p \rightarrow \omega\pi^0 p$

<sup>1</sup> From a phenomenological model based on vector meson dominance with interfering  $\rho(770)$ ,  $\rho(1450)$ , and  $\rho(1700)$ . The  $\rho(1700)$  mass and width are fixed at 1720 MeV and 250 MeV, respectively. Systematic uncertainties not estimated. Supersedes ACHASOV 13.<sup>2</sup> Using Breit-Wigner parameterization of the  $\rho(1450)$  and assuming equal probabilities of the  $\rho(1450) \rightarrow \pi\pi$  and  $\rho(1450) \rightarrow \omega\pi$  decays.<sup>3</sup> From a phenomenological model based on vector meson dominance with the interfering  $\rho(1450)$  and  $\rho(1700)$  and their widths fixed at 400 and 250 MeV, respectively. Systematic uncertainty not estimated.<sup>4</sup> Using the data of AKHMETSHIN 03B and BISELLO 91B assuming the  $\omega\pi^0$  and  $\pi^+\pi^-$  mass dependence of the total width.  $\rho(1700)$  mass and width fixed at 1700 MeV and 240 MeV, respectively.<sup>5</sup> Using Breit-Wigner parameterization of the  $\rho(1450)$  and assuming the  $\omega\pi^-$  mass dependence for the total width.

<sup>6</sup> Mass-independent width parameterization.  $\rho(1700)$  mass and width fixed at 1700 MeV and 235 MeV respectively.

<sup>7</sup> Using data from BISELLO 91B, DOLINSKY 86 and ALBRECHT 87L.

<sup>8</sup> Not separated from  $b_1(1235)$ , not pure  $J^P = 1^-$  effect.

## 4 $\pi$ MODE

VALUE (MeV)		DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1435 ± 40		ABELE	01B CBAR	0.0 $\bar{p}n \rightarrow 2\pi^- 2\pi^0 \pi^+$
1350 ± 50		ACHASOV	97 RVUE	$e^+e^- \rightarrow 2(\pi^+\pi^-)$
1449 ± 4		<sup>1</sup> ARMSTRONG	89E OMEG 300	$pp \rightarrow p\rho 2(\pi^+\pi^-)$

<sup>1</sup> Not clear whether this observation has  $l=1$  or 0.

## $\pi\pi$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1350 ± 20	$\begin{smallmatrix} +20 \\ -30 \end{smallmatrix}$ 63.5k	<sup>1</sup> ABRAMOWICZ12	ZEUS	$ep \rightarrow e\pi^+\pi^-p$
1493 ± 15		<sup>2</sup> LEES	12G BABR	$e^+e^- \rightarrow \pi^+\pi^-\gamma$
1446 ± 7	±28 5.4M	<sup>3,4</sup> FUJIKAWA	08 BELL	$\tau^- \rightarrow \pi^-\pi^0\nu_\tau$
1328 ± 15		<sup>5</sup> SCHAEEL	05C ALEP	$\tau^- \rightarrow \pi^-\pi^0\nu_\tau$
1406 ± 15	87k	<sup>3,6</sup> ANDERSON	00A CLE2	$\tau^- \rightarrow \pi^-\pi^0\nu_\tau$
~ 1368		<sup>7</sup> ABELE	99C CBAR	0.0 $\bar{p}d \rightarrow \pi^+\pi^-\pi^-p$
1348 ± 33		BERTIN	98 OBLX	0.05–0.405 $\bar{p}p \rightarrow 2\pi^+\pi^-$
1411 ± 14		<sup>8</sup> ABELE	97 CBAR	$\bar{p}n \rightarrow \pi^-\pi^0\pi^0$
1370 $\begin{smallmatrix} +90 \\ -70 \end{smallmatrix}$		ACHASOV	97 RVUE	$e^+e^- \rightarrow \pi^+\pi^-$
1359 ± 40		<sup>6</sup> BERTIN	97C OBLX	0.0 $\bar{p}p \rightarrow \pi^+\pi^-\pi^0$
1282 ± 37		BERTIN	97D OBLX	0.05 $\bar{p}p \rightarrow 2\pi^+2\pi^-$
1424 ± 25		BISELLO	89 DM2	$e^+e^- \rightarrow \pi^+\pi^-$
1265.5 ± 75.3		DUBNICKA	89 RVUE	$e^+e^- \rightarrow \pi^+\pi^-$
1292 ± 17		<sup>9</sup> KURDADZE	83 OLYA	0.64–1.4 $e^+e^- \rightarrow \pi^+\pi^-$

<sup>1</sup> Using the KUHN 90 parametrization of the pion form factor, neglecting  $\rho$ - $\omega$  interference.

<sup>2</sup> Using the GOUNARIS 68 parametrization of the pion form factor leaving the masses and widths of the  $\rho(1450)$ ,  $\rho(1700)$ , and  $\rho(2150)$  resonances as free parameters of the fit.

<sup>3</sup> From the GOUNARIS 68 parametrization of the pion form factor.

<sup>4</sup>  $|F_\pi(0)|^2$  fixed to 1.

<sup>5</sup> From the combined fit of the  $\tau^-$  data from ANDERSON 00A and SCHAEEL 05C and  $e^+e^-$  data from the compilation of BARKOV 85, AKHMETSHIN 04, and ALOISIO 05.  $\rho(1700)$  mass and width fixed at 1713 MeV and 235 MeV, respectively. Supersedes BARATE 97M.

<sup>6</sup>  $\rho(1700)$  mass and width fixed at 1700 MeV and 235 MeV, respectively.

<sup>7</sup>  $\rho(1700)$  mass and width fixed at 1780 MeV and 275 MeV respectively.

<sup>8</sup> T-matrix pole.

<sup>9</sup> Using for  $\rho(1700)$  mass and width  $1600 \pm 20$  and  $300 \pm 10$  MeV respectively.

**$K\bar{K}$  MODE**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1208 ± 8 ± 9	190k	<sup>1</sup> AAIJ	16N	LHCB	$D^0 \rightarrow K_S^0 K^\pm \pi^\mp$
1422.8 ± 6.5	27k	<sup>2</sup> ABELE	99D	CBAR ±	$0.0 \bar{p} p \rightarrow K^+ K^- \pi^0$

<sup>1</sup> Using the GOUNARIS 68 parameterization with fixed width.

<sup>2</sup> K-matrix pole. Isospin not determined, could be  $\omega(1420)$ .

 **$K\bar{K}^*(892) + c.c.$  MODE**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1505 ± 19 ± 7	AUBERT	08S BABR	$10.6 e^+ e^- \rightarrow K\bar{K}^*(892)\gamma$

 **$\rho(1450)$  WIDTH**

VALUE (MeV) DOCUMENT ID  
**400 ± 60 OUR ESTIMATE** This is only an educated guess; the error given is larger than the error on the average of the published values.

 **$\eta\rho^0$  MODE**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
226 ± 44	<sup>1</sup> AKHMETSHIN 01B	CMD2	$e^+ e^- \rightarrow \eta\gamma$
211 ± 31	<sup>2</sup> AKHMETSHIN 00D	CMD2	$e^+ e^- \rightarrow \eta\pi^+\pi^-$
230 ± 30	ANTONELLI 88	DM2	$e^+ e^- \rightarrow \eta\pi^+\pi^-$
60 ± 15	FUKUI 88	SPEC	$8.95 \pi^- p \rightarrow \eta\pi^+\pi^- n$

<sup>1</sup> Using the data of AKHMETSHIN 01B on  $e^+ e^- \rightarrow \eta\gamma$ , AKHMETSHIN 00D and ANTONELLI 88 on  $e^+ e^- \rightarrow \eta\pi^+\pi^-$ .

<sup>2</sup> Using the data of ANTONELLI 88, DOLINSKY 91, and AKHMETSHIN 00D. The energy-independent width of the  $\rho(1450)$  and  $\rho(1700)$  mesons assumed.

 **$\omega\pi$  MODE**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
440 ± 40	10.2k	<sup>1</sup> ACHASOV	16D	SND $1.05-2.00 e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
$303^{+31}_{-52} \pm 69_{-7}$	821	<sup>2</sup> MATVIENKO	15	BELL $\bar{B}^0 \rightarrow D^{*+} \omega \pi^-$
429 ± 42 ± 10	2382	<sup>3</sup> AKHMETSHIN 03B	CMD2	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
547 ± 86 $^{+46}_{-45}$	341	<sup>4</sup> ALEXANDER	01B	CLE2 $B \rightarrow D^{(*)} \omega \pi^-$
400 ± 35		<sup>5</sup> EDWARDS	00A	CLE2 $\tau^- \rightarrow \omega \pi^- \nu_\tau$
311 ± 62		<sup>6</sup> CLEGG	94	RVUE
300		<sup>7</sup> ASTON	80C	OMEG $20-70 \gamma p \rightarrow \omega \pi^0 p$
320 ± 100		<sup>7</sup> BARBER	80C	SPEC $3-5 \gamma p \rightarrow \omega \pi^0 p$

<sup>1</sup> From a phenomenological model based on vector meson dominance with interfering  $\rho(770)$ ,  $\rho(1450)$ , and  $\rho(1700)$ . The  $\rho(1700)$  mass and width are fixed at 1720 MeV and 250 MeV, respectively. Systematic uncertainties not estimated. Supersedes ACHASOV 13.

- <sup>2</sup> Using Breit-Wigner parameterization of the  $\rho(1450)$  and assuming equal probabilities of the  $\rho(1450) \rightarrow \pi\pi$  and  $\rho(1450) \rightarrow \omega\pi$  decays.
- <sup>3</sup> Using the data of AKHMETSHIN 03B and BISELLO 91B assuming the  $\omega\pi^0$  and  $\pi^+\pi^-$  mass dependence of the total width.  $\rho(1700)$  mass and width fixed at 1700 MeV and 240 MeV, respectively.
- <sup>4</sup> Using Breit-Wigner parameterization of the  $\rho(1450)$  and assuming the  $\omega\pi^-$  mass dependence for the total width.
- <sup>5</sup> Mass-independent width parameterization.  $\rho(1700)$  mass and width fixed at 1700 MeV and 235 MeV respectively.
- <sup>6</sup> Using data from BISELLO 91B, DOLINSKY 86 and ALBRECHT 87L.
- <sup>7</sup> Not separated from  $b_1(1235)$ , not pure  $J^P = 1^-$  effect.

## 4 $\pi$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$325 \pm 100$	ABELE	01B	CBAR $0.0 \bar{p}n \rightarrow 2\pi^- 2\pi^0 \pi^+$
---------------	-------	-----	---

## $\pi\pi$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$460 \pm 30^{+40}_{-45}$	63.5k	<sup>1</sup> ABRAMOWICZ12	ZEUS	$e p \rightarrow e \pi^+ \pi^- p$
$427 \pm 31$		<sup>2</sup> LEES	12G	BABR $e^+ e^- \rightarrow \pi^+ \pi^- \gamma$
$434 \pm 16 \pm 60$	5.4M	<sup>3,4</sup> FUJIKAWA	08	BELL $\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
$468 \pm 41$		<sup>5</sup> SCHAEEL	05C	ALEP $\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
$455 \pm 41$	87k	<sup>3,6</sup> ANDERSON	00A	CLE2 $\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
$\sim 374$		<sup>7</sup> ABELE	99C	CBAR $0.0 \bar{p}d \rightarrow \pi^+ \pi^- \pi^- p$
$275 \pm 10$		BERTIN	98	OBLX $0.05-0.405 \bar{n}p \rightarrow \pi^+ \pi^+ \pi^-$
$343 \pm 20$		<sup>8</sup> ABELE	97	CBAR $\bar{p}n \rightarrow \pi^- \pi^0 \pi^0$
$310 \pm 40$		<sup>6</sup> BERTIN	97C	OBLX $0.0 \bar{p}p \rightarrow \pi^+ \pi^- \pi^0$
$236 \pm 36$		BERTIN	97D	OBLX $0.05 \bar{p}p \rightarrow 2\pi^+ 2\pi^-$
$269 \pm 31$		BISELLO	89	DM2 $e^+ e^- \rightarrow \pi^+ \pi^-$
$391 \pm 70$		DUBNICKA	89	RVUE $e^+ e^- \rightarrow \pi^+ \pi^-$
$218 \pm 46$		<sup>9</sup> KURDADZE	83	OLYA $0.64-1.4 e^+ e^- \rightarrow \pi^+ \pi^-$

<sup>1</sup> Using the KUHN 90 parametrization of the pion form factor, neglecting  $\rho-\omega$  interference.

<sup>2</sup> Using the GOUNARIS 68 parametrization of the pion form factor leaving the masses and widths of the  $\rho(1450)$ ,  $\rho(1700)$ , and  $\rho(2150)$  resonances as free parameters of the fit.

<sup>3</sup> From the GOUNARIS 68 parametrization of the pion form factor.

<sup>4</sup>  $|F_\pi(0)|^2$  fixed to 1.

<sup>5</sup> From the combined fit of the  $\tau^-$  data from ANDERSON 00A and SCHAEEL 05C and  $e^+ e^-$  data from the compilation of BARKOV 85, AKHMETSHIN 04, and ALOISIO 05.  $\rho(1700)$  mass and width fixed at 1713 MeV and 235 MeV, respectively. Supersedes BARATE 97M.

<sup>6</sup>  $\rho(1700)$  mass and width fixed at 1700 MeV and 235 MeV, respectively.

<sup>7</sup>  $\rho(1700)$  mass and width fixed at 1780 MeV and 275 MeV respectively.

<sup>8</sup> T-matrix pole.

<sup>9</sup> Using for  $\rho(1700)$  mass and width  $1600 \pm 20$  and  $300 \pm 10$  MeV respectively.

### $K\bar{K}$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
410 ± 19 ± 35	190k	<sup>1</sup> AAIJ	16N	LHCB	$D^0 \rightarrow K_S^0 K^\pm \pi^\mp$
146.5 ± 10.5	27k	<sup>2</sup> ABELE	99D	CBAR ±	$0.0 \bar{p}p \rightarrow K^+ K^- \pi^0$

<sup>1</sup> Using the GOUNARIS 68 parameterization with fixed mass.

<sup>2</sup> K-matrix pole. Isospin not determined, could be  $\omega(1420)$ .

### $K\bar{K}^*(892) + \text{c.c.}$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
418 ± 25 ± 4	AUBERT	08S BABR	10.6 $e^+ e^- \rightarrow K\bar{K}^*(892)\gamma$

### $\rho(1450)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\pi\pi$	seen
$\Gamma_2$ $4\pi$	seen
$\Gamma_3$ $\omega\pi$	
$\Gamma_4$ $a_1(1260)\pi$	
$\Gamma_5$ $h_1(1170)\pi$	
$\Gamma_6$ $\pi(1300)\pi$	
$\Gamma_7$ $\rho\rho$	
$\Gamma_8$ $\rho(\pi\pi)$ S-wave	
$\Gamma_9$ $e^+ e^-$	seen
$\Gamma_{10}$ $\eta\rho$	seen
$\Gamma_{11}$ $a_2(1320)\pi$	not seen
$\Gamma_{12}$ $K\bar{K}$	not seen
$\Gamma_{13}$ $K\bar{K}^*(892) + \text{c.c.}$	possibly seen
$\Gamma_{14}$ $\eta\gamma$	seen
$\Gamma_{15}$ $f_0(500)\gamma$	not seen
$\Gamma_{16}$ $f_0(980)\gamma$	not seen
$\Gamma_{17}$ $f_0(1370)\gamma$	not seen
$\Gamma_{18}$ $f_2(1270)\gamma$	not seen

### $\rho(1450) \Gamma(i)\Gamma(e^+ e^-)/\Gamma(\text{total})$

$\Gamma(\pi\pi) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$   $\Gamma_1\Gamma_9/\Gamma$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
0.12	<sup>1</sup> DIEKMAN	88	RVUE $e^+ e^- \rightarrow \pi^+ \pi^-$
0.027 <sup>+0.015</sup> <sub>-0.010</sub>	<sup>2</sup> KURDADZE	83	OLYA $0.64-1.4 e^+ e^- \rightarrow \pi^+ \pi^-$

<sup>1</sup> Using total width = 235 MeV.

<sup>2</sup> Using for  $\rho(1700)$  mass and width  $1600 \pm 20$  and  $300 \pm 10$  MeV respectively.

$\Gamma(\eta\rho) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{10}\Gamma_9/\Gamma$

VALUE (eV)                      DOCUMENT ID      TECN      COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

74 ± 20	<sup>1</sup> AKHMETSHIN 00D	CMD2	$e^+e^- \rightarrow \eta\pi^+\pi^-$
91 ± 19	ANTONELLI 88	DM2	$e^+e^- \rightarrow \eta\pi^+\pi^-$

<sup>1</sup> Using the data of ANTONELLI 88, DOLINSKY 91, and AKHMETSHIN 00D. The energy-independent width of the  $\rho(1450)$  and  $\rho(1700)$  mesons assumed.

$\Gamma(\eta\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{14}\Gamma_9/\Gamma$

VALUE (eV)                      DOCUMENT ID      TECN      COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

<16.4	<sup>1</sup> AKHMETSHIN 05	CMD2	0.60-1.38 $e^+e^- \rightarrow \eta\gamma$
2.2 ± 0.5 ± 0.3	<sup>2</sup> AKHMETSHIN 01B	CMD2	$e^+e^- \rightarrow \eta\gamma$

<sup>1</sup> From  $2\gamma$  decay mode of  $\eta$  using 1465 MeV and 310 MeV for the  $\rho(1450)$  mass and width. Recalculated by us.

<sup>2</sup> Using the data of AKHMETSHIN 01B on  $e^+e^- \rightarrow \eta\gamma$ , AKHMETSHIN 00D and ANTONELLI 88 on  $e^+e^- \rightarrow \eta\pi^+\pi^-$ . Recalculated by us using width of 226 MeV.

$\Gamma(K\bar{K}^*(892) + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{13}\Gamma_9/\Gamma$

VALUE (eV)                      DOCUMENT ID      TECN      COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

127 ± 15 ± 6	AUBERT	08s	BABR 10.6 $e^+e^- \rightarrow K\bar{K}^*(892)\gamma$
--------------	--------	-----	--

$\rho(1450) \Gamma(i)/\Gamma(\text{total}) \times \Gamma(e^+e^-)/\Gamma(\text{total})$

$\Gamma(\omega\pi)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_3/\Gamma \times \Gamma_9/\Gamma$

VALUE (units 10<sup>-6</sup>)      EVTS                      DOCUMENT ID      TECN      COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.1 ± 0.4	10.2k	<sup>1</sup> ACHASOV	16D	SND	1.05-2.00 $e^+e^- \rightarrow \pi^0\pi^0\gamma$
5.3 ± 0.4	7815	<sup>2</sup> ACHASOV	13	SND	1.05-2.00 $e^+e^- \rightarrow \pi^0\pi^0\gamma$

<sup>1</sup> From a phenomenological model based on vector meson dominance with interfering  $\rho(770)$ ,  $\rho(1450)$ , and  $\rho(1700)$ . The  $\rho(1700)$  mass and width are fixed at 1720 MeV and 250 MeV, respectively. Systematic uncertainties not estimated. Supersedes ACHASOV 13.

<sup>2</sup> From a phenomenological model based on vector meson dominance with the interfering  $\rho(1450)$  and  $\rho(1700)$  and their widths fixed at 400 and 250 MeV, respectively. Systematic uncertainty not estimated.

$\Gamma(\eta\rho)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{10}/\Gamma \times \Gamma_9/\Gamma$

VALUE (units 10<sup>-7</sup>)      EVTS                      DOCUMENT ID      TECN      COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.3 <sup>+1.1</sup> <sub>-0.9</sub> ± 0.2	4.9k	<sup>1</sup> AULCHENKO	15	SND	1.22-2.00 $e^+e^- \rightarrow \eta\pi^+\pi^-$
---	------	------------------------	----	-----	---

<sup>1</sup> From a fit to the  $e^+e^- \rightarrow \eta\pi^+\pi^-$  cross section with vector meson dominance model including  $\rho(770)$ ,  $\rho(1450)$ , and  $\rho(1700)$  decaying exclusively via  $\eta\rho(770)$ . Masses and widths of vector states are fixed to PDG 14. Coupling constants are assumed to be real.

$$\Gamma(f_0(500)\gamma)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \qquad \Gamma_{15}/\Gamma \times \Gamma_9/\Gamma$$

<u>VALUE (units 10<sup>-9</sup>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<4.0	90	ACHASOV 11	SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$

$$\Gamma(f_0(980)\gamma)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \qquad \Gamma_{16}/\Gamma \times \Gamma_9/\Gamma$$

<u>VALUE (units 10<sup>-9</sup>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2.6	90	ACHASOV 11	SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$

$$\Gamma(f_0(1370)\gamma)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \qquad \Gamma_{17}/\Gamma \times \Gamma_9/\Gamma$$

<u>VALUE (units 10<sup>-9</sup>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<3.5	90	ACHASOV 11	SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$

$$\Gamma(f_2(1270)\gamma)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \qquad \Gamma_{18}/\Gamma \times \Gamma_9/\Gamma$$

<u>VALUE (units 10<sup>-9</sup>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.8	90	<sup>1</sup> ACHASOV 11	SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$

<sup>1</sup> Using Breit-Wigner parametrization of the  $\rho(1450)$  with mass and width of 1465 MeV and 400 MeV, respectively.

### $\rho(1450)$ BRANCHING RATIOS

$$\Gamma(\pi\pi)/\Gamma(4\pi) \qquad \Gamma_1/\Gamma_2$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.37 ± 0.10	1,2 ABELE	01B	CBAR $0.0 \bar{p}n \rightarrow 5\pi$
-------------	-----------	-----	--------------------------------------

<sup>1</sup>  $\omega\pi$  not included.

<sup>2</sup> Using ABELE 97.

$$\Gamma(\omega\pi)/\Gamma_{\text{total}} \qquad \Gamma_3/\Gamma$$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	-------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	821	<sup>1</sup> MATVIENKO 15	BELL	$\bar{B}^0 \rightarrow D^{*+}\omega\pi^-$
seen	1.6k	ACHASOV 12	SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
~ 0.21		CLEGG 94	RVUE	

<sup>1</sup> Using Breit-Wigner parameterization of the  $\rho(1450)$  and assuming equal probabilities of the  $\rho(1450) \rightarrow \pi\pi$  and  $\rho(1450) \rightarrow \omega\pi$  decays.

$$\Gamma(\pi\pi)/\Gamma(\omega\pi) \qquad \Gamma_1/\Gamma_3$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
--------------	--------------------	-------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 0.32	CLEGG 94	RVUE
--------	----------	------

$$\Gamma(\omega\pi)/\Gamma(4\pi) \qquad \Gamma_3/\Gamma_2$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
--------------	--------------------	-------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.14	CLEGG 88	RVUE
-------	----------	------

### $\Gamma(a_1(1260)\pi)/\Gamma(4\pi)$

$\Gamma_4/\Gamma_2$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.27 \pm 0.08$	<sup>1</sup> ABELE	01B	CBAR $0.0 \bar{p}n \rightarrow 5\pi$
<sup>1</sup> $\omega\pi$ not included.			

### $\Gamma(h_1(1170)\pi)/\Gamma(4\pi)$

$\Gamma_5/\Gamma_2$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.08 \pm 0.04$	<sup>1</sup> ABELE	01B	CBAR $0.0 \bar{p}n \rightarrow 5\pi$
<sup>1</sup> $\omega\pi$ not included.			

### $\Gamma(\pi(1300)\pi)/\Gamma(4\pi)$

$\Gamma_6/\Gamma_2$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.37 \pm 0.13$	<sup>1</sup> ABELE	01B	CBAR $0.0 \bar{p}n \rightarrow 5\pi$
<sup>1</sup> $\omega\pi$ not included.			

### $\Gamma(\rho\rho)/\Gamma(4\pi)$

$\Gamma_7/\Gamma_2$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.11 \pm 0.05$	<sup>1</sup> ABELE	01B	CBAR $0.0 \bar{p}n \rightarrow 5\pi$
<sup>1</sup> $\omega\pi$ not included.			

### $\Gamma(\rho(\pi\pi)_{S\text{-wave}})/\Gamma(4\pi)$

$\Gamma_8/\Gamma_2$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.17 \pm 0.09$	<sup>1</sup> ABELE	01B	CBAR $0.0 \bar{p}n \rightarrow 5\pi$
<sup>1</sup> $\omega\pi$ not included.			

### $\Gamma(\eta\rho)/\Gamma_{\text{total}}$

$\Gamma_{10}/\Gamma$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>seen</b>	35	<sup>1</sup> ACHASOV	14	SND $1.15\text{--}2.00 e^+e^- \rightarrow \eta\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<0.04$		DONNACHIE	87B	RVUE
<sup>1</sup> From a phenomenological model based on vector meson dominance with $\rho(1450)$ and $\phi(1680)$ masses and widths from the PDG 12.				

### $\Gamma(\eta\rho)/\Gamma(\omega\pi)$

$\Gamma_{10}/\Gamma_3$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.081 \pm 0.020$	<sup>1,2</sup> AULCHENKO	15	SND $1.22\text{--}2.00 e^+e^- \rightarrow \eta\pi^+\pi^-$
$\sim 0.24$	<sup>3</sup> DONNACHIE	91	RVUE
$>2$	FUKUI	91	SPEC $8.95 \pi^- p \rightarrow \omega\pi^0 n$

<sup>1</sup> From a fit to the  $e^+e^- \rightarrow \eta\pi^+\pi^-$  cross section with vector meson dominance model including  $\rho(770)$ ,  $\rho(1450)$ , and  $\rho(1700)$  decaying exclusively via  $\eta\rho(770)$ . Masses and widths of vector states are fixed to PDG 14. Coupling constants are assumed to be real.

<sup>2</sup> Reports the inverse of the quoted value as  $12.3 \pm 3.1$ .

<sup>3</sup> Using data from BISELLO 91B, DOLINSKY 86 and ALBRECHT 87L.





ALEXANDER	01B	PR D64 092001	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
AKHMETSHIN	00D	PL B489 125	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
AMELIN	00	NP A668 83	D. Amelin <i>et al.</i>	(VES Collab.)
ANDERSON	00A	PR D61 112002	S. Anderson <i>et al.</i>	(CLEO Collab.)
EDWARDS	00A	PR D61 072003	K.W. Edwards <i>et al.</i>	(CLEO Collab.)
ABELE	99C	PL B450 275	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
ABELE	99D	PL B468 178	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
BERTIN	98	PR D57 55	A. Bertin <i>et al.</i>	(OBELIX Collab.)
ABELE	97	PL B391 191	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
ACHASOV	97	PR D55 2663	N.N. Achasov <i>et al.</i>	(NOVM)
BARATE	97M	ZPHY C76 15	R. Barate <i>et al.</i>	(ALEPH Collab.)
BERTIN	97C	PL B408 476	A. Bertin <i>et al.</i>	(OBELIX Collab.)
BERTIN	97D	PL B414 220	A. Bertin <i>et al.</i>	(OBELIX Collab.)
CLEGG	94	ZPHY C62 455	A.B. Clegg, A. Donnachie	(LANC, MCHS)
BISELLO	91B	NPBPS B21 111	D. Bisello	(DM2 Collab.)
DOLINSKY	91	PRPL 202 99	S.I. Dolinsky <i>et al.</i>	(NOVO)
DONNACHIE	91	ZPHY C51 689	A. Donnachie, A.B. Clegg	(MCHS, LANC)
FUKUI	91	PL B257 241	S. Fukui <i>et al.</i>	(SUGI, NAGO, KEK, KYOT+)
KUHN	90	ZPHY C48 445	J.H. Kuhn <i>et al.</i>	(MPIM)
ARMSTRONG	89E	PL B228 536	T.A. Armstrong, M. Benayoun	(ATHU, BARI, BIRM+)
BISELLO	89	PL B220 321	D. Bisello <i>et al.</i>	(DM2 Collab.)
DUBNICKA	89	JP G15 1349	S. Dubnicka <i>et al.</i>	(JINR, SLOV)
ANTONELLI	88	PL B212 133	A. Antonelli <i>et al.</i>	(DM2 Collab.)
CLEGG	88	ZPHY C40 313	A.B. Clegg, A. Donnachie	(MCHS, LANC)
DIEKMAN	88	PRPL 159 99	B. Diekmann	(BONN)
FUKUI	88	PL B202 441	S. Fukui <i>et al.</i>	(SUGI, NAGO, KEK, KYOT+)
ALBRECHT	87L	PL B185 223	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
DONNACHIE	87B	ZPHY C34 257	A. Donnachie, A.B. Clegg	(MCHS, LANC)
DOLINSKY	86	PL B174 453	S.I. Dolinsky <i>et al.</i>	(NOVO)
BARKOV	85	NP B256 365	L.M. Barkov <i>et al.</i>	(NOVO)
KURDADZE	83	JETPL 37 733	L.M. Kurdadze <i>et al.</i>	(NOVO)
ASTON	80C	Translated from ZETFP 37 613 PL 92B 211	D. Aston	(BONN, CERN, EPOL, GLAS, LANC+)
BARBER	80C	ZPHY C4 169	D.P. Barber <i>et al.</i>	(DARE, LANC, SHEF)
GOUNARIS	68	PRL 21 244	G.J. Gounaris, J.J. Sakurai	