

QUARKS

The u -, d -, and s -quark masses are estimates of so-called “current-quark masses,” in a mass-independent subtraction scheme such as $\overline{\text{MS}}$ at a scale $\mu \approx 2$ GeV. The c - and b -quark masses are the “running” masses in the $\overline{\text{MS}}$ scheme. For the b -quark we also quote the 1S mass. These can be different from the heavy quark masses obtained in potential models.

u

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

$$m_u = 2.2^{+0.6}_{-0.4} \text{ MeV} \quad \text{Charge} = \frac{2}{3} e \quad I_z = +\frac{1}{2}$$

$$m_u/m_d = 0.38\text{--}0.58$$

d

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

$$m_d = 4.7^{+0.5}_{-0.4} \text{ MeV} \quad \text{Charge} = -\frac{1}{3} e \quad I_z = -\frac{1}{2}$$

$$m_s/m_d = 17\text{--}22$$

$$\bar{m} = (m_u + m_d)/2 = 3.5^{+0.7}_{-0.3} \text{ MeV}$$

s

$$I(J^P) = 0(\frac{1}{2}^+)$$

$$m_s = 96^{+8}_{-4} \text{ MeV} \quad \text{Charge} = -\frac{1}{3} e \quad \text{Strangeness} = -1$$

$$m_s / ((m_u + m_d)/2) = 27.3 \pm 0.7$$

c

$$I(J^P) = 0(\frac{1}{2}^+)$$

$$m_c = 1.28 \pm 0.03 \text{ GeV} \quad \text{Charge} = \frac{2}{3} e \quad \text{Charm} = +1$$

$$m_c/m_s = 11.72 \pm 0.25$$

$$m_b/m_c = 4.53 \pm 0.05$$

$$m_b - m_c = 3.45 \pm 0.05 \text{ GeV}$$

b

$$I(J^P) = 0(\frac{1}{2}^+)$$

$$\text{Charge} = -\frac{1}{3} e \quad \text{Bottom} = -1$$

$$\text{Mass } m = 4.18^{+0.04}_{-0.03} \text{ GeV} \quad \text{Charge} = -\frac{1}{3} e \quad \text{Bottom} = -1$$

t

$$I(J^P) = 0(\frac{1}{2}^+)$$

$$\text{Charge} = \frac{2}{3} e \quad \text{Top} = +1$$

Mass (direct measurements) $m = 173.1 \pm 0.6 \text{ GeV}^{[a,b]}$ (S = 1.6)

Mass from cross-section measurements) $m = 160_{-4}^{+5} \text{ GeV}^{[a]}$

Mass (Pole from cross-section measurements) $m = 173.5 \pm 1.1 \text{ GeV}$

$m_t - m_{\bar{t}} = -0.2 \pm 0.5 \text{ GeV}$ (S = 1.1)

Full width $\Gamma = 1.41_{-0.15}^{+0.19} \text{ GeV}$ (S = 1.4)

$\Gamma(Wb)/\Gamma(Wq(q = b, s, d)) = 0.957 \pm 0.034$ (S = 1.5)

t-quark EW Couplings

$$F_0 = 0.685 \pm 0.020$$

$$F_- = 0.320 \pm 0.013$$

$$F_+ = 0.002 \pm 0.011$$

$$F_{V+A} < 0.29, \text{ CL} = 95\%$$

t DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	$\frac{p}{\text{MeV}/c}$
$t \rightarrow Wq(q = b, s, d)$			—
$t \rightarrow Wb$			—
$t \rightarrow \ell\nu_\ell \text{ anything}$	[c,d] (9.4±2.4) %		—
$t \rightarrow e\nu_e b$	(13.3±0.6) %		—
$t \rightarrow \mu\nu_\mu b$	(13.4±0.6) %		—
$t \rightarrow q\bar{q}b$	(66.5±1.4) %		—

$\Delta T = 1$ weak neutral current (T1) modes

$t \rightarrow Zq(q=u,c)$	T1 [e] < 5	$\times 10^{-4}$	95%	—
$t \rightarrow \ell^+ \bar{q}q' (q=d,s,b; q'=u,c)$	< 1.6	$\times 10^{-3}$	95%	—

b' (4th Generation) Quark, Searches for

Mass $m > 190 \text{ GeV}$, CL = 95% ($p\bar{p}$, quasi-stable b')

Mass $m > 755 \text{ GeV}$, CL = 95% (pp , neutral-current decays)

Mass $m > 675 \text{ GeV}$, CL = 95% (pp , charged-current decays)

Mass $m > 46.0 \text{ GeV}$, CL = 95% (e^+e^- , all decays)

t' (4th Generation) Quark, Searches for

$$\begin{aligned} m(t'(2/3)) &> 782 \text{ GeV, CL} = 95\% && \text{(neutral-current decays)} \\ m(t'(2/3)) &> 700 \text{ GeV, CL} = 95\% && \text{(charged-current decays)} \\ m(t'(5/3)) &> 800 \text{ GeV, CL} = 95\% \end{aligned}$$

Free Quark Searches

All searches since 1977 have had negative results.

NOTES

- [a] A discussion of the definition of the top quark mass in these measurements can be found in the review “The Top Quark.”
- [b] Based on published top mass measurements using data from Tevatron Run-I and Run-II and LHC at $\sqrt{s} = 7$ TeV. Including the most recent unpublished results from Tevatron Run-II, the Tevatron Electroweak Working Group reports a top mass of 173.2 ± 0.9 GeV. See the note “The Top Quark” in the Quark Particle Listings of this *Review*.
- [c] ℓ means e or μ decay mode, not the sum over them.
- [d] Assumes lepton universality and W -decay acceptance.
- [e] This limit is for $\Gamma(t \rightarrow Zq)/\Gamma(t \rightarrow Wb)$.