

$J/\psi(1S)$

$$J^G(J^{PC}) = 0^-(1^{--})$$

 $J/\psi(1S)$ MASS

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3096.900±0.006 OUR AVERAGE				
3096.900±0.002±0.006		¹ ANASHIN 15	KEDR	$e^+e^- \rightarrow \text{hadrons}$
3096.89 ±0.09	502	² ARTAMONOV 00	OLYA	$e^+e^- \rightarrow \text{hadrons}$
3096.91 ±0.03 ±0.01		³ ARMSTRONG 93B	E760	$\bar{p}p \rightarrow e^+e^-$
3096.95 ±0.1 ±0.3	193	BAGLIN 87	SPEC	$\bar{p}p \rightarrow e^+e^-X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
3096.66 ±0.19 ±0.02	6.1k	⁴ AAIJ 15BI	LHCB	$pp \rightarrow J/\psi X$
3096.917±0.010±0.007		AULCHENKO 03	KEDR	$e^+e^- \rightarrow \text{hadrons}$
3097.5 ±0.3		GRIBUSHIN 96	FMPS	515 $\pi^- \text{Be} \rightarrow 2\mu X$
3098.4 ±2.0	38k	LEMOIGNE 82	GOLI	185 $\pi^- \text{Be} \rightarrow \gamma\mu^+\mu^- A$
3096.93 ±0.09	502	⁵ ZHOLENTZ 80	REDE	e^+e^-
3097.0 ±1		⁶ BRANDELIK 79C	DASP	e^+e^-

¹ Supersedes AULCHENKO 03.² Reanalysis of ZHOLENTZ 80 using new electron mass (COHEN 87) and radiative corrections (KURAEV 85).³ Mass central value and systematic error recalculated by us according to Eq. (16) in ARMSTRONG 93B, using the value for the $\psi(2S)$ mass from AULCHENKO 03.⁴ From a sample of $\eta_c(1S)$ and J/ψ produced in b -hadron decays. Systematic uncertainties not estimated.⁵ Superseded by ARTAMONOV 00.⁶ From a simultaneous fit to e^+e^- , $\mu^+\mu^-$ and hadronic channels assuming $\Gamma(e^+e^-) = \Gamma(\mu^+\mu^-)$. **$J/\psi(1S)$ WIDTH**

<u>VALUE (keV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
92.9± 2.8 OUR AVERAGE Error includes scale factor of 1.1.				
96.1± 3.2	13k	¹ ADAMS 06A	CLEO	$e^+e^- \rightarrow \mu^+\mu^-\gamma$
84.4± 8.9		BAI 95B	BES	e^+e^-
91 ±11 ±6		² ARMSTRONG 93B	E760	$\bar{p}p \rightarrow e^+e^-$
85.5 ^{+6.1} _{-5.8}		³ HSUEH 92	RVUE	See Υ mini-review

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

94.1± 2.7 ⁴ ANASHIN 10 KEDR 3.097 $e^+e^- \rightarrow e^+e^-, \mu^+\mu^-$ 93.7± 3.5 7.8k ¹ AUBERT 04 BABR $e^+e^- \rightarrow \mu^+\mu^-\gamma$ ¹ Calculated by us from the reported values of $\Gamma(e^+e^-) \times B(\mu^+\mu^-)$ using $B(e^+e^-) = (5.94 \pm 0.06)\%$ and $B(\mu^+\mu^-) = (5.93 \pm 0.06)\%$.² The initial-state radiation correction reevaluated by ANDREOTTI 07 in its Ref. [4].³ Using data from COFFMAN 92, BALDINI-CELIO 75, BOYARSKI 75, ESPOSITO 75B, BRANDELIK 79C.⁴ Assuming $\Gamma(e^+e^-) = \Gamma(\mu^+\mu^-)$ and using $\Gamma(e^+e^-)/\Gamma_{\text{total}} = (5.94 \pm 0.06)\%$.

$J/\psi(1S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 hadrons	(87.7 \pm 0.5) %	
Γ_2 virtual $\gamma \rightarrow$ hadrons	(13.50 \pm 0.30) %	
Γ_3 $g g g$	(64.1 \pm 1.0) %	
Γ_4 $\gamma g g$	(8.8 \pm 1.1) %	
Γ_5 $e^+ e^-$	(5.971 \pm 0.032) %	
Γ_6 $e^+ e^- \gamma$	[a] (8.8 \pm 1.4) $\times 10^{-3}$	
Γ_7 $\mu^+ \mu^-$	(5.961 \pm 0.033) %	

Decays involving hadronic resonances

Γ_8 $\rho \pi$	(1.69 \pm 0.15) %	S=2.4
Γ_9 $\rho^0 \pi^0$	(5.6 \pm 0.7) $\times 10^{-3}$	
Γ_{10} $\rho(770)^\mp K^\pm K_S^0$	(1.9 \pm 0.4) $\times 10^{-3}$	
Γ_{11} $\rho(1450) \pi$		
Γ_{12} $\rho(1450) \pi \rightarrow \pi^+ \pi^- \pi^0$	(2.3 \pm 0.7) $\times 10^{-3}$	
Γ_{13} $\rho(1450)^\pm \pi^\mp \rightarrow K_S^0 K^\pm \pi^\mp$	(3.5 \pm 0.6) $\times 10^{-4}$	
Γ_{14} $\rho(1450)^0 \pi^0 \rightarrow K^+ K^- \pi^0$	(2.0 \pm 0.5) $\times 10^{-4}$	
Γ_{15} $\rho(1450) \eta'(958) \rightarrow$ $\pi^+ \pi^- \eta'(958)$	(3.3 \pm 0.7) $\times 10^{-6}$	
Γ_{16} $\rho(1700) \pi$		
Γ_{17} $\rho(1700) \pi \rightarrow \pi^+ \pi^- \pi^0$	(1.7 \pm 1.1) $\times 10^{-4}$	
Γ_{18} $\rho(2150) \pi$		
Γ_{19} $\rho(2150) \pi \rightarrow \pi^+ \pi^- \pi^0$	(8 \pm 40) $\times 10^{-6}$	
Γ_{20} $\rho_3(1690) \pi \rightarrow \pi^+ \pi^- \pi^0$		
Γ_{21} $a_2(1320) \rho$	(1.09 \pm 0.22) %	
Γ_{22} $\omega \pi^+ \pi^+ \pi^- \pi^-$	(8.5 \pm 3.4) $\times 10^{-3}$	
Γ_{23} $\omega \pi^+ \pi^- \pi^0$	(4.0 \pm 0.7) $\times 10^{-3}$	
Γ_{24} $\omega \pi^+ \pi^-$	(8.6 \pm 0.7) $\times 10^{-3}$	S=1.1
Γ_{25} $\omega f_2(1270)$	(4.3 \pm 0.6) $\times 10^{-3}$	
Γ_{26} $K^*(892)^0 \bar{K}^*(892)^0$	(2.3 \pm 0.6) $\times 10^{-4}$	
Γ_{27} $K^*(892)^\pm K^*(892)^\mp$	(1.00 $^{+0.22}_{-0.40}$) $\times 10^{-3}$	
Γ_{28} $K^*(892)^\pm K^*(700)^\mp$	(1.1 $^{+1.0}_{-0.6}$) $\times 10^{-3}$	
Γ_{29} $K_S^0 \pi^- K^*(892)^+ + \text{c.c.}$	(2.0 \pm 0.5) $\times 10^{-3}$	
Γ_{30} $K_S^0 \pi^- K^*(892)^+ + \text{c.c.} \rightarrow$ $K_S^0 K_S^0 \pi^+ \pi^-$	(6.7 \pm 2.2) $\times 10^{-4}$	
Γ_{31} $\eta K^*(892)^0 \bar{K}^*(892)^0$	(1.15 \pm 0.26) $\times 10^{-3}$	
Γ_{32} $K^*(1410) \bar{K} + \text{c.c.}$		
Γ_{33} $K^*(1410) \bar{K} + \text{c.c.} \rightarrow$ $K^\pm K^\mp \pi^0$	(4.9 \pm 2.8) $\times 10^{-5}$	
Γ_{34} $K^*(1410) \bar{K} + \text{c.c.} \rightarrow$ $K_S^0 K^\pm \pi^\mp$	(8 \pm 6) $\times 10^{-5}$	

Γ_{35}	$K_2^*(1430)\bar{K} + \text{c.c.}$		
Γ_{36}	$K_2^*(1430)\bar{K} + \text{c.c.} \rightarrow$ $K^\pm K^\mp \pi^0$	$(7.5 \pm 3.5) \times 10^{-5}$	
Γ_{37}	$K_2^*(1430)\bar{K} + \text{c.c.} \rightarrow$ $K_S^0 K^\pm \pi^\mp$	$(4.0 \pm 1.0) \times 10^{-4}$	
Γ_{38}	$K^*(892)^0 \bar{K}_2^*(1430)^0 + \text{c.c.}$	$(4.66 \pm 0.31) \times 10^{-3}$	
Γ_{39}	$K^*(892)^+ K_2^*(1430)^- + \text{c.c.}$	$(3.4 \pm 2.9) \times 10^{-3}$	
Γ_{40}	$K^*(892)^+ K_2^*(1430)^- + \text{c.c.} \rightarrow$ $K^*(892)^+ K_S^0 \pi^- + \text{c.c.}$	$(4 \pm 4) \times 10^{-4}$	
Γ_{41}	$K^*(892)^0 \bar{K}_2(1770)^0 + \text{c.c.} \rightarrow$ $K^*(892)^0 K^- \pi^+ + \text{c.c.}$	$(6.9 \pm 0.9) \times 10^{-4}$	
Γ_{42}	$\omega K^*(892)\bar{K} + \text{c.c.}$	$(6.1 \pm 0.9) \times 10^{-3}$	
Γ_{43}	$\bar{K} K^*(892) + \text{c.c.}$		
Γ_{44}	$\bar{K} K^*(892) + \text{c.c.} \rightarrow$ $K_S^0 K^\pm \pi^\mp$	$(5.1 \pm 0.5) \times 10^{-3}$	
Γ_{45}	$K^+ K^*(892)^- + \text{c.c.}$	$(5.12 \pm 0.30) \times 10^{-3}$	
Γ_{46}	$K^+ K^*(892)^- + \text{c.c.} \rightarrow$ $K^+ K^- \pi^0$	$(1.97 \pm 0.20) \times 10^{-3}$	
Γ_{47}	$K^+ K^*(892)^- + \text{c.c.} \rightarrow$ $K^0 K^\pm \pi^\mp + \text{c.c.}$	$(3.0 \pm 0.4) \times 10^{-3}$	
Γ_{48}	$K^0 \bar{K}^*(892)^0 + \text{c.c.}$	$(4.39 \pm 0.31) \times 10^{-3}$	
Γ_{49}	$K^0 \bar{K}^*(892)^0 + \text{c.c.} \rightarrow$ $K^0 K^\pm \pi^\mp + \text{c.c.}$	$(3.2 \pm 0.4) \times 10^{-3}$	
Γ_{50}	$K_1(1400)^\pm K^\mp$	$(3.8 \pm 1.4) \times 10^{-3}$	
Γ_{51}	$\bar{K}^*(892)^0 K^+ \pi^- + \text{c.c.}$	seen	
Γ_{52}	$K^*(892)^\pm K^\mp \pi^0$	$(4.1 \pm 1.3) \times 10^{-3}$	
Γ_{53}	$K^*(892)^0 K_S^0 \pi^0$	$(6 \pm 4) \times 10^{-4}$	
Γ_{54}	$\omega \pi^0 \pi^0$	$(3.4 \pm 0.8) \times 10^{-3}$	
Γ_{55}	$b_1(1235)^\pm \pi^\mp$	[b] $(3.0 \pm 0.5) \times 10^{-3}$	
Γ_{56}	$\omega K^\pm K_S^0 \pi^\mp$	[b] $(3.4 \pm 0.5) \times 10^{-3}$	
Γ_{57}	$b_1(1235)^0 \pi^0$	$(2.3 \pm 0.6) \times 10^{-3}$	
Γ_{58}	$\eta K^\pm K_S^0 \pi^\mp$	[b] $(2.2 \pm 0.4) \times 10^{-3}$	
Γ_{59}	$\phi K^*(892)\bar{K} + \text{c.c.}$	$(2.18 \pm 0.23) \times 10^{-3}$	
Γ_{60}	$\omega K \bar{K}$	$(1.70 \pm 0.32) \times 10^{-3}$	
Γ_{61}	$\omega f_0(1710) \rightarrow \omega K \bar{K}$	$(4.8 \pm 1.1) \times 10^{-4}$	
Γ_{62}	$\phi 2(\pi^+ \pi^-)$	$(1.66 \pm 0.23) \times 10^{-3}$	
Γ_{63}	$\Delta(1232)^{++} \bar{p} \pi^-$	$(1.6 \pm 0.5) \times 10^{-3}$	
Γ_{64}	$\omega \eta$	$(1.74 \pm 0.20) \times 10^{-3}$	S=1.6
Γ_{65}	$\phi K \bar{K}$	$(1.77 \pm 0.16) \times 10^{-3}$	S=1.3
Γ_{66}	$\phi K_S^0 K_S^0$	$(5.9 \pm 1.5) \times 10^{-4}$	
Γ_{67}	$\phi f_0(1710) \rightarrow \phi K \bar{K}$	$(3.6 \pm 0.6) \times 10^{-4}$	
Γ_{68}	$\phi K^+ K^-$	$(8.3 \pm 1.2) \times 10^{-4}$	
Γ_{69}	$\phi f_2(1270)$	$(3.2 \pm 0.6) \times 10^{-4}$	

Γ_{70}	$\Delta(1232)^{++}\bar{\Delta}(1232)^{--}$		$(1.10 \pm 0.29) \times 10^{-3}$	
Γ_{71}	$\Sigma(1385)^-\bar{\Sigma}(1385)^+$ (or c.c.)	[b]	$(1.16 \pm 0.05) \times 10^{-3}$	
Γ_{72}	$\Sigma(1385)^0\bar{\Sigma}(1385)^0$		$(1.07 \pm 0.08) \times 10^{-3}$	
Γ_{73}	$K^+K^-f_2'(1525)$		$(1.04 \pm 0.35) \times 10^{-3}$	
Γ_{74}	$\phi f_2'(1525)$		$(8 \pm 4) \times 10^{-4}$	S=2.7
Γ_{75}	$\phi\pi^+\pi^-$		$(8.7 \pm 0.9) \times 10^{-4}$	S=1.4
Γ_{76}	$\phi\pi^0\pi^0$		$(5.0 \pm 1.0) \times 10^{-4}$	
Γ_{77}	$\phi K^\pm K_S^0\pi^\mp$	[b]	$(7.2 \pm 0.8) \times 10^{-4}$	
Γ_{78}	$\omega f_1(1420)$		$(6.8 \pm 2.4) \times 10^{-4}$	
Γ_{79}	$\phi\eta$		$(7.5 \pm 0.8) \times 10^{-4}$	S=1.5
Γ_{80}	$\Xi^0\Xi^0$		$(1.17 \pm 0.04) \times 10^{-3}$	
Γ_{81}	$\Xi(1530)^-\Xi^+$		$(5.9 \pm 1.5) \times 10^{-4}$	
Γ_{82}	$\rho K^-\bar{\Sigma}(1385)^0$		$(5.1 \pm 3.2) \times 10^{-4}$	
Γ_{83}	$\omega\pi^0$		$(4.5 \pm 0.5) \times 10^{-4}$	S=1.4
Γ_{84}	$\omega\pi^0 \rightarrow \pi^+\pi^-\pi^0$		$(1.7 \pm 0.8) \times 10^{-5}$	
Γ_{85}	$\phi\eta'(958)$		$(4.6 \pm 0.5) \times 10^{-4}$	S=2.2
Γ_{86}	$\phi f_0(980)$		$(3.2 \pm 0.9) \times 10^{-4}$	S=1.9
Γ_{87}	$\phi f_0(980) \rightarrow \phi\pi^+\pi^-$		$(2.59 \pm 0.34) \times 10^{-4}$	
Γ_{88}	$\phi f_0(980) \rightarrow \phi\pi^0\pi^0$		$(1.8 \pm 0.5) \times 10^{-4}$	
Γ_{89}	$\phi\pi^0 f_0(980) \rightarrow \phi\pi^0\pi^+\pi^-$		$(4.5 \pm 1.0) \times 10^{-6}$	
Γ_{90}	$\phi\pi^0 f_0(980) \rightarrow \phi\pi^0\rho^0\pi^0$		$(1.7 \pm 0.6) \times 10^{-6}$	
Γ_{91}	$\eta\phi f_0(980) \rightarrow \eta\phi\pi^+\pi^-$		$(3.2 \pm 1.0) \times 10^{-4}$	
Γ_{92}	$\phi a_0(980)^0 \rightarrow \phi\eta\pi^0$		$(5 \pm 4) \times 10^{-6}$	
Γ_{93}	$\Xi(1530)^0\Xi^0$		$(3.2 \pm 1.4) \times 10^{-4}$	
Γ_{94}	$\Sigma(1385)^-\bar{\Sigma}^+$ (or c.c.)	[b]	$(3.1 \pm 0.5) \times 10^{-4}$	
Γ_{95}	$\phi f_1(1285)$		$(2.6 \pm 0.5) \times 10^{-4}$	
Γ_{96}	$\phi f_1(1285) \rightarrow \phi\pi^0 f_0(980) \rightarrow$ $\phi\pi^0\pi^+\pi^-$		$(9.4 \pm 2.8) \times 10^{-7}$	
Γ_{97}	$\phi f_1(1285) \rightarrow \phi\pi^0 f_0(980) \rightarrow$ $\phi\pi^0\pi^0\pi^0$		$(2.1 \pm 2.2) \times 10^{-7}$	
Γ_{98}	$\eta\pi^+\pi^-$		$(4.0 \pm 1.7) \times 10^{-4}$	
Γ_{99}	$\eta\rho$		$(1.93 \pm 0.23) \times 10^{-4}$	
Γ_{100}	$\omega\eta'(958)$		$(1.89 \pm 0.18) \times 10^{-4}$	
Γ_{101}	$\omega f_0(980)$		$(1.4 \pm 0.5) \times 10^{-4}$	
Γ_{102}	$\rho\eta'(958)$		$(8.1 \pm 0.8) \times 10^{-5}$	S=1.6
Γ_{103}	$a_2(1320)^\pm\pi^\mp$	[b]	$< 4.3 \times 10^{-3}$	CL=90%
Γ_{104}	$K\bar{K}_2^*(1430)+$ c.c.		$< 4.0 \times 10^{-3}$	CL=90%
Γ_{105}	$K_1(1270)^\pm K^\mp$		$< 3.0 \times 10^{-3}$	CL=90%
Γ_{106}	$K_S^0\pi^-K_2^*(1430)^++$ c.c.		$(3.6 \pm 1.8) \times 10^{-3}$	
Γ_{107}	$K_S^0\pi^-K_2^*(1430)^++$ c.c. \rightarrow $K_S^0K_S^0\pi^+\pi^-$		$(4.5 \pm 2.2) \times 10^{-4}$	
Γ_{108}	$K_2^*(1430)^0\bar{K}_2^*(1430)^0$		$< 2.9 \times 10^{-3}$	CL=90%
Γ_{109}	$\phi\pi^0$		3×10^{-6} or 1×10^{-7}	

Γ_{110}	$\phi\eta(1405) \rightarrow \phi\eta\pi^+\pi^-$	$(2.0 \pm 1.0) \times 10^{-5}$	
Γ_{111}	$\omega f'_2(1525)$	< 2.2	$\times 10^{-4}$ CL=90%
Γ_{112}	$\omega X(1835) \rightarrow \omega p\bar{p}$	< 3.9	$\times 10^{-6}$ CL=95%
Γ_{113}	$\phi X(1835) \rightarrow \phi p\bar{p}$	< 2.1	$\times 10^{-7}$ CL=90%
Γ_{114}	$\phi X(1835) \rightarrow \phi\eta\pi^+\pi^-$	< 2.8	$\times 10^{-4}$ CL=90%
Γ_{115}	$\phi X(1870) \rightarrow \phi\eta\pi^+\pi^-$	< 6.13	$\times 10^{-5}$ CL=90%
Γ_{116}	$\eta\phi(2170) \rightarrow \eta\phi f_0(980) \rightarrow$ $\eta\phi\pi^+\pi^-$	$(1.2 \pm 0.4) \times 10^{-4}$	
Γ_{117}	$\eta\phi(2170) \rightarrow$ $\eta K^*(892)^0 \bar{K}^*(892)^0$	< 2.52	$\times 10^{-4}$ CL=90%
Γ_{118}	$\Sigma(1385)^0 \bar{\Lambda} + \text{c.c.}$	< 8.2	$\times 10^{-6}$ CL=90%
Γ_{119}	$\Delta(1232)^+ \bar{p}$	< 1	$\times 10^{-4}$ CL=90%
Γ_{120}	$\Lambda(1520) \bar{\Lambda} + \text{c.c.} \rightarrow \gamma \Lambda \bar{\Lambda}$	< 4.1	$\times 10^{-6}$ CL=90%
Γ_{121}	$\Theta(1540) \bar{\Theta}(1540) \rightarrow$ $K_S^0 p K^- \bar{n} + \text{c.c.}$	< 1.1	$\times 10^{-5}$ CL=90%
Γ_{122}	$\Theta(1540) K^- \bar{n} \rightarrow K_S^0 p K^- \bar{n}$	< 2.1	$\times 10^{-5}$ CL=90%
Γ_{123}	$\Theta(1540) K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n$	< 1.6	$\times 10^{-5}$ CL=90%
Γ_{124}	$\bar{\Theta}(1540) K^+ n \rightarrow K_S^0 \bar{p} K^+ n$	< 5.6	$\times 10^{-5}$ CL=90%
Γ_{125}	$\bar{\Theta}(1540) K_S^0 p \rightarrow K_S^0 p K^- \bar{n}$	< 1.1	$\times 10^{-5}$ CL=90%
Γ_{126}	$\Sigma^0 \bar{\Lambda}$	< 9	$\times 10^{-5}$ CL=90%

Decays into stable hadrons

Γ_{127}	$2(\pi^+\pi^-)\pi^0$	$(4.1 \pm 0.5) \%$	S=2.4
Γ_{128}	$3(\pi^+\pi^-)\pi^0$	$(2.9 \pm 0.6) \%$	
Γ_{129}	$\pi^+\pi^-\pi^0$	$(2.11 \pm 0.07) \%$	S=1.5
Γ_{130}	$\pi^+\pi^-\pi^0 K^+ K^-$	$(1.79 \pm 0.29) \%$	S=2.2
Γ_{131}	$4(\pi^+\pi^-)\pi^0$	$(9.0 \pm 3.0) \times 10^{-3}$	
Γ_{132}	$\pi^+\pi^- K^+ K^-$	$(6.84 \pm 0.32) \times 10^{-3}$	
Γ_{133}	$\pi^+\pi^- K_S^0 K_L^0$	$(3.8 \pm 0.6) \times 10^{-3}$	
Γ_{134}	$\pi^+\pi^- K_S^0 K_S^0$	$(1.68 \pm 0.19) \times 10^{-3}$	
Γ_{135}	$\pi^\pm \pi^0 K^\mp K_S^0$	$(5.7 \pm 0.5) \times 10^{-3}$	
Γ_{136}	$K^+ K^- K_S^0 K_S^0$	$(4.1 \pm 0.8) \times 10^{-4}$	
Γ_{137}	$\pi^+\pi^- K^+ K^- \eta$	$(1.84 \pm 0.28) \times 10^{-3}$	
Γ_{138}	$\pi^0 \pi^0 K^+ K^-$	$(2.12 \pm 0.23) \times 10^{-3}$	
Γ_{139}	$\pi^0 \pi^0 K_S^0 K_L^0$	$(1.9 \pm 0.4) \times 10^{-3}$	
Γ_{140}	$K \bar{K} \pi$	$(6.1 \pm 1.0) \times 10^{-3}$	
Γ_{141}	$K^+ K^- \pi^0$	$(2.14 \pm 0.24) \times 10^{-3}$	
Γ_{142}	$K_S^0 K^\pm \pi^\mp$	$(5.6 \pm 0.5) \times 10^{-3}$	
Γ_{143}	$K_S^0 K_L^0 \pi^0$	$(2.06 \pm 0.27) \times 10^{-3}$	
Γ_{144}	$K^*(892)^0 \bar{K}^0 + \text{c.c.} \rightarrow$ $K_S^0 K_L^0 \pi^0$	$(1.21 \pm 0.18) \times 10^{-3}$	
Γ_{145}	$K_2^*(1430)^0 \bar{K}^0 + \text{c.c.} \rightarrow$ $K_S^0 K_L^0 \pi^0$	$(4.3 \pm 1.3) \times 10^{-4}$	

Γ_{146}	$K_S^0 K_L^0 \eta$	$(1.44 \pm 0.34) \times 10^{-3}$	
Γ_{147}	$2(\pi^+ \pi^-)$	$(3.57 \pm 0.30) \times 10^{-3}$	
Γ_{148}	$3(\pi^+ \pi^-)$	$(4.3 \pm 0.4) \times 10^{-3}$	
Γ_{149}	$2(\pi^+ \pi^- \pi^0)$	$(1.62 \pm 0.21) \%$	
Γ_{150}	$2(\pi^+ \pi^-) \eta$	$(2.29 \pm 0.24) \times 10^{-3}$	
Γ_{151}	$3(\pi^+ \pi^-) \eta$	$(7.2 \pm 1.5) \times 10^{-4}$	
Γ_{152}	$\rho \bar{\rho}$	$(2.121 \pm 0.029) \times 10^{-3}$	
Γ_{153}	$\rho \bar{\rho} \pi^0$	$(1.19 \pm 0.08) \times 10^{-3}$	S=1.1
Γ_{154}	$\rho \bar{\rho} \pi^+ \pi^-$	$(6.0 \pm 0.5) \times 10^{-3}$	S=1.3
Γ_{155}	$\rho \bar{\rho} \pi^+ \pi^- \pi^0$	[c] $(2.3 \pm 0.9) \times 10^{-3}$	S=1.9
Γ_{156}	$\rho \bar{\rho} \eta$	$(2.00 \pm 0.12) \times 10^{-3}$	
Γ_{157}	$\rho \bar{\rho} \rho$	< 3.1 $\times 10^{-4}$	CL=90%
Γ_{158}	$\rho \bar{\rho} \omega$	$(9.8 \pm 1.0) \times 10^{-4}$	S=1.3
Γ_{159}	$\rho \bar{\rho} \eta'(958)$	$(2.1 \pm 0.4) \times 10^{-4}$	
Γ_{160}	$\rho \bar{\rho} a_0(980) \rightarrow \rho \bar{\rho} \pi^0 \eta$	$(6.8 \pm 1.8) \times 10^{-5}$	
Γ_{161}	$\rho \bar{\rho} \phi$	$(5.19 \pm 0.33) \times 10^{-5}$	
Γ_{162}	$n \bar{n}$	$(2.09 \pm 0.16) \times 10^{-3}$	
Γ_{163}	$n \bar{n} \pi^+ \pi^-$	$(4 \pm 4) \times 10^{-3}$	
Γ_{164}	$\Sigma^+ \bar{\Sigma}^-$	$(1.50 \pm 0.24) \times 10^{-3}$	
Γ_{165}	$\Sigma^0 \bar{\Sigma}^0$	$(1.172 \pm 0.031) \times 10^{-3}$	S=1.4
Γ_{166}	$2(\pi^+ \pi^-) K^+ K^-$	$(4.7 \pm 0.7) \times 10^{-3}$	S=1.3
Γ_{167}	$\rho \bar{n} \pi^-$	$(2.12 \pm 0.09) \times 10^{-3}$	
Γ_{168}	$n N(1440)$	seen	
Γ_{169}	$n N(1520)$	seen	
Γ_{170}	$n N(1535)$	seen	
Γ_{171}	$\Xi^- \bar{\Xi}^+$	$(9.7 \pm 0.8) \times 10^{-4}$	S=1.4
Γ_{172}	$\Lambda \bar{\Lambda}$	$(1.89 \pm 0.08) \times 10^{-3}$	S=2.5
Γ_{173}	$\Lambda \bar{\Sigma}^- \pi^+$ (or c.c.)	[b] $(8.3 \pm 0.7) \times 10^{-4}$	S=1.2
Γ_{174}	$\rho K^- \bar{\Lambda}$	$(8.9 \pm 1.6) \times 10^{-4}$	
Γ_{175}	$2(K^+ K^-)$	$(7.4 \pm 0.7) \times 10^{-4}$	
Γ_{176}	$\rho K^- \bar{\Sigma}^0$	$(2.9 \pm 0.8) \times 10^{-4}$	
Γ_{177}	$K^+ K^-$	$(2.86 \pm 0.21) \times 10^{-4}$	
Γ_{178}	$K_S^0 K_L^0$	$(1.95 \pm 0.11) \times 10^{-4}$	S=2.4
Γ_{179}	$\Lambda \bar{\Lambda} \pi^+ \pi^-$	$(4.3 \pm 1.0) \times 10^{-3}$	
Γ_{180}	$\Lambda \bar{\Lambda} \eta$	$(1.62 \pm 0.17) \times 10^{-4}$	
Γ_{181}	$\Lambda \bar{\Lambda} \pi^0$	$(3.8 \pm 0.4) \times 10^{-5}$	
Γ_{182}	$\bar{\Lambda} n K_S^0 + \text{c.c.}$	$(6.5 \pm 1.1) \times 10^{-4}$	
Γ_{183}	$\pi^+ \pi^-$	$(1.47 \pm 0.14) \times 10^{-4}$	
Γ_{184}	$\Lambda \bar{\Sigma}^+ + \text{c.c.}$	$(2.83 \pm 0.23) \times 10^{-5}$	
Γ_{185}	$K_S^0 K_S^0$	< 1.4 $\times 10^{-8}$	CL=95%

Radiative decays

Γ_{186}	3γ	$(1.16 \pm 0.22) \times 10^{-5}$	
Γ_{187}	4γ	$< 9 \times 10^{-6}$	CL=90%
Γ_{188}	5γ	$< 1.5 \times 10^{-5}$	CL=90%
Γ_{189}	$\gamma\pi^0\pi^0$	$(1.15 \pm 0.05) \times 10^{-3}$	
Γ_{190}	$\gamma\eta\pi^0$	$(2.14 \pm 0.31) \times 10^{-5}$	
Γ_{191}	$\gamma a_0(980)^0 \rightarrow \gamma\eta\pi^0$	$< 2.5 \times 10^{-6}$	CL=95%
Γ_{192}	$\gamma a_2(1320)^0 \rightarrow \gamma\eta\pi^0$	$< 6.6 \times 10^{-6}$	CL=95%
Γ_{193}	$\gamma\eta_c(1S)$	$(1.7 \pm 0.4)\%$	S=1.5
Γ_{194}	$\gamma\eta_c(1S) \rightarrow 3\gamma$	$(3.8 \begin{smallmatrix} +1.3 \\ -1.0 \end{smallmatrix}) \times 10^{-6}$	S=1.1
Γ_{195}	$\gamma\pi^+\pi^-2\pi^0$	$(8.3 \pm 3.1) \times 10^{-3}$	
Γ_{196}	$\gamma\eta\pi\pi$	$(6.1 \pm 1.0) \times 10^{-3}$	
Γ_{197}	$\gamma\eta_2(1870) \rightarrow \gamma\eta\pi^+\pi^-$	$(6.2 \pm 2.4) \times 10^{-4}$	
Γ_{198}	$\gamma\eta(1405/1475) \rightarrow \gamma K\bar{K}\pi$	[d] $(2.8 \pm 0.6) \times 10^{-3}$	S=1.6
Γ_{199}	$\gamma\eta(1405/1475) \rightarrow \gamma\gamma\rho^0$	$(7.8 \pm 2.0) \times 10^{-5}$	S=1.8
Γ_{200}	$\gamma\eta(1405/1475) \rightarrow \gamma\eta\pi^+\pi^-$	$(3.0 \pm 0.5) \times 10^{-4}$	
Γ_{201}	$\gamma\eta(1405/1475) \rightarrow \gamma\gamma\phi$	$< 8.2 \times 10^{-5}$	CL=95%
Γ_{202}	$\gamma\rho\rho$	$(4.5 \pm 0.8) \times 10^{-3}$	
Γ_{203}	$\gamma\rho\omega$	$< 5.4 \times 10^{-4}$	CL=90%
Γ_{204}	$\gamma\rho\phi$	$< 8.8 \times 10^{-5}$	CL=90%
Γ_{205}	$\gamma\eta'(958)$	$(5.13 \pm 0.17) \times 10^{-3}$	S=1.3
Γ_{206}	$\gamma 2\pi^+ 2\pi^-$	$(2.8 \pm 0.5) \times 10^{-3}$	S=1.9
Γ_{207}	$\gamma f_2(1270) f_2(1270)$	$(9.5 \pm 1.7) \times 10^{-4}$	
Γ_{208}	$\gamma f_2(1270) f_2(1270)$ (non resonant)	$(8.2 \pm 1.9) \times 10^{-4}$	
Γ_{209}	$\gamma K^+ K^- \pi^+ \pi^-$	$(2.1 \pm 0.6) \times 10^{-3}$	
Γ_{210}	$\gamma f_4(2050)$	$(2.7 \pm 0.7) \times 10^{-3}$	
Γ_{211}	$\gamma\omega\omega$	$(1.61 \pm 0.33) \times 10^{-3}$	
Γ_{212}	$\gamma\eta(1405/1475) \rightarrow \gamma\rho^0\rho^0$	$(1.7 \pm 0.4) \times 10^{-3}$	S=1.3
Γ_{213}	$\gamma f_2(1270)$	$(1.64 \pm 0.12) \times 10^{-3}$	S=1.3
Γ_{214}	$\gamma f_0(1370) \rightarrow \gamma K\bar{K}$	$(4.2 \pm 1.5) \times 10^{-4}$	
Γ_{215}	$\gamma f_0(1710) \rightarrow \gamma K\bar{K}$	$(1.00 \begin{smallmatrix} +0.11 \\ -0.09 \end{smallmatrix}) \times 10^{-3}$	S=1.5
Γ_{216}	$\gamma f_0(1710) \rightarrow \gamma\pi\pi$	$(3.8 \pm 0.5) \times 10^{-4}$	
Γ_{217}	$\gamma f_0(1710) \rightarrow \gamma\omega\omega$	$(3.1 \pm 1.0) \times 10^{-4}$	
Γ_{218}	$\gamma f_0(1710) \rightarrow \gamma\eta\eta$	$(2.4 \begin{smallmatrix} +1.2 \\ -0.7 \end{smallmatrix}) \times 10^{-4}$	
Γ_{219}	$\gamma\eta$	$(1.104 \pm 0.034) \times 10^{-3}$	
Γ_{220}	$\gamma f_1(1420) \rightarrow \gamma K\bar{K}\pi$	$(7.9 \pm 1.3) \times 10^{-4}$	
Γ_{221}	$\gamma f_1(1285)$	$(6.1 \pm 0.8) \times 10^{-4}$	
Γ_{222}	$\gamma f_1(1510) \rightarrow \gamma\eta\pi^+\pi^-$	$(4.5 \pm 1.2) \times 10^{-4}$	
Γ_{223}	$\gamma f_2'(1525)$	$(5.7 \begin{smallmatrix} +0.8 \\ -0.5 \end{smallmatrix}) \times 10^{-4}$	S=1.5
Γ_{224}	$\gamma f_2'(1525) \rightarrow \gamma\eta\eta$	$(3.4 \pm 1.4) \times 10^{-5}$	

Γ_{225}	$\gamma f_2(1640) \rightarrow \gamma \omega \omega$	$(2.8 \pm 1.8) \times 10^{-4}$	
Γ_{226}	$\gamma f_2(1910) \rightarrow \gamma \omega \omega$	$(2.0 \pm 1.4) \times 10^{-4}$	
Γ_{227}	$\gamma f_0(1800) \rightarrow \gamma \omega \phi$	$(2.5 \pm 0.6) \times 10^{-4}$	
Γ_{228}	$\gamma f_2(1810) \rightarrow \gamma \eta \eta$	$(5.4 \begin{smallmatrix} + 3.5 \\ - 2.4 \end{smallmatrix}) \times 10^{-5}$	
Γ_{229}	$\gamma f_2(1950) \rightarrow$ $\gamma K^*(892) \bar{K}^*(892)$	$(7.0 \pm 2.2) \times 10^{-4}$	
Γ_{230}	$\gamma K^*(892) \bar{K}^*(892)$	$(4.0 \pm 1.3) \times 10^{-3}$	
Γ_{231}	$\gamma \phi \phi$	$(4.0 \pm 1.2) \times 10^{-4}$	S=2.1
Γ_{232}	$\gamma p \bar{p}$	$(3.8 \pm 1.0) \times 10^{-4}$	
Γ_{233}	$\gamma \eta(2225)$	$(3.14 \begin{smallmatrix} + 0.50 \\ - 0.19 \end{smallmatrix}) \times 10^{-4}$	
Γ_{234}	$\gamma \eta(1760) \rightarrow \gamma \rho^0 \rho^0$	$(1.3 \pm 0.9) \times 10^{-4}$	
Γ_{235}	$\gamma \eta(1760) \rightarrow \gamma \omega \omega$	$(1.98 \pm 0.33) \times 10^{-3}$	
Γ_{236}	$\gamma X(1835) \rightarrow \gamma \pi^+ \pi^- \eta'$	$(2.77 \begin{smallmatrix} + 0.34 \\ - 0.40 \end{smallmatrix}) \times 10^{-4}$	S=1.1
Γ_{237}	$\gamma X(1835) \rightarrow \gamma p \bar{p}$	$(7.7 \begin{smallmatrix} + 1.5 \\ - 0.9 \end{smallmatrix}) \times 10^{-5}$	
Γ_{238}	$\gamma X(1835) \rightarrow \gamma K_S^0 K_S^0 \eta$	$(3.3 \begin{smallmatrix} + 2.0 \\ - 1.3 \end{smallmatrix}) \times 10^{-5}$	
Γ_{239}	$\gamma X(1840) \rightarrow \gamma 3(\pi^+ \pi^-)$	$(2.4 \begin{smallmatrix} + 0.7 \\ - 0.8 \end{smallmatrix}) \times 10^{-5}$	
Γ_{240}	$\gamma (K \bar{K} \pi) [J^{PC} = 0^- +]$	$(7 \pm 4) \times 10^{-4}$	S=2.1
Γ_{241}	$\gamma \pi^0$	$(3.49 \begin{smallmatrix} + 0.33 \\ - 0.30 \end{smallmatrix}) \times 10^{-5}$	
Γ_{242}	$\gamma p \bar{p} \pi^+ \pi^-$	$< 7.9 \times 10^{-4}$	CL=90%
Γ_{243}	$\gamma \Lambda \bar{\Lambda}$	$< 1.3 \times 10^{-4}$	CL=90%
Γ_{244}	$\gamma f_0(2100) \rightarrow \gamma \eta \eta$	$(1.13 \begin{smallmatrix} + 0.60 \\ - 0.30 \end{smallmatrix}) \times 10^{-4}$	
Γ_{245}	$\gamma f_0(2100) \rightarrow \gamma \pi \pi$	$(6.2 \pm 1.0) \times 10^{-4}$	
Γ_{246}	$\gamma f_0(2200)$		
Γ_{247}	$\gamma f_0(2200) \rightarrow \gamma K \bar{K}$	$(5.9 \pm 1.3) \times 10^{-4}$	
Γ_{248}	$\gamma f_J(2220)$		
Γ_{249}	$\gamma f_J(2220) \rightarrow \gamma \pi \pi$	$< 3.9 \times 10^{-5}$	CL=90%
Γ_{250}	$\gamma f_J(2220) \rightarrow \gamma K \bar{K}$	$< 4.1 \times 10^{-5}$	CL=90%
Γ_{251}	$\gamma f_J(2220) \rightarrow \gamma p \bar{p}$	$(1.5 \pm 0.8) \times 10^{-5}$	
Γ_{252}	$\gamma f_2(2340) \rightarrow \gamma \eta \eta$	$(5.6 \begin{smallmatrix} + 2.4 \\ - 2.2 \end{smallmatrix}) \times 10^{-5}$	
Γ_{253}	$\gamma f_0(1500) \rightarrow \gamma \pi \pi$	$(1.09 \pm 0.24) \times 10^{-4}$	
Γ_{254}	$\gamma f_0(1500) \rightarrow \gamma \eta \eta$	$(1.7 \begin{smallmatrix} + 0.6 \\ - 1.4 \end{smallmatrix}) \times 10^{-5}$	
Γ_{255}	$\gamma A \rightarrow \gamma \text{invisible}$	$[e] < 6.3 \times 10^{-6}$	CL=90%
Γ_{256}	$\gamma A^0 \rightarrow \gamma \mu^+ \mu^-$	$[f] < 5 \times 10^{-6}$	CL=90%

Dalitz decays

Γ_{257}	$\pi^0 e^+ e^-$	$(7.6 \pm 1.4) \times 10^{-7}$
Γ_{258}	$\eta e^+ e^-$	$(1.16 \pm 0.09) \times 10^{-5}$
Γ_{259}	$\eta'(958) e^+ e^-$	$(5.81 \pm 0.35) \times 10^{-5}$

Weak decays

Γ_{260}	$D^- e^+ \nu_e + \text{c.c.}$	< 1.2	$\times 10^{-5}$	CL=90%
Γ_{261}	$\overline{D}^0 e^+ e^- + \text{c.c.}$	< 8.5	$\times 10^{-8}$	CL=90%
Γ_{262}	$D_s^- e^+ \nu_e + \text{c.c.}$	< 1.3	$\times 10^{-6}$	CL=90%
Γ_{263}	$D_s^{*-} e^+ \nu_e + \text{c.c.}$	< 1.8	$\times 10^{-6}$	CL=90%
Γ_{264}	$D^- \pi^+ + \text{c.c.}$	< 7.5	$\times 10^{-5}$	CL=90%
Γ_{265}	$\overline{D}^0 \overline{K}^0 + \text{c.c.}$	< 1.7	$\times 10^{-4}$	CL=90%
Γ_{266}	$\overline{D}^0 \overline{K}^{*0} + \text{c.c.}$	< 2.5	$\times 10^{-6}$	CL=90%
Γ_{267}	$D_s^- \pi^+ + \text{c.c.}$	< 1.3	$\times 10^{-4}$	CL=90%
Γ_{268}	$D_s^- \rho^+ + \text{c.c.}$	< 1.3	$\times 10^{-5}$	CL=90%

**Charge conjugation (C), Parity (P),
Lepton Family number (LF) violating modes**

Γ_{269}	$\gamma\gamma$	C	< 2.7	$\times 10^{-7}$	CL=90%
Γ_{270}	$\gamma\phi$	C	< 1.4	$\times 10^{-6}$	CL=90%
Γ_{271}	$e^\pm \mu^\mp$	LF	< 1.6	$\times 10^{-7}$	CL=90%
Γ_{272}	$e^\pm \tau^\mp$	LF	< 8.3	$\times 10^{-6}$	CL=90%
Γ_{273}	$\mu^\pm \tau^\mp$	LF	< 2.0	$\times 10^{-6}$	CL=90%

Other decays

Γ_{274}	invisible	< 7	$\times 10^{-4}$	CL=90%
----------------	-----------	-----	------------------	--------

[a] For $E_\gamma > 100$ MeV.

[b] The value is for the sum of the charge states or particle/antiparticle states indicated.

[c] Includes $p\bar{p}\pi^+\pi^-\gamma$ and excludes $p\bar{p}\eta$, $p\bar{p}\omega$, $p\bar{p}\eta'$.

[d] See the "Note on the $\eta(1405)$ " in the $\eta(1405)$ Particle Listings.

[e] For a narrow state A with mass less than 960 MeV.

[f] For a narrow scalar or pseudoscalar A^0 with mass 0.21–3.0 GeV.

 $J/\psi(1S)$ PARTIAL WIDTHS

$\Gamma(\text{hadrons})$					Γ_1
VALUE (keV)	DOCUMENT ID	TECN	COMMENT		
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
74.1 ± 8.1	BAI	95B	BES	$e^+ e^-$	
59 ± 24	BALDINI-...	75	FRAG	$e^+ e^-$	
59 ± 14	BOYARSKI	75	MRK1	$e^+ e^-$	
50 ± 25	ESPOSITO	75B	FRAM	$e^+ e^-$	

$\Gamma(e^+e^-)$ Γ_5

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
5.55±0.14±0.02 OUR EVALUATION				
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
5.58±0.05±0.08		¹ ABLIKIM	16Q BES3	3.773 $e^+e^- \rightarrow \mu^+\mu^-\gamma$
5.71±0.16	13k	² ADAMS	06A CLEO	$e^+e^- \rightarrow \mu^+\mu^-\gamma$
5.57±0.19	7.8k	² AUBERT	04 BABR	$e^+e^- \rightarrow \mu^+\mu^-\gamma$
5.14±0.39		BAI	95B BES	e^+e^-
5.36 ^{+0.29} _{-0.28}		³ HSUEH	92 RVUE	See \mathcal{T} mini-review
4.72±0.35		ALEXANDER	89 RVUE	See \mathcal{T} mini-review
4.4 ±0.6		³ BRANDELIK	79C DASP	e^+e^-
4.6 ±0.8		⁴ BALDINI-...	75 FRAG	e^+e^-
4.8 ±0.6		BOYARSKI	75 MRK1	e^+e^-
4.6 ±1.0		ESPOSITO	75B FRAM	e^+e^-

¹ Using $B(J/\psi \rightarrow \mu^+\mu^-) = (5.973 \pm 0.007 \pm 0.037)\%$ from ABLIKIM 13R.² Calculated by us from the reported values of $\Gamma(e^+e^-) \times B(\mu^+\mu^-)$ using $B(\mu^+\mu^-) = (5.93 \pm 0.06)\%$.³ From a simultaneous fit to e^+e^- , $\mu^+\mu^-$, and hadronic channels assuming $\Gamma(e^+e^-) = \Gamma(\mu^+\mu^-)$.⁴ Assuming equal partial widths for e^+e^- and $\mu^+\mu^-$. $\Gamma(\mu^+\mu^-)$ Γ_7

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
5.13±0.52	BAI	95B BES	e^+e^-
4.8 ±0.6	BOYARSKI	75 MRK1	e^+e^-
5 ±1	ESPOSITO	75B FRAM	e^+e^-

 $\Gamma(\gamma\gamma)$ Γ_{269}

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<5.4	90	BRANDELIK	79C DASP	e^+e^-

 $J/\psi(1S) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

This combination of a partial width with the partial width into e^+e^- and with the total width is obtained from the integrated cross section into channel i in the e^+e^- annihilation.

 $\Gamma(\text{hadrons}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_1\Gamma_5/\Gamma$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
4 ±0.8	¹ BALDINI-...	75 FRAG	e^+e^-
3.9±0.8	¹ ESPOSITO	75B FRAM	e^+e^-

¹ Data redundant with branching ratios or partial widths above.

$\Gamma(e^+e^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_5\Gamma_5/\Gamma$

VALUE (eV)		DOCUMENT ID	TECN	COMMENT
332.3 ± 6.4 ± 4.8		ANASHIN	10	KEDR 3.097 $e^+e^- \rightarrow e^+e^-$
• • •	We do not use the following data for averages, fits, limits, etc. • • •			
350 ± 20		BRANDELIK	79C	DASP e^+e^-
320 ± 70		¹ BALDINI-...	75	FRAG e^+e^-
340 ± 90		¹ ESPOSITO	75B	FRAM e^+e^-
360 ± 100		¹ FORD	75	SPEC e^+e^-

¹Data redundant with branching ratios or partial widths above.

$\Gamma(\mu^+\mu^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_7\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
333 ± 4	OUR AVERAGE			
333.4 ± 2.5 ± 4.4		ABLIKIM	16Q	BES3 3.773 $e^+e^- \rightarrow \mu^+\mu^-\gamma$
331.8 ± 5.2 ± 6.3		ANASHIN	10	KEDR 3.097 $e^+e^- \rightarrow \mu^+\mu^-$
338.4 ± 5.8 ± 7.1	13k	ADAMS	06A	CLEO $e^+e^- \rightarrow \mu^+\mu^-\gamma$
330.1 ± 7.7 ± 7.3	7.8k	AUBERT	04	BABR $e^+e^- \rightarrow \mu^+\mu^-\gamma$
• • •	We do not use the following data for averages, fits, limits, etc. • • •			
510 ± 90		DASP	75	DASP e^+e^-
380 ± 50		¹ ESPOSITO	75B	FRAM e^+e^-

¹Data redundant with branching ratios or partial widths above.

$\Gamma(\rho(770)^\mp K^\pm K_S^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{10}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
10.4 ± 1.0 ± 1.9	130	LEES	17D	BABR $e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$

$\Gamma(\omega\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{23}\Gamma_5/\Gamma$

VALUE (10 ⁻² keV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.2 ± 0.3 ± 0.2	170	AUBERT	06D	BABR 10.6 $e^+e^- \rightarrow \omega\pi^+\pi^-\pi^0\gamma$

$\Gamma(\omega\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{24}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
53.6 ± 5.0 ± 0.4	788	¹ AUBERT	07AU	BABR 10.6 $e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$

¹AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow \omega\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = 47.8 \pm 3.1 \pm 3.2$ eV which we divide by our best value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^*(892)^0 \bar{K}^*(892)^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{26}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
1.28 ± 0.34 ± 0.07	47 ± 12	¹ LEES	12F	BABR 10.6 $e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.28 ± 0.40 ± 0.11	25 ± 8	^{1,2} AUBERT	07AK	BABR 10.6 $e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$
--------------------	--------	-----------------------	------	---

¹Dividing by $(2/3)^2$ to take twice into account that $B(K^{*0} \rightarrow K^+\pi^-) = 2/3 B(K^{*0} \rightarrow K\pi)$.

²Superseded by LEES 12F.

$$\Gamma(K^*(892)^0 \bar{K}_2^*(1430)^0 + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{38} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

25.8 ± 1.4 ± 0.6	710	1,2,3 LEES	12F BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$
-------------------------	-----	------------	----------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

33 ± 4 ± 1	317	2,4 AUBERT	07AK BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$
------------	-----	------------	-----------	---

¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow K^*(892)^0 \bar{K}_2^*(1430)^0 + \text{c.c.}) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(K_2^*(1430) \rightarrow K\pi)] = 12.89 \pm 0.54 \pm 0.41$ eV which we divide by our best value $B(K_2^*(1430) \rightarrow K\pi) = (49.9 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Dividing by 2/3 to take into account that $B(K^*0 \rightarrow K^+ \pi^-) = 2/3 B(K^*0 \rightarrow K\pi)$.

³ The $K_2^*(1430)$ cannot be distinguished from the $K_0^*(1430)$.

⁴ Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow K^*(892)^0 \bar{K}_2^*(1430)^0 + \text{c.c.}) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(K_2^*(1430) \rightarrow K\pi)] = 16.4 \pm 1.1 \pm 1.4$ eV which we divide by our best value $B(K_2^*(1430) \rightarrow K\pi) = (49.9 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(K^*(892)^0 \bar{K}_2(1770)^0 + \text{c.c.} \rightarrow K^*(892)^0 K^- \pi^+ + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{41} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

3.8 ± 0.4 ± 0.3	110 ± 14	¹ AUBERT	07AK BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$
------------------------	----------	---------------------	-----------	---

¹ Dividing by 2/3 to take into account that $B(K^*0 \rightarrow K^+ \pi^-) = 2/3$.

$$\Gamma(K^+ K^*(892)^- + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{45} \Gamma_5 / \Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

29.0 ± 1.7 ± 1.3	AUBERT	08S BABR	10.6 $e^+ e^- \rightarrow K^+ K^*(892)^- \gamma$
-------------------------	--------	----------	--

$$\Gamma(K^+ K^*(892)^- + \text{c.c.} \rightarrow K^+ K^- \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{46} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

10.96 ± 0.85 ± 0.70	155	AUBERT	08S BABR	10.6 $e^+ e^- \rightarrow K^+ K^- \pi^0 \gamma$
----------------------------	-----	--------	----------	---

$$\Gamma(K^+ K^*(892)^- + \text{c.c.} \rightarrow K^0 K^\pm \pi^\mp + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{47} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

16.76 ± 1.70 ± 1.00	89	AUBERT	08S BABR	10.6 $e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \gamma$
----------------------------	----	--------	----------	---

$$\Gamma(K^0 \bar{K}^*(892)^0 + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{48} \Gamma_5 / \Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

26.6 ± 2.5 ± 1.5	AUBERT	08S BABR	10.6 $e^+ e^- \rightarrow K^0 \bar{K}^*(892)^0 \gamma$
-------------------------	--------	----------	--

$$\Gamma(K^0 \bar{K}^*(892)^0 + \text{c.c.} \rightarrow K^0 K^\pm \pi^\mp + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{49} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

17.70 ± 1.70 ± 1.00	94	AUBERT	08S BABR	10.6 $e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \gamma$
----------------------------	----	--------	----------	---

$$\Gamma(K^*(892)^\pm K^\mp \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{52} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
22.8 ± 2.8 ± 6.8	80	¹ LEES	17D BABR	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$

¹ Dividing by 1/4 to account for $B(K^*(892)^\pm \rightarrow K_S^0 \pi^\pm) = 1/4$.

$$\Gamma(K^*(892)^0 K_S^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{53} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
3.60 ± 0.75 ± 2.25	34	¹ LEES	17D BABR	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$

¹ Dividing by 2/3 to account for $B(K^*(892)^0 \rightarrow K^+ \pi^-) = 2/3$.

$$\Gamma(\omega K \bar{K}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{60} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
3.70 ± 1.98 ± 0.03	24	¹ AUBERT	07AU BABR	$10.6 e^+ e^- \rightarrow \omega K^+ K^- \gamma$

¹ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow \omega K \bar{K}) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0)] = 3.3 \pm 1.3 \pm 1.2$ eV which we divide by our best value $B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\phi 2(\pi^+ \pi^-)) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{62} \Gamma_5 / \Gamma$$

VALUE (10^{-2} keV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.95 ± 0.19 ± 0.01	35	¹ AUBERT	06D BABR	$10.6 e^+ e^- \rightarrow \phi 2(\pi^+ \pi^-) \gamma$

¹ AUBERT 06D reports $[\Gamma(J/\psi(1S) \rightarrow \phi 2(\pi^+ \pi^-)) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = (0.47 \pm 0.09 \pm 0.03) \times 10^{-2}$ keV which we divide by our best value $B(\phi(1020) \rightarrow K^+ K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\phi K^+ K^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{68} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
4.59 ± 0.62 ± 0.05	163	¹ LEES	12F BABR	$10.6 e^+ e^- \rightarrow K^+ K^- K^+ K^- \gamma$

¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi K^+ K^-) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 2.26 \pm 0.26 \pm 0.16$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+ K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\phi f_2(1270)) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{69} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
1.79 ± 0.32^{+0.02}_{-0.06}	61 ± 10	^{1,2,3} LEES	12F BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.08 ± 0.73 ^{+0.04} _{-0.14}	44 ± 7	^{2,4} AUBERT	07AK BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$
---	--------	-----------------------	-----------	---

¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_2(1270)) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi \pi)] = 1.51 \pm 0.25 \pm 0.10$ eV which we divide by our best value $B(f_2(1270) \rightarrow \pi \pi) = (84.2^{+2.9}_{-0.9}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Using $B(\phi \rightarrow K^+ K^-) = (48.9 \pm 0.5)\%$.

³ Using $\pi^+ \pi^-$ invariant mass between 1.1 and 1.5 GeV. May include other sources such as $f_0(1370)$.

⁴ Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_2(1270)) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = 3.44 \pm 0.55 \pm 0.28$ eV which we divide by our best value $B(f_2(1270) \rightarrow \pi\pi) = (84.2^{+2.9}_{-0.9}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi\pi^+\pi^-) \times \Gamma(e^+e^-) / \Gamma_{\text{total}}$ $\Gamma_{75}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
4.47±0.35 OUR AVERAGE				
4.45±0.49±0.05	181	¹ LEES	12F BABR	10.6 $e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
4.50±0.48±0.05	254 ± 23	² SHEN	09 BELL	10.6 $e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
5.3 ± 0.7 ± 0.1	103	³ AUBERT, BE 06D	BABR	10.6 $e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$

¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-) / \Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 2.19 \pm 0.23 \pm 0.07$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² SHEN 09 reports $4.50 \pm 0.41 \pm 0.26$ eV from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \phi\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-) / \Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)]$ assuming $B(\phi(1020) \rightarrow K^+K^-) = (49.2 \pm 0.6) \times 10^{-2}$, which we rescale to our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Superseded by LEES 12F. AUBERT, BE 06D reports $[\Gamma(J/\psi(1S) \rightarrow \phi\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-) / \Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 2.61 \pm 0.30 \pm 0.18$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi\pi^0\pi^0) \times \Gamma(e^+e^-) / \Gamma_{\text{total}}$ $\Gamma_{76}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.76±0.57±0.03				
3.13±0.88±0.03	23	² AUBERT, BE 06D	BABR	10.6 $e^+e^- \rightarrow K^+K^-\pi^0\pi^0\gamma$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi\pi^0\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-) / \Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 1.36 \pm 0.27 \pm 0.07$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Superseded by LEES 12F. AUBERT, BE 06D reports $[\Gamma(J/\psi(1S) \rightarrow \phi\pi^0\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-) / \Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 1.54 \pm 0.40 \pm 0.16$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\phi\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{79}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
6.1±2.7±0.4	6	¹ AUBERT	07AU BABR	10.6 e ⁺ e ⁻ → φηγ
¹ AUBERT 07AU quotes $\Gamma_{ee}^{J/\psi} \cdot B(J/\psi \rightarrow \phi\eta) \cdot B(\phi \rightarrow K^+K^-) \cdot B(\eta \rightarrow 3\pi) = 0.84 \pm 0.37 \pm 0.05$ eV.				

$$\Gamma(\phi f_0(980) \rightarrow \phi\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{87}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
1.44±0.19 OUR AVERAGE				
1.40±0.25±0.02	57 ± 9	¹ LEES 12F	BABR	10.6 e ⁺ e ⁻ → π ⁺ π ⁻ K ⁺ K ⁻ γ
1.48±0.27±0.09	60 ± 11	² SHEN 09	BELL	10.6 e ⁺ e ⁻ → K ⁺ K ⁻ π ⁺ π ⁻ γ
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1.02±0.24±0.01	20 ± 5	³ AUBERT 07AK	BABR	10.6 e ⁺ e ⁻ → π ⁺ π ⁻ K ⁺ K ⁻ γ
¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)]/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 0.69 \pm 0.11 \pm 0.05$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				
² Multiplied by 2/3 to take into account the φπ ⁺ π ⁻ mode only. Using $B(\phi \rightarrow K^+K^-) = (49.2 \pm 0.6)\%$.				
³ Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)]/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 0.50 \pm 0.11 \pm 0.04$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

$$\Gamma(\phi f_0(980) \rightarrow \phi\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{88}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.98±0.26±0.01	16 ± 4	¹ LEES 12F	BABR	10.6 e ⁺ e ⁻ → π ⁰ π ⁰ K ⁺ K ⁻ γ
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.95±0.40±0.01	7.0 ± 2.8	² AUBERT 07AK	BABR	10.6 e ⁺ e ⁻ → π ⁰ π ⁰ K ⁺ K ⁻ γ
¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi\pi^0\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)]/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 0.48 \pm 0.12 \pm 0.05$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				
² Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi\pi^0\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)]/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 0.47 \pm 0.19 \pm 0.05$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

$$\Gamma(\eta\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{98}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.23±0.97±0.03	9	¹ AUBERT	07AU BABR	10.6 e ⁺ e ⁻ → ηπ ⁺ π ⁻ γ
¹ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow \eta\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)]/\Gamma_{\text{total}}] \times [B(\eta \rightarrow \pi^+\pi^-\pi^0)] = 0.51 \pm 0.22 \pm 0.03$ eV which we divide by our best value $B(\eta \rightarrow \pi^+\pi^-\pi^0) = (22.92 \pm 0.28) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

$$\Gamma(2(\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{127}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
303±5±18	4990	AUBERT	07AU BABR	10.6 e ⁺ e ⁻ → 2(π ⁺ π ⁻)π ⁰ γ

$$\Gamma(\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{129}\Gamma_5/\Gamma$$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
0.122±0.005±0.008	AUBERT,B	04N BABR	10.6 e ⁺ e ⁻ → π ⁺ π ⁻ π ⁰ γ

$$\Gamma(\pi^+\pi^-\pi^0K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{130}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
107.0±4.3±6.4	768	AUBERT	07AU BABR	10.6 e ⁺ e ⁻ → K ⁺ K ⁻ π ⁺ π ⁻ π ⁰ γ

$$\Gamma(\pi^+\pi^-K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{132}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
37.94±0.81±1.10	3.1k	LEES	12F BABR	10.6 e ⁺ e ⁻ → π ⁺ π ⁻ K ⁺ K ⁻ γ

• • • We do not use the following data for averages, fits, limits, etc. • • •

36.3 ±1.3 ±2.1 1.5k ¹ AUBERT 07AK BABR 10.6 e⁺e⁻ → π⁺π⁻K⁺K⁻γ

33.6 ±2.7 ±2.7 233 ² AUBERT 05D BABR 10.6 e⁺e⁻ → K⁺K⁻π⁺π⁻γ

¹ Superseded by LEES 12F.

² Superseded by AUBERT 07AK.

$$\Gamma(\pi^\pm\pi^0K^\mp K_S^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{135}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
31.7±1.9±1.8	393	LEES	17D BABR	e ⁺ e ⁻ → K _S ⁰ K [±] π [∓] π ⁰ γ

$$\Gamma(\pi^+\pi^-K^+K^-\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{137}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
25.9±3.9±0.1	73	¹ AUBERT	07AU BABR	10.6 e ⁺ e ⁻ → K ⁺ K ⁻ π ⁺ π ⁻ ηγ

¹ AUBERT 07AU reports [Γ(J/ψ(1S) → π⁺π⁻K⁺K⁻η) × Γ(J/ψ(1S) → e⁺e⁻)/Γ_{total}] × [B(η → 2γ)] = 10.2 ± 1.3 ± 0.8 eV which we divide by our best value B(η → 2γ) = (39.41 ± 0.20) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\pi^0\pi^0K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{138}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
11.75±0.81±0.90	388	LEES	12F BABR	10.6 e ⁺ e ⁻ → π ⁰ π ⁰ K ⁺ K ⁻ γ

• • • We do not use the following data for averages, fits, limits, etc. • • •

13.6 ±1.1 ±1.3 203 ¹ AUBERT 07AK BABR 10.6 e⁺e⁻ → π⁰π⁰K⁺K⁻γ

¹ Superseded by LEES 12F.

$$\Gamma(\pi^+\pi^-K_S^0K_L^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{133}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
20.8±2.3±2.1	248	LEES	14H BABR	e ⁺ e ⁻ → π ⁺ π ⁻ K _S ⁰ K _L ⁰ γ

$$\Gamma(\pi^+\pi^-K_S^0K_S^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{134}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
9.3±0.9±0.5	133	LEES	14H BABR	e ⁺ e ⁻ → π ⁺ π ⁻ K _S ⁰ K _S ⁰ γ

$$\Gamma(K^+ K^- K_S^0 K_S^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{136} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.3±0.4±0.1	29	LEES	14H BABR	$e^+ e^- \rightarrow K_S^0 K_S^0 K^+ K^- \gamma$

$$\Gamma(K_S^0 \pi^- K^*(892)^+ + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{29} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
11.0±2.8 OUR AVERAGE				
9.2±1.2±3.2	64	¹ LEES	17D BABR	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$
14.8±4.8±1.2	53	² LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$

¹ Dividing by 1/2 to take into account $B(K^*(892)^\pm \rightarrow K^\pm \pi^\mp) = 1/2$.

² Dividing by 1/4 to take into account $B(K^*(892) \rightarrow K_S^0 \pi) = 1/4$.

$$\Gamma(K_S^0 \pi^- K^*(892)^+ + \text{c.c.} \rightarrow K_S^0 K_S^0 \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{30} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
3.7±1.2±0.3	53	LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$

$$\Gamma(K_S^0 \pi^- K_2^*(1430)^+ + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{106} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
20.1±9.8±0.5	35	^{1,2} LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$

¹ Dividing by 1/4 to take into account $B(K^*(1430) \rightarrow K_S^0 \pi) = 1/4$ $B(K^*(1430) \rightarrow K \pi)$.

² LEES 14H reports $[\Gamma(J/\psi(1S) \rightarrow K_S^0 \pi^- K_2^*(1430)^+ + \text{c.c.}) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(K_2^*(1430) \rightarrow K \pi)] = 10.0 \pm 4.8 \pm 0.8$ eV which we divide by our best value $B(K_2^*(1430) \rightarrow K \pi) = (49.9 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(K_S^0 \pi^- K_2^*(1430)^+ + \text{c.c.} \rightarrow K_S^0 K_S^0 \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{107} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.5±1.2±0.2	35	LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$

$$\Gamma(K^*(892)^\pm K^*(892)^\mp) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{27} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.80±0.48±0.32	1±5	¹ LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$

¹ Dividing by $(1/4)^2$ to take twice into account $B(K^*(892) \rightarrow K_S^0 \pi) = 1/4$.

$$\Gamma(K^*(892)^+ K_2^*(1430)^- + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{39} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
18.6±16.1±0.4	8±8	^{1,2} LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$

¹ Dividing by $(1/4)^2$ to take into account $B(K^*(892) \rightarrow K_S^0 \pi) = 1/4$ and $B(K^*(1430) \rightarrow K_S^0 \pi) = 1/4$ $B(K^*(1430) \rightarrow K \pi)$.

² LEES 14H reports $[\Gamma(J/\psi(1S) \rightarrow K^*(892)^+ K_2^*(1430)^- + \text{c.c.}) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(K_2^*(1430) \rightarrow K \pi)] = 9.28 \pm 8.0 \pm 0.32$ eV which we divide by our best value $B(K_2^*(1430) \rightarrow K \pi) = (49.9 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\frac{\Gamma(K^*(892)^+ K_2^*(1430)^- + \text{c.c.} \rightarrow K^*(892)^+ K_S^0 \pi^- + \text{c.c.}) \times \Gamma(e^+ e^-)}{\Gamma_{\text{total}} \Gamma_{40} \Gamma_5 / \Gamma}$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.32±2.00±0.08	8 ± 8	¹ LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$

¹ Dividing by 1/4 to take into account $B(K^*(892) \rightarrow K_S^0 \pi) = 1/4$.

$$\frac{\Gamma(\phi K_S^0 K_S^0) \times \Gamma(e^+ e^-)}{\Gamma_{\text{total}} \Gamma_{66} \Gamma_5 / \Gamma}$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
3.25±0.84±0.03	29	¹ LEES	14H BABR	$e^+ e^- \rightarrow K_S^0 K_S^0 K^+ K^- \gamma$

¹ LEES 14H reports $[\Gamma(J/\psi(1S) \rightarrow \phi K_S^0 K_S^0) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 1.6 \pm 0.4 \pm 0.1$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+ K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\frac{\Gamma(\phi f_2'(1525)) \times \Gamma(e^+ e^-)}{\Gamma_{\text{total}} \Gamma_{74} \Gamma_5 / \Gamma}$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
8.1±3.2±0.2	11	^{1,2} LEES	14H BABR	$e^+ e^- \rightarrow K_S^0 K_S^0 K^+ K^- \gamma$

¹ Dividing by 1/4 to take into account $B(f_2'(1525) \rightarrow K_S^0 K_S^0) = 1/4$ $B(f_2'(1525) \rightarrow K \bar{K})$ and using $B(\phi \rightarrow K^+ K^-) = (48.9 \pm 0.5)\%$.

² LEES 14H reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_2'(1525)) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(f_2'(1525) \rightarrow K \bar{K})] = 7.2 \pm 2.8 \pm 0.3$ eV which we divide by our best value $B(f_2'(1525) \rightarrow K \bar{K}) = (88.7 \pm 2.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\frac{\Gamma(K^+ K^- f_2'(1525)) \times \Gamma(e^+ e^-)}{\Gamma_{\text{total}} \Gamma_{73} \Gamma_5 / \Gamma}$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
5.8±1.9±0.1	16	^{1,2} LEES	14H BABR	$e^+ e^- \rightarrow K_S^0 K_S^0 K^+ K^- \gamma$

¹ Dividing by 1/4 to take into account $B(f_2'(1525) \rightarrow K_S^0 K_S^0) = 1/4$ $B(f_2'(1525) \rightarrow K \bar{K})$.

² LEES 14H reports $[\Gamma(J/\psi(1S) \rightarrow K^+ K^- f_2'(1525)) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(f_2'(1525) \rightarrow K \bar{K})] = 5.12 \pm 1.68 \pm 0.20$ eV which we divide by our best value $B(f_2'(1525) \rightarrow K \bar{K}) = (88.7 \pm 2.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\frac{\Gamma(K_S^0 K_L^0 \eta) \times \Gamma(e^+ e^-)}{\Gamma_{\text{total}} \Gamma_{146} \Gamma_5 / \Gamma}$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
8.0±1.8±0.4	45	LEES	17A BABR	$e^+ e^- \rightarrow K_S^0 K_L^0 \eta \gamma$

$$\frac{\Gamma(\eta K^\pm K_S^0 \pi^\mp) \times \Gamma(e^+ e^-)}{\Gamma_{\text{total}} \Gamma_{58} \Gamma_5 / \Gamma}$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
7.3±1.4±0.4	44	LEES	17D BABR	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$

$$\Gamma(\overline{K}^*(892)^0 K^+ \pi^- + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{51} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
42.6 ± 4.8 ± 7.2	99	¹ LEES	17D BABR	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$

¹ Dividing by 1/6 to account for $B(K^*(892)^0 \rightarrow K_S^0 \pi^0) = 1/6$.

$$\Gamma(\pi^0 \pi^0 K_S^0 K_L^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{139} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
10.3 ± 2.3 ± 0.5	47	LEES	17A BABR	$e^+ e^- \rightarrow K_S^0 K_L^0 \pi^0 \pi^0 \gamma$

$$\Gamma(K_S^0 K_L^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{143} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
11.4 ± 1.3 ± 0.6	182	LEES	17A BABR	$e^+ e^- \rightarrow K_S^0 K_L^0 \pi^0 \gamma$

$$\Gamma(K^*(892)^0 \overline{K}^0 + \text{c.c.} \rightarrow K_S^0 K_L^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{144} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
6.7 ± 0.9 ± 0.4	106	LEES	17A BABR	$e^+ e^- \rightarrow K_S^0 K_L^0 \pi^0 \gamma$

$$\Gamma(K_2^*(1430)^0 \overline{K}^0 + \text{c.c.} \rightarrow K_S^0 K_L^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{145} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.4 ± 0.7 ± 0.1	37	LEES	17A BABR	$e^+ e^- \rightarrow K_S^0 K_L^0 \pi^0 \gamma$

$$\Gamma(2(\pi^+ \pi^-)) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{147} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
20.4 ± 0.9 ± 0.4		LEES	12E BABR	$10.6 e^+ e^- \rightarrow 2\pi^+ 2\pi^- \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

19.5 ± 1.4 ± 1.3	270	¹ AUBERT	05D BABR	$10.6 e^+ e^- \rightarrow 2(\pi^+ \pi^-) \gamma$
------------------	-----	---------------------	----------	--

¹ Superseded by LEES 12E.

$$\Gamma(3(\pi^+ \pi^-)) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{148} \Gamma_5 / \Gamma$$

VALUE (10 ⁻² keV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.37 ± 0.16 ± 0.14	496	AUBERT	06D BABR	$10.6 e^+ e^- \rightarrow 3(\pi^+ \pi^-) \gamma$

$$\Gamma(2(\pi^+ \pi^- \pi^0)) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{149} \Gamma_5 / \Gamma$$

VALUE (10 ⁻² keV)	EVTS	DOCUMENT ID	TECN	COMMENT
8.9 ± 0.5 ± 1.0	761	AUBERT	06D BABR	$10.6 e^+ e^- \rightarrow 2(\pi^+ \pi^- \pi^0) \gamma$

$$\Gamma(2(\pi^+ \pi^-) \eta) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{150} \Gamma_5 / \Gamma$$


VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
13.1 ± 2.4 ± 0.1	85	¹ AUBERT	07AU BABR	$10.6 e^+ e^- \rightarrow 2(\pi^+ \pi^-) \eta \gamma$

¹ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow 2(\pi^+ \pi^-) \eta) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] = 5.16 \pm 0.85 \pm 0.39$ eV which we divide by our best value $B(\eta \rightarrow 2\gamma) = (39.41 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(p\bar{p}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{152}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
11.9±0.6 OUR AVERAGE		Error includes scale factor of 1.8. See the ideogram below.		
11.3±0.4±0.3	821	¹ LEES	13O	BABR $e^+e^- \rightarrow p\bar{p}\gamma$
12.9±0.4±0.4	918	² LEES	13Y	BABR $e^+e^- \rightarrow p\bar{p}\gamma$
9.7±1.7		³ ARMSTRONG	93B	E760 $\bar{p}p \rightarrow e^+e^-$
12.0±0.6±0.5	438	⁴ AUBERT	06B	BABR $e^+e^- \rightarrow p\bar{p}\gamma$

- • • We do not use the following data for averages, fits, limits, etc. • • •
- ¹ ISR photon reconstructed in the detector
- ² ISR photon undetected
- ³ Using $\Gamma_{\text{total}} = 85.5^{+6.1}_{-5.8}$ MeV.
- ⁴ Superseded by LEES 130



$\Gamma(\Sigma^0\bar{\Sigma}^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{165}\Gamma_5/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
6.4±1.2±0.6	AUBERT	07BD	BABR 10.6 $e^+e^- \rightarrow \Sigma^0\bar{\Sigma}^0\gamma$

$\Gamma(2(\pi^+\pi^-)K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{166}\Gamma_5/\Gamma$

VALUE (10 ⁻² keV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.75±0.23±0.17	205	AUBERT	06D	BABR 10.6 $e^+e^- \rightarrow K^+K^-2(\pi^+\pi^-)\gamma$

$\Gamma(\Lambda\bar{\Lambda}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{172}\Gamma_5/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
10.7±0.9±0.7	AUBERT	07BD	BABR 10.6 $e^+e^- \rightarrow \Lambda\bar{\Lambda}\gamma$

$\Gamma(2(K^+ K^-)) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{175} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
4.00 ± 0.33 ± 0.29	287 ± 24	LEES	12F	BABR 10.6 e ⁺ e ⁻ → 2(K ⁺ K ⁻)γ
• • • We do not use the following data for averages, fits, limits, etc. • • •				
4.11 ± 0.39 ± 0.30	156 ± 15	¹ AUBERT	07AK	BABR 10.6 e ⁺ e ⁻ → 2(K ⁺ K ⁻)γ
4.0 ± 0.7 ± 0.6	38	² AUBERT	05D	BABR 10.6 e ⁺ e ⁻ → 2(K ⁺ K ⁻)γ

¹ Superseded by LEES 12F.

² Superseded by AUBERT 07AK.

$\Gamma(K^+ K^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{177} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.78 ± 0.11 ± 0.05	462	¹ LEES	15J	BABR e ⁺ e ⁻ → K ⁺ K ⁻ γ
1.94 ± 0.11 ± 0.05	462	² LEES	15J	BABR e ⁺ e ⁻ → K ⁺ K ⁻ γ
1.42 ± 0.23 ± 0.08	51	³ LEES	13Q	BABR e ⁺ e ⁻ → K ⁺ K ⁻ γ

¹ sin φ > 0.

² sin φ < 0.

³ Interference with non-resonant K⁺K⁻ production not taken into account.

J/ψ(1S) BRANCHING RATIOS

For the first four branching ratios, see also the partial widths, and (partial widths) × Γ(e⁺e⁻)/Γ_{total} above.

$\Gamma(\text{hadrons}) / \Gamma_{\text{total}}$ Γ_1 / Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.877 ± 0.005 OUR AVERAGE			
0.878 ± 0.005	BAI	95B	BES e ⁺ e ⁻
0.86 ± 0.02	BOYARSKI	75	MRK1 e ⁺ e ⁻

$\Gamma(\text{virtual } \gamma \rightarrow \text{hadrons}) / \Gamma_{\text{total}}$ Γ_2 / Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.135 ± 0.003	^{1,2} SETH	04	RVUE e ⁺ e ⁻
0.17 ± 0.02	¹ BOYARSKI	75	MRK1 e ⁺ e ⁻

¹ Included in Γ(hadrons)/Γ_{total}.

² Using B(J/ψ → ℓ⁺ℓ⁻) = (5.90 ± 0.09)% from RPP-2002 and R = 2.28 ± 0.04 determined by a fit to data from BAI 00 and BAI 02C.

$\Gamma(g g g) / \Gamma_{\text{total}}$ Γ_3 / Γ

VALUE (units 10 ⁻²)	EVTS	DOCUMENT ID	TECN	COMMENT
64.1 ± 1.0	6 M	¹ BESSION	08	CLEO ψ(2S) → π ⁺ π ⁻ + hadrons

¹ Calculated using the value Γ(γγg)/Γ(ggg) = 0.137 ± 0.001 ± 0.016 ± 0.004 from BESSION 08 and the PDG 08 values of B(ℓ⁺ℓ⁻), B(virtual γ → hadrons), and B(γ η_C). The statistical error is negligible and the systematic error is partially correlated with that of Γ(γγg)/Γ_{total} measurement of BESSION 08.

$\Gamma(\gamma g g)/\Gamma_{\text{total}}$ Γ_4/Γ

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
8.79 ± 1.05	200 k	¹ BESSON	08 CLEO	$\psi(2S) \rightarrow \pi^+ \pi^- \gamma + \text{hadrons}$

¹ Calculated using the value $\Gamma(\gamma g g)/\Gamma(g g g) = 0.137 \pm 0.001 \pm 0.016 \pm 0.004$ from BESSON 08 and the value of $\Gamma(g g g)/\Gamma_{\text{total}}$. The statistical error is negligible and the systematic error is partially correlated with that of $\Gamma(g g g)/\Gamma_{\text{total}}$ measurement of BESSON 08.

 $\Gamma(\gamma g g)/\Gamma(g g g)$ Γ_4/Γ_3

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
13.7 ± 0.1 ± 0.7	6 M	BESSON	08 CLEO	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$

 $\Gamma(e^+ e^-)/\Gamma_{\text{total}}$ Γ_5/Γ

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
5.971 ± 0.032 OUR AVERAGE				
5.983 ± 0.007 ± 0.037	720k	ABLIKIM	13R BES3	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
5.945 ± 0.067 ± 0.042	15k	LI	05C CLEO	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
5.90 ± 0.05 ± 0.10		BAI	98D BES	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
6.09 ± 0.33		BAI	95B BES	$e^+ e^-$
5.92 ± 0.15 ± 0.20		COFFMAN	92 MRK3	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
6.9 ± 0.9		BOYARSKI	75 MRK1	$e^+ e^-$

 $\Gamma(e^+ e^- \gamma)/\Gamma_{\text{total}}$ Γ_6/Γ

<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
8.8 ± 1.3 ± 0.4	¹ ARMSTRONG	96 E760	$\bar{p} p \rightarrow e^+ e^- \gamma$

¹ For $E_\gamma > 100$ MeV.

 $\Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_7/Γ

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
5.961 ± 0.033 OUR AVERAGE				
5.973 ± 0.007 ± 0.038	770k	ABLIKIM	13R BES3	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
5.960 ± 0.065 ± 0.050	17k	LI	05C CLEO	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
5.84 ± 0.06 ± 0.10		BAI	98D BES	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
6.08 ± 0.33		BAI	95B BES	$e^+ e^-$
5.90 ± 0.15 ± 0.19		COFFMAN	92 MRK3	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
6.9 ± 0.9		BOYARSKI	75 MRK1	$e^+ e^-$

 $\Gamma(e^+ e^-)/\Gamma(\mu^+ \mu^-)$ Γ_5/Γ_7

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.0016 ± 0.0031 OUR AVERAGE			
1.0022 ± 0.0044 ± 0.0048	¹ AULCHENKO	14 KEDR	3.097 $e^+ e^- \rightarrow e^+ e^-, \mu^+ \mu^-$
1.0017 ± 0.0017 ± 0.0033	² ABLIKIM	13R BES3	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
1.002 ± 0.021 ± 0.013	³ ANASHIN	10 KEDR	3.097 $e^+ e^- \rightarrow e^+ e^-, \mu^+ \mu^-$
0.997 ± 0.012 ± 0.006	LI	05C CLEO	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1.011 ± 0.013 ± 0.016	BAI	98D BES	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
1.00 ± 0.07	BAI	95B BES	$e^+ e^-$
1.00 ± 0.05	BOYARSKI	75 MRK1	$e^+ e^-$

0.91 ±0.15 ESPOSITO 75B FRAM e^+e^-
 0.93 ±0.10 FORD 75 SPEC e^+e^-

¹ From 235.3k $J/\psi \rightarrow e^+e^-$ and 156.6k $J/\psi \rightarrow \mu^+\mu^-$ observed events.


² Not independent of the corresponding measurements of $\Gamma(e^+e^-)/\Gamma_{\text{total}}$ and $\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$.

³ Not independent of the corresponding measurements of $\Gamma(e^+e^-) \times \Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$ and $\Gamma(\mu^+\mu^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$.

HADRONIC DECAYS

$\Gamma(\rho\pi)/\Gamma_{\text{total}}$					Γ_8/Γ
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	
1.69 ±0.15	OUR AVERAGE	Error includes scale factor of 2.4. See the ideogram below.			
2.18 ±0.19		1,2 AUBERT,B	04N BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$	
2.184±0.005±0.201	220k	2,3 BAI	04H BES	$e^+e^- \rightarrow J/\psi \rightarrow \pi^+\pi^-\pi^0$	
2.091±0.021±0.116		2,4 BAI	04H BES	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$	
1.21 ±0.20		BAI	96D BES	$e^+e^- \rightarrow \rho\pi$	
1.42 ±0.01 ±0.19		COFFMAN	88 MRK3	e^+e^-	
1.3 ±0.3	150	FRANKLIN	83 MRK2	e^+e^-	
1.6 ±0.4	183	ALEXANDER	78 PLUT	e^+e^-	
1.33 ±0.21		BRANDELIK	78B DASP	e^+e^-	
1.0 ±0.2	543	BARTEL	76 CNTR	e^+e^-	
1.3 ±0.3	153	JEAN-MARIE	76 MRK1	e^+e^-	

WEIGHTED AVERAGE
 1.69±0.15 (Error scaled by 2.4)



¹ From the ratio of $\Gamma(e^+e^-)B(\pi^+\pi^-\pi^0)$ and $\Gamma(e^+e^-)B(\mu^+\mu^-)$ (AUBERT 04).

² Not independent of their $B(\pi^+\pi^-\pi^0)$.

³ From $J/\psi \rightarrow \pi^+\pi^-\pi^0$ events directly.

⁴ Obtained comparing the rates for $\pi^+\pi^-\pi^0$ and $\mu^+\mu^-$, using J/ψ events produced via $\psi(2S) \rightarrow \pi^+\pi^-J/\psi$ and with $B(J/\psi \rightarrow \mu^+\mu^-) = 5.88 \pm 0.10\%$.

$\Gamma(\rho^0\pi^0)/\Gamma(\rho\pi)$

Γ_9/Γ_8

VALUE	DOCUMENT ID	TECN	COMMENT
0.328±0.005±0.027	COFFMAN 88	MRK3	e^+e^-
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.35 ±0.08	ALEXANDER 78	PLUT	e^+e^-
0.32 ±0.08	BRANDELIK 78B	DASP	e^+e^-
0.39 ±0.11	BARTEL 76	CNTR	e^+e^-
0.37 ±0.09	JEAN-MARIE 76	MRK1	e^+e^-

$\Gamma(\rho\pi)/\Gamma(\pi^+\pi^-\pi^0)$

Γ_8/Γ_{129}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.142±0.011±0.026	20K	¹ LEES	17C	BABR $J/\psi \rightarrow \pi^+\pi^-\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.331±0.033	20K	² LEES	17C	BABR $J/\psi \rightarrow \pi^+\pi^-\pi^0$
¹ From a Dalitz plot analysis in an isobar model.				
² From a Dalitz plot analysis in a Veneziano model.				

$\Gamma(\rho(1450)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma(\pi^+\pi^-\pi^0)$

Γ_{12}/Γ_{129}

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
10.9 ±1.7 ±2.7	20K	¹ LEES	17C	BABR $J/\psi \rightarrow \pi^+\pi^-\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.80±0.27	20K	² LEES	17C	BABR $J/\psi \rightarrow \pi^+\pi^-\pi^0$
¹ From a Dalitz plot analysis in an isobar model.				
² From a Dalitz plot analysis in a Veneziano model.				

$\Gamma(\rho(1450)^0\pi^0 \rightarrow K^+K^-\pi^0)/\Gamma(K^+K^-\pi^0)$

Γ_{14}/Γ_{141}

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
9.3±2.0±0.6	2K	¹ LEES	17C	BABR $J/\psi \rightarrow K^+K^-\pi^0$
¹ From a Dalitz plot analysis in an isobar model.				

$\Gamma(\rho(1450)^\pm\pi^\mp \rightarrow K_S^0 K^\pm\pi^\mp)/\Gamma(K_S^0 K^\pm\pi^\mp)$

Γ_{13}/Γ_{142}

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
6.3±0.8±0.6	4K	¹ LEES	17C	BABR $J/\psi \rightarrow K_S^0 K^\pm\pi^\mp$
¹ From a Dalitz plot analysis in an isobar model.				

$\Gamma(\rho(1450)\eta'(958) \rightarrow \pi^+\pi^-\eta'(958))/\Gamma_{\text{total}}$

Γ_{15}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
3.28±0.55±0.44	119	¹ ABLIKIM	17AK	BES3 $J/\psi \rightarrow \pi^+\pi^-\eta'$
¹ From a partial wave analysis of the decay $J/\psi \rightarrow \pi^+\pi^-\eta'$.				

$\Gamma(\rho_3(1690)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma(\pi^+\pi^-\pi^0)$ Γ_{20}/Γ_{129}

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.0 ± 0.8	20K	¹ LEES	17C	BABR $J/\psi \rightarrow \pi^+\pi^-\pi^0$
---------------	-----	-------------------	-----	---

¹ From a Dalitz plot analysis in a Veneziano model.

$\Gamma(\rho(1700)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma(\pi^+\pi^-\pi^0)$ Γ_{17}/Γ_{129}

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$8 \pm 2 \pm 5$	20K	¹ LEES	17C	BABR $J/\psi \rightarrow \pi^+\pi^-\pi^0$
-----------------------------------	-----	-------------------	-----	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

22 ± 6	20K	² LEES	17C	BABR $J/\psi \rightarrow \pi^+\pi^-\pi^0$
------------	-----	-------------------	-----	---

¹ From a Dalitz plot analysis in an isobar model.

² From a Dalitz plot analysis in a Veneziano model.

$\Gamma(\rho(2150)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma(\pi^+\pi^-\pi^0)$ Γ_{19}/Γ_{129}

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$4 \pm 1 \pm 20$	20K	¹ LEES	17C	BABR $J/\psi \rightarrow \pi^+\pi^-\pi^0$
------------------------------------	-----	-------------------	-----	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

600 ± 250	20K	² LEES	17C	BABR $J/\psi \rightarrow \pi^+\pi^-\pi^0$
---------------	-----	-------------------	-----	---

¹ From a Dalitz plot analysis in an isobar model.

² From a Dalitz plot analysis in a Veneziano model.

$\Gamma(a_2(1320)\rho)/\Gamma_{\text{total}}$ Γ_{21}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

10.9 ± 2.2 OUR AVERAGE

$11.7 \pm 0.7 \pm 2.5$	7584	AUGUSTIN	89	DM2 $J/\psi \rightarrow \rho^0 \rho^\pm \pi^\mp$
------------------------	------	----------	----	--

8.4 ± 4.5	36	VANNUCCI	77	MRK1 $e^+e^- \rightarrow 2(\pi^+\pi^-)\pi^0$
---------------	----	----------	----	--

$\Gamma(\omega\pi^+\pi^+\pi^-\pi^-)/\Gamma_{\text{total}}$ Γ_{22}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

85 ± 34	140	VANNUCCI	77	MRK1 $e^+e^- \rightarrow 3(\pi^+\pi^-)\pi^0$
-------------------------------	-----	----------	----	--

$\Gamma(\omega\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{23}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$0.40 \pm 0.06 \pm 0.04$	170	¹ AUBERT	06D	BABR $10.6 e^+e^- \rightarrow \omega\pi^+\pi^-\pi^0\gamma$
--	-----	---------------------	-----	--

¹ Using $\Gamma(J/\psi \rightarrow e^+e^-) = 5.52 \pm 0.14 \pm 0.04$ keV.

$\Gamma(\omega\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{24}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

8.6 ± 0.7 OUR AVERAGE Error includes scale factor of 1.1.

$9.7 \pm 0.6 \pm 0.6$	788	¹ AUBERT	07AU	BABR $10.6 e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$
-----------------------	-----	---------------------	------	---

7.0 ± 1.6	18058	AUGUSTIN	89	DM2 $J/\psi \rightarrow 2(\pi^+\pi^-)\pi^0$
---------------	-------	----------	----	---

7.8 ± 1.6	215	BURMESTER	77D	PLUT e^+e^-
---------------	-----	-----------	-----	---------------

6.8 ± 1.9	348	VANNUCCI	77	MRK1 $e^+e^- \rightarrow 2(\pi^+\pi^-)\pi^0$
---------------	-----	----------	----	--

¹ AUBERT 07AU quotes $\Gamma_{ee}^{J/\psi} \cdot B(J/\psi \rightarrow \omega\pi^+\pi^-) \cdot B(\omega \rightarrow 3\pi) = 47.8 \pm 3.1 \pm 3.2$ eV.

$\Gamma(\omega f_2(1270))/\Gamma_{\text{total}}$ Γ_{25}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
4.3±0.6 OUR AVERAGE				
4.3±0.2±0.6	5860	AUGUSTIN	89 DM2	e^+e^-
4.0±1.6	70	BURMESTER	77D PLUT	e^+e^-
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.9±0.8	81	VANNUCCI	77 MRK1	$e^+e^- \rightarrow 2(\pi^+\pi^-)\pi^0$

 $\Gamma(K^*(892)^0\bar{K}^*(892)^0)/\Gamma_{\text{total}}$ Γ_{26}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
2.3±0.7±0.1		25 ± 8	¹ AUBERT	07AK BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$
<5		90	VANNUCCI	77 MRK1	$e^+e^- \rightarrow \pi^+\pi^-K^+K^-$
¹ Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow K^*(892)^0\bar{K}^*(892)^0)/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+e^-)] = (1.28 \pm 0.40 \pm 0.11) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

 $\Gamma(K^*(892)^\pm K^*(892)^\mp)/\Gamma_{\text{total}}$ Γ_{27}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.00±0.19^{+0.11}_{-0.32}	323	ABLIKIM	10E BES2	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp \pi^0$

 $\Gamma(K^*(892)^\pm K^*(700)^\mp)/\Gamma_{\text{total}}$ Γ_{28}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.09±0.18^{+0.94}_{-0.54}	655	ABLIKIM	10E BES2	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp \pi^0$

 $\Gamma(\eta K^*(892)^0\bar{K}^*(892)^0)/\Gamma_{\text{total}}$ Γ_{31}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.15±0.13±0.22	209	ABLIKIM	10C BES2	$J/\psi \rightarrow \eta K^+\pi^- K^-\pi^+$

 $\Gamma(K^*(892)^0\bar{K}_2^*(1430)^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{38}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
5.9±0.6±0.2	317 ± 23	^{1,2} AUBERT	07AK BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$
6.7±2.6	40	VANNUCCI	77 MRK1	$e^+e^- \rightarrow \pi^+\pi^-K^+K^-$

¹Using $B(K_2^*(1430)^0 \rightarrow K\pi) = (49.9 \pm 1.2)\%$.

²Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow K^*(892)^0\bar{K}_2^*(1430)^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+e^-)] = (32.9 \pm 2.3 \pm 2.7) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\omega K^*(892)\bar{K} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{42}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
61 ± 9 OUR AVERAGE				
62.0 ± 6.8 ± 10.6	899 ± 98	ABLIKIM	08E BES2	$J/\psi \rightarrow \omega K_S^0 K^\pm \pi^\mp$
65.3 ± 10.2 ± 13.5	176 ± 28	ABLIKIM	08E BES2	$J/\psi \rightarrow \omega K^+ K^- \pi^0$
53 ± 14 ± 14	530 ± 140	BECKER	87 MRK3	$e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(K^+ K^*(892)^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{45}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
5.12 ± 0.30 OUR AVERAGE				
5.2 ± 0.4 ± 0.1		¹ AUBERT	08S BABR	10.6 $e^+ e^- \rightarrow K^+ K^*(892)^- \gamma$
4.57 ± 0.17 ± 0.70	2285	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
5.26 ± 0.13 ± 0.53		COFFMAN	88 MRK3	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp, K^+ K^- \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.6 ± 0.6	24	FRANKLIN	83 MRK2	$J/\psi \rightarrow K^+ K^- \pi^0$
3.2 ± 0.6	48	VANNUCCI	77 MRK1	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp$
4.1 ± 1.2	39	BRAUNSCH...	76 DASP	$J/\psi \rightarrow K^\pm X$

¹ AUBERT 08S reports $[\Gamma(J/\psi(1S) \rightarrow K^+ K^*(892)^- + \text{c.c.})/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = (29.0 \pm 1.7 \pm 1.3) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^+ K^*(892)^- + \text{c.c.} \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{46}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.97 ± 0.20 ± 0.05	155	¹ AUBERT	08S BABR	10.6 $e^+ e^- \rightarrow K^+ K^- \pi^0 \gamma$

¹ AUBERT 08S reports $[\Gamma(J/\psi(1S) \rightarrow K^+ K^*(892)^- + \text{c.c.} \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = (10.96 \pm 0.85 \pm 0.70) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^+ K^*(892)^- + \text{c.c.} \rightarrow K^+ K^- \pi^0)/\Gamma(K^+ K^- \pi^0)$ Γ_{46}/Γ_{141}

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
92.4 ± 1.5 ± 3.4	2K	¹ LEES	17C BABR	$J/\psi \rightarrow K^+ K^- \pi^0$

¹ From a Dalitz plot analysis in an isobar model.

$\Gamma(\bar{K} K^*(892) + \text{c.c.} \rightarrow K_S^0 K^\pm \pi^\mp)/\Gamma(K_S^0 K^\pm \pi^\mp)$ Γ_{44}/Γ_{142}

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
90.5 ± 0.9 ± 3.8	4K	¹ LEES	17C BABR	$J/\psi \rightarrow K_S^0 K^\pm \pi^\mp$

¹ From a Dalitz plot analysis in an isobar model.

$\Gamma(K^+ K^*(892)^- + \text{c.c.} \rightarrow K^0 K^\pm \pi^\mp + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{47}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
3.0 ± 0.4 ± 0.1	89	¹ AUBERT	08S BABR	10.6 $e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \gamma$

¹AUBERT 08S reports $[\Gamma(J/\psi(1S) \rightarrow K^+ K^*(892)^- + \text{c.c.} \rightarrow K^0 K^\pm \pi^\mp + \text{c.c.})/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = (16.76 \pm 1.70 \pm 1.00) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^0 \bar{K}^*(892)^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{48}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
4.39±0.31 OUR AVERAGE				
4.8 ± 0.5 ± 0.1		¹ AUBERT	08S BABR	10.6 $e^+ e^- \rightarrow K^0 \bar{K}^*(892)^0 \gamma$
3.96 ± 0.15 ± 0.60	1192	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
4.33 ± 0.12 ± 0.45		COFFMAN	88 MRK3	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.7 ± 0.6 45 VANNUCCI 77 MRK1 $J/\psi \rightarrow K^\pm K_S^0 \pi^\mp$

¹AUBERT 08S reports $[\Gamma(J/\psi(1S) \rightarrow K^0 \bar{K}^*(892)^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = (26.6 \pm 2.5 \pm 1.5) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^0 \bar{K}^*(892)^0 + \text{c.c.})/\Gamma(K^+ K^*(892)^- + \text{c.c.})$ Γ_{48}/Γ_{45}

VALUE	DOCUMENT ID	TECN	COMMENT
0.82±0.05±0.09	COFFMAN	88 MRK3	$J/\psi \rightarrow K \bar{K}^*(892) + \text{c.c.}$

$\Gamma(K^0 \bar{K}^*(892)^0 + \text{c.c.} \rightarrow K^0 K^\pm \pi^\mp + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{49}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
3.2±0.4±0.1	94	¹ AUBERT	08S BABR	10.6 $e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \gamma$

¹AUBERT 08S reports $[\Gamma(J/\psi(1S) \rightarrow K^0 \bar{K}^*(892)^0 + \text{c.c.} \rightarrow K^0 K^\pm \pi^\mp + \text{c.c.})/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = (17.70 \pm 1.70 \pm 1.00) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K_1(1400)^\pm K^\mp)/\Gamma_{\text{total}}$ Γ_{50}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
3.8±0.8±1.2	¹ BAI	99C BES	$e^+ e^-$

¹Assuming $B(K_1(1400) \rightarrow K^* \pi) = 0.94 \pm 0.06$

$\Gamma(\bar{K}^*(892)^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{51}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	¹ ABLIKIM	06C BES2	$J/\psi \rightarrow \bar{K}^*(892)^0 K^+ \pi^-$

¹A $K_0^*(700)$ is observed by ABLIKIM 06C in the $K^+ \pi^-$ mass spectrum of the $\bar{K}^*(892)^0 K^+ \pi^-$ final state against the $\bar{K}^*(892)$. A corresponding branching fraction of the $J/\psi(1S)$ is not presented.

$\Gamma(K^*(1410) \bar{K} + \text{c.c.} \rightarrow K^\pm K^\mp \pi^0)/\Gamma(K^+ K^- \pi^0)$ Γ_{33}/Γ_{141}

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
2.3±1.1±0.7	2K	¹ LEES	17C BABR	$J/\psi \rightarrow K^+ K^- \pi^0$

¹From a Dalitz plot analysis in an isobar model.

$\Gamma(K^*(1410)\bar{K} + \text{c.c.} \rightarrow K_S^0 K^\pm \pi^\mp) / \Gamma(K_S^0 K^\pm \pi^\mp)$ $\Gamma_{34} / \Gamma_{142}$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
1.5 ± 0.5 ± 0.9	4K	¹ LEES	17C BABR	$J/\psi \rightarrow K_S^0 K^\pm \pi^\mp$

¹ From a Dalitz plot analysis in an isobar model.

$\Gamma(K_2^*(1430)\bar{K} + \text{c.c.} \rightarrow K^\pm K^\mp \pi^0) / \Gamma(K^+ K^- \pi^0)$ $\Gamma_{36} / \Gamma_{141}$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
3.5 ± 1.3 ± 0.9	2K	¹ LEES	17C BABR	$J/\psi \rightarrow K^+ K^- \pi^0$

¹ From a Dalitz plot analysis in an isobar model.

$\Gamma(K_2^*(1430)\bar{K} + \text{c.c.} \rightarrow K_S^0 K^\pm \pi^\mp) / \Gamma(K_S^0 K^\pm \pi^\mp)$ $\Gamma_{37} / \Gamma_{142}$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
7.1 ± 1.3 ± 1.2	4K	¹ LEES	17C BABR	$J/\psi \rightarrow K_S^0 K^\pm \pi^\mp$

¹ From a Dalitz plot analysis in an isobar model.

$\Gamma(\omega \pi^0 \pi^0) / \Gamma_{\text{total}}$ Γ_{54} / Γ

VALUE (units 10 ⁻³)	EVTS	DOCUMENT ID	TECN	COMMENT
3.4 ± 0.3 ± 0.7	509	AUGUSTIN	89 DM2	$J/\psi \rightarrow \pi^+ \pi^- 3\pi^0$

$\Gamma(b_1(1235)^\pm \pi^\mp) / \Gamma_{\text{total}}$ Γ_{55} / Γ

VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT
30 ± 5 OUR AVERAGE				
31 ± 6	4600	AUGUSTIN	89 DM2	$J/\psi \rightarrow 2(\pi^+ \pi^-) \pi^0$
29 ± 7	87	BURMESTER	77D PLUT	$e^+ e^-$

$\Gamma(\omega K^\pm K_S^0 \pi^\mp) / \Gamma_{\text{total}}$ Γ_{56} / Γ

VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT
34 ± 5 OUR AVERAGE				
37.7 ± 0.8 ± 5.8	1972 ± 41	ABLIKIM	08E BES2	$e^+ e^- \rightarrow J/\psi$
29.5 ± 1.4 ± 7.0	879 ± 41	BECKER	87 MRK3	$e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(b_1(1235)^0 \pi^0) / \Gamma_{\text{total}}$ Γ_{57} / Γ

VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT
23 ± 3 ± 5	229	AUGUSTIN	89 DM2	$e^+ e^-$

$\Gamma(\eta K^\pm K_S^0 \pi^\mp) / \Gamma_{\text{total}}$ Γ_{58} / Γ

VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT
21.8 ± 2.2 ± 3.4	232 ± 23	ABLIKIM	08E BES2	$e^+ e^- \rightarrow J/\psi$

$\Gamma(\phi K^*(892)\bar{K} + \text{c.c.}) / \Gamma_{\text{total}}$ Γ_{59} / Γ

VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT
21.8 ± 2.3 OUR AVERAGE				
20.8 ± 2.7 ± 3.9	195 ± 25	ABLIKIM	08E BES2	$J/\psi \rightarrow \phi K_S^0 K^\pm \pi^\mp$
29.6 ± 3.7 ± 4.7	238 ± 30	ABLIKIM	08E BES2	$J/\psi \rightarrow \phi K^+ K^- \pi^0$
20.7 ± 2.4 ± 3.0		FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
20 ± 3 ± 3	155 ± 20	BECKER	87 MRK3	$e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(\omega K\bar{K})/\Gamma_{\text{total}}$ **Γ_{60}/Γ**

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
17.0 ± 3.2 OUR AVERAGE				
13.6 ± 5.0 ± 1.0	24	¹ AUBERT 07AU BABR		10.6 $e^+e^- \rightarrow \omega K^+ K^- \gamma$
19.8 ± 2.1 ± 3.9		² FALVARD 88 DM2		$J/\psi \rightarrow \text{hadrons}$
16 ± 10	22	FELDMAN 77 MRK1		e^+e^-

¹ AUBERT 07AU quotes $\Gamma_{ee}^{J/\psi} \cdot B(J/\psi \rightarrow \omega K^+ K^-) \cdot B(\eta \rightarrow 3\pi) = 3.3 \pm 1.3 \pm 0.2 \text{ eV}$.
² Addition of $\omega K^+ K^-$ and $\omega K^0 \bar{K}^0$ branching ratios.

$\Gamma(\omega f_0(1710) \rightarrow \omega K\bar{K})/\Gamma_{\text{total}}$ **Γ_{61}/Γ**

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
4.8 ± 1.1 ± 0.3	^{1,2} FALVARD 88 DM2		$J/\psi \rightarrow \text{hadrons}$

¹ Includes unknown branching fraction $f_0(1710) \rightarrow K\bar{K}$.
² Addition of $f_0(1710) \rightarrow K^+ K^-$ and $f_0(1710) \rightarrow K^0 \bar{K}^0$ branching ratios.

$\Gamma(\phi 2(\pi^+ \pi^-))/\Gamma_{\text{total}}$ **Γ_{62}/Γ**

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
16.6 ± 2.3 OUR AVERAGE				
17.3 ± 3.3 ± 1.2	35	¹ AUBERT 06D BABR		10.6 $e^+e^- \rightarrow \phi 2(\pi^+ \pi^-) \gamma$
16.0 ± 1.0 ± 3.0		FALVARD 88 DM2		$J/\psi \rightarrow \text{hadrons}$

¹ Using $\Gamma(J/\psi \rightarrow e^+e^-) = 5.52 \pm 0.14 \pm 0.04 \text{ keV}$.

$\Gamma(\Delta(1232)^{++} \bar{p}\pi^-)/\Gamma_{\text{total}}$ **Γ_{63}/Γ**


VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.58 ± 0.23 ± 0.40	332	EATON 84 MRK2		e^+e^-

$\Gamma(\omega\eta)/\Gamma_{\text{total}}$ **Γ_{64}/Γ**

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.74 ± 0.20 OUR AVERAGE				Error includes scale factor of 1.6. See the ideogram below.
2.352 ± 0.273	5k	¹ ABLIKIM 06F BES2		$J/\psi \rightarrow \omega\eta$
1.44 ± 0.40 ± 0.14	13	² AUBERT 06D BABR		10.6 $e^+e^- \rightarrow \omega\eta\gamma$
1.43 ± 0.10 ± 0.21	378	JOUSSET 90 DM2		$J/\psi \rightarrow \text{hadrons}$
1.71 ± 0.08 ± 0.20		COFFMAN 88 MRK3		$e^+e^- \rightarrow 3\pi\eta$

¹ Using $B(\eta \rightarrow 2\gamma) = (39.43 \pm 0.26)\%$, $B(\eta \rightarrow \pi^+ \pi^- \pi^0) = 22.6 \pm 0.4\%$, $B(\eta \rightarrow \pi^+ \pi^- \gamma) = 4.68 \pm 0.11\%$, and $B(\omega \rightarrow \pi^+ \pi^- \pi^0) = (89.1 \pm 0.7)\%$.
² Using $\Gamma(J/\psi \rightarrow e^+e^-) = 5.52 \pm 0.14 \pm 0.04 \text{ keV}$.

WEIGHTED AVERAGE
 1.74 ± 0.20 (Error scaled by 1.6)



$\Gamma(\phi K \bar{K})/\Gamma_{\text{total}}$

Γ_{65}/Γ


VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
17.7 ± 1.6	OUR AVERAGE	Error includes scale factor of 1.3. See the ideogram below.		
$16.6 \pm 1.9 \pm 1.2$	163 ± 19	LEES	12F BABR	$10.6 e^+ e^- \rightarrow 2(K^+ K^-) \gamma$
$21.4 \pm 0.4 \pm 2.2$		ABLIKIM	05 BES2	$J/\psi \rightarrow \phi \pi^+ \pi^-$
$48^{+20}_{-16} \pm 6$	$9.0^{+3.7}_{-3.0}$	^{1,2} HUANG	03 BELL	$B^+ \rightarrow (\phi K^+ K^-) K^+$
$14.6 \pm 0.8 \pm 2.1$		³ FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
18 ± 8	14	FELDMAN	77 MRK1	$e^+ e^-$

¹ We have multiplied $K^+ K^-$ measurement by 2 to obtain $K \bar{K}$.

² Using $B(B^+ \rightarrow J/\psi K^+) = (1.01 \pm 0.05) \times 10^{-3}$.

³ Addition of $\phi K^+ K^-$ and $\phi K^0 \bar{K}^0$ branching ratios.

WEIGHTED AVERAGE
 17.7 ± 1.6 (Error scaled by 1.3)



$\Gamma(\phi f_0(1710) \rightarrow \phi K \bar{K})/\Gamma_{\text{total}}$ **Γ_{67}/Γ**

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$3.6 \pm 0.2 \pm 0.6$	1,2 FALVARD	88 DM2	$J/\psi \rightarrow$ hadrons

¹ Including interference with $f'_2(1525)$.
² Includes unknown branching fraction $f_0(1710) \rightarrow K \bar{K}$.

$\Gamma(\phi f_2(1270))/\Gamma_{\text{total}}$ **Γ_{69}/Γ**

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
$0.72 \pm 0.13 \pm 0.02$		44 ± 7	1,2 AUBERT	07AK BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$
< 0.45	90		FALVARD	88 DM2	$J/\psi \rightarrow$ hadrons
< 0.37	90		VANNUCCI	77 MRK1	$e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^-$

¹ Using $B(f_2(1270) \rightarrow \pi\pi) = (84.8^{+2.4}_{-1.2})\%$
² Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_2(1270))/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = (4.02 \pm 0.65 \pm 0.33) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\Delta(1232)^{++} \bar{\Delta}(1232)^{--})/\Gamma_{\text{total}}$ **Γ_{70}/Γ**

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.10 \pm 0.09 \pm 0.28$	233	EATON	84 MRK2	$e^+ e^-$

$\Gamma(\Sigma(1385)^-\bar{\Sigma}(1385)^+ \text{ (or c.c.)})/\Gamma_{\text{total}}$ Γ_{71}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.16 ± 0.05	OUR AVERAGE			
1.096 ± 0.012 ± 0.071	43K	ABLIKIM	16L BES3	$J/\psi \rightarrow \Sigma(1385)^-\bar{\Sigma}(1385)^+$
1.258 ± 0.014 ± 0.078	53k	ABLIKIM	16L BES3	$J/\psi \rightarrow \Sigma(1385)^+\bar{\Sigma}(1385)^-$
1.23 ± 0.07 ± 0.30	0.8k	ABLIKIM	12P BES2	$J/\psi \rightarrow \Sigma(1385)^-\bar{\Sigma}(1385)^+$
1.50 ± 0.08 ± 0.38	1k	ABLIKIM	12P BES2	$J/\psi \rightarrow \Sigma(1385)^+\bar{\Sigma}(1385)^-$
1.00 ± 0.04 ± 0.21	0.6k	HENRARD	87 DM2	$e^+e^- \rightarrow \Sigma^{*-}$
1.19 ± 0.04 ± 0.25	0.7k	HENRARD	87 DM2	$e^+e^- \rightarrow \Sigma^{*+}$
0.86 ± 0.18 ± 0.22	56	EATON	84 MRK2	$e^+e^- \rightarrow \Sigma^{*-}$
1.03 ± 0.24 ± 0.25	68	EATON	84 MRK2	$e^+e^- \rightarrow \Sigma^{*+}$

 $\Gamma(\Sigma(1385)^0\bar{\Sigma}(1385)^0)/\Gamma_{\text{total}}$ Γ_{72}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.071 ± 0.009 ± 0.082	103k	ABLIKIM	17E BES3	$e^+e^- \rightarrow J/\psi \rightarrow$ hadrons


 $\Gamma(\phi f'_2(1525))/\Gamma_{\text{total}}$ Γ_{74}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
8 ± 4	OUR AVERAGE	Error includes scale factor of 2.7.		
12.3 ± 0.6 ± 2.0		^{1,2} FALVARD	88 DM2	$J/\psi \rightarrow$ hadrons
4.8 ± 1.8	46	¹ GIDAL	81 MRK2	$J/\psi \rightarrow K^+K^-K^+K^-$
¹ Re-evaluated using $B(f'_2(1525) \rightarrow K\bar{K}) = 0.713$.				
² Including interference with $f_0(1710)$.				

 $\Gamma(\phi\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{75}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.87 ± 0.09	OUR AVERAGE	Error includes scale factor of 1.4. See the ideogram below.		
0.81 ± 0.08 ± 0.03	181	LEES	12F BABR	10.6 $e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
1.09 ± 0.02 ± 0.13		ABLIKIM	05 BES2	$J/\psi \rightarrow \phi\pi^+\pi^-$
0.78 ± 0.03 ± 0.12		FALVARD	88 DM2	$J/\psi \rightarrow$ hadrons
2.1 ± 0.9	23	FELDMAN	77 MRK1	e^+e^-
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.96 ± 0.13	103	¹ AUBERT, BE	06D BABR	10.6 $e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
¹ Superseded by LEES 12F. Derived by us. AUBERT, BE 06D measures $\Gamma(J/\psi \rightarrow e^+e^-)$ $\times B(J/\psi \rightarrow \phi\pi^+\pi^-) \times B(\phi \rightarrow K^+K^-) = (2.61 \pm 0.30 \pm 0.18)$ eV				

WEIGHTED AVERAGE
 0.87 ± 0.09 (Error scaled by 1.4)



$\Gamma(\phi\pi^0\pi^0)/\Gamma_{\text{total}}$

Γ_{76}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.56 ± 0.16	23	¹ AUBERT, BE 06D	BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^0 \pi^0 \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •
¹ Superseded by LEES 12F. Derived by us. AUBERT, BE 06D measures $\Gamma(J/\psi \rightarrow e^+ e^-) \times B(J/\psi \rightarrow \phi \pi^0 \pi^0) \times B(\phi \rightarrow K^+ K^-) = (1.54 \pm 0.40 \pm 0.16) \text{ eV}$

$\Gamma(\phi K^\pm K_S^0 \pi^\mp)/\Gamma_{\text{total}}$

Γ_{77}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
7.2 ± 0.8 OUR AVERAGE				
$7.4 \pm 0.6 \pm 1.4$	227 ± 19	ABLIKIM	08E	BES2 $e^+ e^- \rightarrow J/\psi$
$7.4 \pm 0.9 \pm 1.1$		FALVARD	88	DM2 $J/\psi \rightarrow \text{hadrons}$
$7 \pm 0.6 \pm 1.0$	163 ± 15	BECKER	87	MRK3 $e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(\omega f_1(1420))/\Gamma_{\text{total}}$

Γ_{78}/Γ


VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$6.8^{+1.9}_{-1.6} \pm 1.7$	111^{+31}_{-26}	BECKER	87	MRK3 $e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(\phi\eta)/\Gamma_{\text{total}}$

Γ_{79}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.75 ± 0.08 OUR AVERAGE				Error includes scale factor of 1.5. See the ideogram below.
$1.4 \pm 0.6 \pm 0.1$	6	¹ AUBERT	07AU	BABR $10.6 e^+ e^- \rightarrow \phi \eta \gamma$
$0.898 \pm 0.024 \pm 0.089$		ABLIKIM	05B	BES2 $e^+ e^- \rightarrow J/\psi \rightarrow \text{hadr}$
$0.64 \pm 0.04 \pm 0.11$	346	JOUSSET	90	DM2 $J/\psi \rightarrow \text{hadrons}$
$0.661 \pm 0.045 \pm 0.078$		COFFMAN	88	MRK3 $e^+ e^- \rightarrow K^+ K^- \eta$

¹ AUBERT 07AU quotes $\Gamma_{ee}^{J/\psi} \cdot B(J/\psi \rightarrow \phi\eta) \cdot B(\phi \rightarrow K^+K^-) \cdot B(\eta \rightarrow \gamma\gamma) = 0.84 \pm 0.37 \pm 0.05$ eV.



$\Gamma(\Xi^0 \Xi^0)/\Gamma_{\text{total}}$ Γ_{80}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.17 ± 0.04	OUR AVERAGE			
1.165 ± 0.004 ± 0.043	135K	ABLIKIM 17E	BES3	$e^+e^- \rightarrow J/\psi \rightarrow$ hadrons
1.20 ± 0.12 ± 0.21	206	ABLIKIM 080	BES2	$e^+e^- \rightarrow J/\psi$

$\Gamma(\Xi(1530)^- \Xi^+)/\Gamma_{\text{total}}$ Γ_{81}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.59 ± 0.09 ± 0.12	75 ± 11	HENRARD 87	DM2	e^+e^-

$\Gamma(pK^- \bar{\Sigma}(1385)^0)/\Gamma_{\text{total}}$ Γ_{82}/Γ


VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.51 ± 0.26 ± 0.18	89	EATON 84	MRK2	e^+e^-

$\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$ Γ_{83}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.45 ± 0.05	OUR AVERAGE			
0.538 ± 0.012 ± 0.065	2090	¹ ABLIKIM 06F	BES2	$J/\psi \rightarrow \omega\pi^0$
0.360 ± 0.028 ± 0.054	222	JOUSSET 90	DM2	$J/\psi \rightarrow$ hadrons
0.482 ± 0.019 ± 0.064		COFFMAN 88	MRK3	$e^+e^- \rightarrow \pi^0\pi^+\pi^-\pi^0$

Error includes scale factor of 1.4. See the ideogram below.

¹ Using $B(\omega \rightarrow \pi^+ \pi^- \pi^0) = (89.1 \pm 0.7)\%$.



$\Gamma(\omega\pi^0 \rightarrow \pi^+\pi^-\pi^0)/\Gamma(\pi^+\pi^-\pi^0)$ Γ_{84}/Γ_{129}

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$8 \pm 3 \pm 2$	20K	¹ LEES	17C BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$

¹ From a Dalitz plot analysis in an isobar model and significance 4.9σ .


$\Gamma(\phi\eta'(958))/\Gamma_{\text{total}}$ Γ_{85}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
4.6 ± 0.5	OUR AVERAGE		Error includes scale factor of 2.2. See the ideogram below.		
$5.10 \pm 0.03 \pm 0.32$		31k	ABLIKIM 16M	BES3	$e^+e^- \rightarrow J/\psi \rightarrow \text{hadrons}$
$5.46 \pm 0.31 \pm 0.56$			ABLIKIM 05B	BES2	$e^+e^- \rightarrow J/\psi \rightarrow \text{hadrons}$
$4.1 \pm 0.3 \pm 0.8$		167	JOUSSET 90	DM2	$J/\psi \rightarrow \text{hadrons}$
$3.08 \pm 0.34 \pm 0.36$			COFFMAN 88	MRK3	$e^+e^- \rightarrow K^+K^-\eta'$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 13	90	VANNUCCI 77	MRK1	e^+e^-
------	----	-------------	------	----------

WEIGHTED AVERAGE
 4.6 ± 0.5 (Error scaled by 2.2)



$\Gamma(\phi f_0(980)) / \Gamma_{\text{total}}$

Γ_{86} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.2 ± 0.9 OUR AVERAGE				Error includes scale factor of 1.9.
$4.6 \pm 0.4 \pm 0.8$		¹ FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
2.6 ± 0.6	50	¹ GIDAL	81 MRK2	$J/\psi \rightarrow K^+ K^- K^+ K^-$

¹ Assuming $B(f_0(980) \rightarrow \pi\pi) = 0.78$.

$\Gamma(\phi f_0(980) \rightarrow \phi \pi^+ \pi^-) / \Gamma_{\text{total}}$

Γ_{87} / Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$0.182 \pm 0.042 \pm 0.005$	19.5 ± 4.5	^{1,2} AUBERT	07AK BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

¹ Using $B(\phi \rightarrow K^+ K^-) = (49.3 \pm 0.6)\%$.

² Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi \pi^+ \pi^-) / \Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = (1.01 \pm 0.22 \pm 0.08) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi f_0(980) \rightarrow \phi \pi^0 \pi^0) / \Gamma_{\text{total}}$

Γ_{88} / Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$0.171 \pm 0.073 \pm 0.004$	7.0 ± 2.8	^{1,2} AUBERT	07AK BABR	$10.6 e^+ e^- \rightarrow \pi^0 \pi^0 K^+ K^- \gamma$

¹ Using $B(\phi \rightarrow K^+ K^-) = (49.3 \pm 0.6)\%$.

² Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi \pi^0 \pi^0) / \Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = (0.95 \pm 0.39 \pm 0.10) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi \pi^0 f_0(980) \rightarrow \phi \pi^0 \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{89} / Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
4.50 ± 0.80 ± 0.61	355	ABLIKIM	15P	BES3 $J/\psi \rightarrow K^+ K^- 3\pi$

$\Gamma(\phi \pi^0 f_0(980) \rightarrow \phi \pi^0 \rho^0 \pi^0) / \Gamma_{\text{total}}$ Γ_{90} / Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
1.67 ± 0.50 ± 0.24	70	ABLIKIM	15P	BES3 $J/\psi \rightarrow K^+ K^- 3\pi$

$\Gamma(\eta \phi f_0(980) \rightarrow \eta \phi \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{91} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.23 ± 0.75 ± 0.73	52	ABLIKIM	08F	BES $J/\psi \rightarrow \eta \phi f_0(980)$

$\Gamma(\phi a_0(980)^0 \rightarrow \phi \eta \pi^0) / \Gamma_{\text{total}}$ Γ_{92} / Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
5.0 ± 2.7 ± 2.5	¹ ABLIKIM	11D	BES3 $J/\psi \rightarrow \phi \eta \pi^0$

¹ Assuming $a_0(980) - f_0(980)$ mixing and isospin breaking via γ^* and $K^* K$ loops.

$\Gamma(\Xi(1530)^0 \Xi^0) / \Gamma_{\text{total}}$ Γ_{93} / Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.32 ± 0.12 ± 0.07	24 ± 9	HENRARD	87	DM2 $e^+ e^-$

$\Gamma(\Sigma(1385)^- \bar{\Sigma}^+ (\text{or c.c.})) / \Gamma_{\text{total}}$ Γ_{94} / Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.31 ± 0.05 OUR AVERAGE				
0.30 ± 0.03 ± 0.07	74 ± 8	HENRARD	87	DM2 $e^+ e^- \rightarrow \Sigma^{*-}$
0.34 ± 0.04 ± 0.07	77 ± 9	HENRARD	87	DM2 $e^+ e^- \rightarrow \Sigma^{*+}$
0.29 ± 0.11 ± 0.10	26	EATON	84	MRK2 $e^+ e^- \rightarrow \Sigma^{*-}$
0.31 ± 0.11 ± 0.11	28	EATON	84	MRK2 $e^+ e^- \rightarrow \Sigma^{*+}$

$\Gamma(\phi f_1(1285)) / \Gamma_{\text{total}}$ Γ_{95} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.6 ± 0.5 OUR AVERAGE				
3.4 ± 1.8 ± 1.5	1.1k	¹ ABLIKIM	15H	BES3 $e^+ e^- \rightarrow J/\psi \rightarrow \phi \eta \pi^+ \pi^-$
3.2 ± 0.6 ± 0.4		JOUSSET	90	DM2 $J/\psi \rightarrow \phi 2(\pi^+ \pi^-)$
2.1 ± 0.5 ± 0.4	25	² JOUSSET	90	DM2 $J/\psi \rightarrow \phi \eta \pi^+ \pi^-$
• • •				We do not use the following data for averages, fits, limits, etc. • • •
0.6 ± 0.2 ± 0.1	16	BECKER	87	MRK3 $J/\psi \rightarrow \phi K \bar{K} \pi$

¹ ABLIKIM 15H reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_1(1285))/\Gamma_{\text{total}}] \times [B(f_1(1285) \rightarrow \eta\pi^+\pi^-)] = (1.20 \pm 0.6 \pm 0.14) \times 10^{-4}$ which we divide by our best value $B(f_1(1285) \rightarrow \eta\pi^+\pi^-) = (35 \pm 15) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² We attribute to the $f_1(1285)$ the signal observed in the $\pi^+\pi^-\eta$ invariant mass distribution at 1297 MeV.

$\Gamma(\phi f_1(1285) \rightarrow \phi\pi^0 f_0(980) \rightarrow \phi\pi^0\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{96}/Γ

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
9.36±2.31±1.54	78	ABLIKIM	15P	BES3 $J/\psi \rightarrow K^+K^-3\pi$

$\Gamma(\phi f_1(1285) \rightarrow \phi\pi^0 f_0(980) \rightarrow \phi\pi^0\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_{97}/Γ

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
2.08±1.63±1.47	9	ABLIKIM	15P	BES3 $J/\psi \rightarrow K^+K^-3\pi$

$\Gamma(\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{98}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.40±0.17±0.03	9	¹ AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow \eta\pi^+\pi^-\gamma$

¹ AUBERT 07AU quotes $\Gamma_{ee}^{J/\psi} \cdot B(J/\psi \rightarrow \eta\pi^+\pi^-) \cdot B(\eta \rightarrow 3\pi) = 0.51 \pm 0.22 \pm 0.03$ eV.

$\Gamma(\eta\rho)/\Gamma_{\text{total}}$ Γ_{99}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.193±0.023 OUR AVERAGE				
0.194±0.017±0.029	299	JOUSSET	90	DM2 $J/\psi \rightarrow \text{hadrons}$
0.193±0.013±0.029		COFFMAN	88	MRK3 $e^+e^- \rightarrow \pi^+\pi^-\eta$

$\Gamma(\omega\eta'(958))/\Gamma_{\text{total}}$ Γ_{100}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.89±0.18 OUR AVERAGE				
2.08±0.30±0.14	137	¹ ABLIKIM	17AK	BES3 $J/\psi \rightarrow \pi^+\pi^-\eta'$
2.26±0.43	218	² ABLIKIM	06F	BES2 $J/\psi \rightarrow \omega\eta'$
1.8 $^{+1.0}_{-0.8}$ ±0.3	6	JOUSSET	90	DM2 $J/\psi \rightarrow \text{hadrons}$
1.66±0.17±0.19		COFFMAN	88	MRK3 $e^+e^- \rightarrow 3\pi\eta'$

¹ From a partial wave analysis of the decay $J/\psi \rightarrow \pi^+\pi^-\eta'$.

² Using $B(\eta' \rightarrow \pi^+\pi^-\eta) = (44.3 \pm 1.5)\%$, $B(\eta' \rightarrow \pi^+\pi^-\gamma) = 29.5 \pm 1.0\%$, $B(\eta \rightarrow 2\gamma) = 39.43 \pm 0.26\%$, and $B(\omega \rightarrow \pi^+\pi^-\pi^0) = (89.1 \pm 0.7)\%$.

$\Gamma(\omega f_0(980))/\Gamma_{\text{total}}$ Γ_{101}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
1.41±0.27±0.47	¹ AUGUSTIN	89	DM2 $J/\psi \rightarrow 2(\pi^+\pi^-)\pi^0$

¹ Assuming $B(f_0(980) \rightarrow \pi\pi) = 0.78$.

$\Gamma(\rho\eta'(958))/\Gamma_{\text{total}}$ Γ_{102}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
8.1 ± 0.8 OUR AVERAGE	Error	includes scale factor of 1.6.		
7.90 ± 0.19 ± 0.49	3476	¹ ABLIKIM	17AK BES3	$J/\psi \rightarrow \pi^+ \pi^- \eta'$
8.3 ± 3.0 ± 1.2	19	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
11.4 ± 1.4 ± 1.6		COFFMAN	88 MRK3	$J/\psi \rightarrow \pi^+ \pi^- \eta'$

¹ From a partial wave analysis of the decay $J/\psi \rightarrow \pi^+ \pi^- \eta'$.

$\Gamma(a_2(1320)^\pm \pi^\mp)/\Gamma_{\text{total}}$ Γ_{103}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<43	90	BRAUNSCH...	76 DASP	$e^+ e^-$

$\Gamma(K \bar{K}_2^*(1430) + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{104}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<40	90	VANNUCCI	77 MRK1	$e^+ e^- \rightarrow K^0 \bar{K}_2^{*0}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<66	90	BRAUNSCH...	76 DASP	$e^+ e^- \rightarrow K^\pm \bar{K}_2^{*\mp}$
-----	----	-------------	---------	--

$\Gamma(K_1(1270)^\pm K^\mp)/\Gamma_{\text{total}}$ Γ_{105}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<3.0	90	¹ BAI	99C BES	$e^+ e^-$

¹ Assuming $B(K_1(1270) \rightarrow K\rho) = 0.42 \pm 0.06$

$\Gamma(K_2^*(1430)^0 \bar{K}_2^*(1430)^0)/\Gamma_{\text{total}}$ Γ_{108}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<29	90	VANNUCCI	77 MRK1	$e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^-$

$\Gamma(\phi\pi^0)/\Gamma_{\text{total}}$ Γ_{109}/Γ

The two different fit values of ABLIKIM 15K below have the same statistical significance of 6.4σ and cannot be distinguished at this moment.

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.94 ± 0.16 ± 0.16		0.8k	¹ ABLIKIM	15K BES3	$e^+ e^- \rightarrow J/\psi \rightarrow K^+ K^- \gamma\gamma$
0.124 ± 0.033 ± 0.030		35 ± 9	² ABLIKIM	15K BES3	$e^+ e^- \rightarrow J/\psi \rightarrow K^+ K^- \gamma\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6.4	90	³ ABLIKIM	05B BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \phi\gamma\gamma$
<6.8	90	COFFMAN	88 MRK3	$e^+ e^- \rightarrow K^+ K^- \pi^0$

¹ Corresponding to one of the two fit solutions with $\delta = (-95.9 \pm 1.5)^\circ$ for the phase angle between the resonant $J/\psi \rightarrow \phi\pi^0$ and non-phi $J/\psi \rightarrow K^+ K^- \pi^0$ contributions.

² Corresponding to one of the two fit solutions with $\delta = (-152.1 \pm 7.7)^\circ$ for the phase angle between the resonant $J/\psi \rightarrow \phi\pi^0$ and non-phi $J/\psi \rightarrow K^+ K^- \pi^0$ contributions.

³ Superseded by ABLIKIM 15K.

$\Gamma(\phi\eta(1405) \rightarrow \phi\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{110}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$2.01 \pm 0.58 \pm 0.82$		172	¹ ABLIKIM	15H BES3	$e^+e^- \rightarrow J/\psi \rightarrow \phi\eta\pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 17 90 ² FALVARD 88 DM2 $J/\psi \rightarrow \text{hadrons}$

¹ With 3.6σ significance.

² Includes unknown branching fraction $\eta(1405) \rightarrow \eta\pi\pi$.

 $\Gamma(\omega f'_2(1525))/\Gamma_{\text{total}}$ Γ_{111}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 2.2	90	¹ VANNUCCI 77	MRK1	$e^+e^- \rightarrow \pi^+\pi^-\pi^0 K^+K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 2.8 90 ¹ FALVARD 88 DM2 $J/\psi \rightarrow \text{hadrons}$

¹ Re-evaluated assuming $B(f'_2(1525) \rightarrow K\bar{K}) = 0.713$.

 $\Gamma(\omega X(1835) \rightarrow \omega p\bar{p})/\Gamma_{\text{total}}$ Γ_{112}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 3.9	95	ABLIKIM	13P BES3	$J/\psi \rightarrow \gamma\pi^0 p\bar{p}$

 $\Gamma(\phi X(1835) \rightarrow \phi p\bar{p})/\Gamma_{\text{total}}$ Γ_{113}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 2.1×10^{-7}	90	¹ ABLIKIM	16K BES3	$J/\psi \rightarrow p\bar{p}K_S^0 K_L^0, p\bar{p}K^+K^-$

¹ Upper limit applies to any $p\bar{p}$ mass enhancement near threshold.

 $\Gamma(\phi X(1835) \rightarrow \phi\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{114}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 2.8×10^{-4}	90	ABLIKIM	15H BES3	$e^+e^- \rightarrow J/\psi \rightarrow \phi\eta\pi^+\pi^-$

 $\Gamma(\phi X(1870) \rightarrow \phi\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{115}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 6.13×10^{-5}	90	ABLIKIM	15H BES3	$e^+e^- \rightarrow J/\psi \rightarrow \phi\eta\pi^+\pi^-$

 $\Gamma(\eta\phi(2170) \rightarrow \eta\phi f_0(980) \rightarrow \eta\phi\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{116}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.20 \pm 0.14 \pm 0.37$	471	ABLIKIM	15H BES3	$e^+e^- \rightarrow J/\psi \rightarrow \phi\eta\pi^+\pi^-$

 $\Gamma(\eta\phi(2170) \rightarrow \eta K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}$ Γ_{117}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 2.52	90	ABLIKIM	10C BES2	$J/\psi \rightarrow \eta K^+\pi^- K^-\pi^+$

 $\Gamma(\Sigma(1385)^0 \bar{\Lambda} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{118}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
< 0.82	90	ABLIKIM	13F BES3	$J/\psi \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 20 90 HENRARD 87 DM2 e^+e^-

$\Gamma(\Delta(1232)^+\bar{p})/\Gamma_{\text{total}} \quad \Gamma_{119}/\Gamma$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.1	90	HENRARD 87	DM2	e^+e^-

 $\Gamma(\Lambda(1520)\bar{\Lambda} + \text{c.c.} \rightarrow \gamma\Lambda\bar{\Lambda})/\Gamma_{\text{total}} \quad \Gamma_{120}/\Gamma$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<4.1	90	ABLIKIM 12B	BES3	$J/\psi \rightarrow \Lambda\bar{\Lambda}\gamma$

 $\Gamma(\Theta(1540)\bar{\Theta}(1540) \rightarrow K_S^0 p K^- \bar{n} + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_{121}/\Gamma$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<1.1	90	BAI 04G	BES2	e^+e^-

 $\Gamma(\Theta(1540)K^- \bar{n} \rightarrow K_S^0 p K^- \bar{n})/\Gamma_{\text{total}} \quad \Gamma_{122}/\Gamma$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<2.1	90	BAI 04G	BES2	e^+e^-

 $\Gamma(\Theta(1540)K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n)/\Gamma_{\text{total}} \quad \Gamma_{123}/\Gamma$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<1.6	90	BAI 04G	BES2	e^+e^-

 $\Gamma(\bar{\Theta}(1540)K^+ n \rightarrow K_S^0 \bar{p} K^+ n)/\Gamma_{\text{total}} \quad \Gamma_{124}/\Gamma$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<5.6	90	BAI 04G	BES2	e^+e^-

 $\Gamma(\bar{\Theta}(1540)K_S^0 p \rightarrow K_S^0 p K^- \bar{n})/\Gamma_{\text{total}} \quad \Gamma_{125}/\Gamma$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<1.1	90	BAI 04G	BES2	e^+e^-

 $\Gamma(\Sigma^0 \bar{\Lambda})/\Gamma_{\text{total}} \quad \Gamma_{126}/\Gamma$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.9	90	HENRARD 87	DM2	e^+e^-


————— STABLE HADRONS —————

 $\Gamma(2(\pi^+ \pi^-) \pi^0)/\Gamma_{\text{total}} \quad \Gamma_{127}/\Gamma$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
4.1 ± 0.5 OUR AVERAGE		Error includes scale factor of 2.4. See the ideogram below.		
5.46 ± 0.34 ± 0.14	4990	¹ AUBERT 07AU	BABR	10.6 $e^+e^- \rightarrow 2(\pi^+ \pi^-) \pi^0 \gamma$
3.25 ± 0.49	46055	AUGUSTIN 89	DM2	$J/\psi \rightarrow 2(\pi^+ \pi^-) \pi^0$
3.17 ± 0.42	147	FRANKLIN 83	MRK2	$e^+e^- \rightarrow \text{hadrons}$
3.64 ± 0.52	1500	BURMESTER 77D	PLUT	e^+e^-
4 ± 1	675	JEAN-MARIE 76	MRK1	e^+e^-

¹ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow 2(\pi^+ \pi^-) \pi^0)/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = 0.303 \pm 0.005 \pm 0.018 \text{ keV}$ which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02 \text{ keV}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

WEIGHTED AVERAGE
 4.1 ± 0.5 (Error scaled by 2.4)



$\Gamma(\omega \pi^+ \pi^-) / \Gamma(2(\pi^+ \pi^-) \pi^0)$

$\Gamma_{24} / \Gamma_{127}$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.3	¹ JEAN-MARIE	76	MRK1	e ⁺ e ⁻
-----	-------------------------	----	------	-------------------------------

¹ Final state $(\pi^+ \pi^-) \pi^0$ under the assumption that $\pi \pi$ is isospin 0.

$\Gamma(3(\pi^+ \pi^-) \pi^0) / \Gamma_{total}$

Γ_{128} / Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.029 ± 0.006 OUR AVERAGE

0.028 ± 0.009	11	FRANKLIN	83	MRK2 e ⁺ e ⁻ → hadrons
0.029 ± 0.007	181	JEAN-MARIE	76	MRK1 e ⁺ e ⁻

$\Gamma(\pi^+ \pi^- \pi^0) / \Gamma_{total}$

Γ_{129} / Γ

VALUE (units 10 ⁻³)	EVTS	DOCUMENT ID	TECN	COMMENT
---------------------------------	------	-------------	------	---------


21.1 ± 0.7 OUR AVERAGE Error includes scale factor of 1.5. See the ideogram below.

21.37 ± 0.04 ^{+0.64} _{-0.62}	1.8M	^{1,2} ABLIKIM	12H	BES3 e ⁺ e ⁻ → J/ψ
23.0 ± 2.0 ± 0.4	256	³ AUBERT	07AU	BABR 10.6 e ⁺ e ⁻ → J/ψ π ⁺ π ⁻ γ
21.8 ± 1.9		^{4,5} AUBERT,B	04N	BABR 10.6 e ⁺ e ⁻ → π ⁺ π ⁻ π ⁰ γ
21.84 ± 0.05 ± 2.01	220k	^{1,5} BAI	04H	BES e ⁺ e ⁻
20.91 ± 0.21 ± 1.16		^{5,6} BAI	04H	BES e ⁺ e ⁻
15 ± 2	168	FRANKLIN	83	MRK2 e ⁺ e ⁻

¹ From J/ψ → π⁺ π⁻ π⁰ events directly.

² The quoted systematic error includes a contribution of 1.23% (added in quadrature) from the uncertainty on the number of J/ψ events.

- ³ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] = (18.6 \pm 1.2 \pm 1.1) \times 10^{-3}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}} = 0.809 \pm 0.013$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ⁴ From the ratio of $\Gamma(e^+e^-) B(\pi^+\pi^-\pi^0)$ and $\Gamma(e^+e^-) B(\mu^+\mu^-)$ (AUBERT 04).
- ⁵ Mostly $\rho\pi$, see also $\rho\pi$ subsection.
- ⁶ Obtained comparing the rates for $\pi^+\pi^-\pi^0$ and $\mu^+\mu^-$, using J/ψ events produced via $\psi(2S) \rightarrow \pi^+\pi^- J/\psi$ and with $B(J/\psi \rightarrow \mu^+\mu^-) = 5.88 \pm 0.10\%$.



$\Gamma(K^+K^-\pi^0)/\Gamma(\pi^+\pi^-\pi^0)$ $\Gamma_{141}/\Gamma_{129}$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
12.0±0.3±0.9	23K	LEES	17C BABR	$J/\psi \rightarrow h^+h^-\pi^0$

$\Gamma(K_S^0 K^\pm \pi^\mp)/\Gamma(\pi^+\pi^-\pi^0)$ $\Gamma_{142}/\Gamma_{129}$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
26.5±0.5±2.1	24K	LEES	17C BABR	$J/\psi \rightarrow h^0 h^+ h^-$

$\Gamma(\pi^+\pi^-\pi^0 K^+K^-)/\Gamma_{\text{total}}$ Γ_{130}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.79±0.29 OUR AVERAGE				Error includes scale factor of 2.2.
1.93±0.14±0.05	768	¹ AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow K^+K^-\pi^+\pi^-\pi^0\gamma$
1.2 ±0.3	309	VANNUCCI	77 MRK1	e^+e^-

¹ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow \pi^+ \pi^- \pi^0 K^+ K^-)/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = 0.1070 \pm 0.0043 \pm 0.0064$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(4(\pi^+ \pi^-) \pi^0)/\Gamma_{\text{total}}$ Γ_{131}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
90 ± 30	13	JEAN-MARIE 76	MRK1	$e^+ e^-$

$\Gamma(\pi^+ \pi^- K^+ K^-)/\Gamma_{\text{total}}$ Γ_{132}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$6.5 \pm 0.4 \pm 0.2$	1.6k	¹ AUBERT 07AK BABR	10.6	$e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$
$6.1 \pm 0.7 \pm 0.2$	233	² AUBERT 05D BABR	10.6	$e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$
7.2 ± 2.3	205	VANNUCCI 77 MRK1		$e^+ e^-$

¹ Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \pi^+ \pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = (36.3 \pm 1.3 \pm 2.1) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Superseded by AUBERT 07AK. AUBERT 05D reports $[\Gamma(J/\psi(1S) \rightarrow \pi^+ \pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = (33.6 \pm 2.7 \pm 2.7) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\pi^+ \pi^- K^+ K^- \eta)/\Gamma_{\text{total}}$ Γ_{137}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.84 ± 0.28 ± 0.05	73	¹ AUBERT 07AU BABR	10.6	$e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \eta \gamma$

¹ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow \pi^+ \pi^- K^+ K^- \eta)/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = (10.2 \pm 1.3 \pm 0.8) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\pi^0 \pi^0 K^+ K^-)/\Gamma_{\text{total}}$ Γ_{138}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.45 \pm 0.31 \pm 0.06$	203 ± 16	¹ AUBERT 07AK BABR	10.6	$e^+ e^- \rightarrow \pi^0 \pi^0 K^+ K^- \gamma$
--------------------------	--------------	-------------------------------	------	--

¹ Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \pi^0 \pi^0 K^+ K^-)/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = (13.6 \pm 1.1 \pm 1.3) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K \bar{K} \pi)/\Gamma_{\text{total}}$ Γ_{140}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
61 ± 10 OUR AVERAGE				
55.2 ± 12.0	25	FRANKLIN 83	MRK2	$e^+ e^- \rightarrow K^+ K^- \pi^0$
78.0 ± 21.0	126	VANNUCCI 77	MRK1	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp$

$\Gamma(2(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{147}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

3.57±0.30 OUR AVERAGE

3.53±0.12±0.29	1107	¹ ABLIKIM	05H BES2	$e^+e^- \rightarrow \psi(2S) \rightarrow J/\psi\pi^+\pi^-, J/\psi \rightarrow 2(\pi^+\pi^-)$
4.0 ±1.0	76	JEAN-MARIE	76 MRK1	e^+e^-

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.51±0.34±0.09	270	² AUBERT	05D BABR	10.6 $e^+e^- \rightarrow 2(\pi^+\pi^-)\gamma$
----------------	-----	---------------------	----------	---

¹ Computed using $B(J/\psi \rightarrow \mu^+\mu^-) = 0.0588 \pm 0.0010$.² AUBERT 05D reports $[\Gamma(J/\psi(1S) \rightarrow 2(\pi^+\pi^-))/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+e^-)] = (19.5 \pm 1.4 \pm 1.3) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value. Superseded by LEES 12E. $\Gamma(3(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{148}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

43 ± 4 OUR AVERAGE

43.0 ± 2.9 ± 2.8	496	¹ AUBERT	06D BABR	10.6 $e^+e^- \rightarrow 3(\pi^+\pi^-)\gamma$
40 ± 20	32	JEAN-MARIE	76 MRK1	e^+e^-

¹ Using $\Gamma(J/\psi \rightarrow e^+e^-) = 5.52 \pm 0.14 \pm 0.04$ keV. $\Gamma(2(\pi^+\pi^-\pi^0))/\Gamma_{\text{total}}$ Γ_{149}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

1.62±0.09±0.19	761	¹ AUBERT	06D BABR	10.6 $e^+e^- \rightarrow 2(\pi^+\pi^-\pi^0)\gamma$
-----------------------	-----	---------------------	----------	--

¹ Using $\Gamma(J/\psi \rightarrow e^+e^-) = 5.52 \pm 0.14 \pm 0.04$ keV. $\Gamma(2(\pi^+\pi^-\eta))/\Gamma_{\text{total}}$ Γ_{150}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.29±0.24 OUR AVERAGE

2.35±0.39±0.20	85	¹ AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow 2(\pi^+\pi^-\eta)\gamma$
2.26±0.08±0.27	4839	ABLIKIM	05C BES2	$e^+e^- \rightarrow 2(\pi^+\pi^-\eta)$

¹ AUBERT 07AU quotes $\Gamma_{ee}^{J/\psi} \cdot B(J/\psi \rightarrow 2(\pi^+\pi^-\eta)) \cdot B(\eta \rightarrow \gamma\gamma) = 5.16 \pm 0.85 \pm 0.39$ eV. $\Gamma(3(\pi^+\pi^-\eta))/\Gamma_{\text{total}}$ Γ_{151}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

7.24±0.96±1.11	616	ABLIKIM	05C BES2	$e^+e^- \rightarrow 3(\pi^+\pi^-\eta)$
-----------------------	-----	---------	----------	--

 $\Gamma(p\bar{p})/\Gamma_{\text{total}}$ Γ_{152}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.121±0.029 OUR AVERAGE

2.112±0.004±0.031	314k	ABLIKIM	12C BES3	e^+e^-
2.19 ±0.16 ±0.06	317	¹ WU	06 BELL	$B^+ \rightarrow p\bar{p}K^+$
2.26 ±0.01 ±0.14	63316	BAI	04E BES2	$e^+e^- \rightarrow J/\psi$
1.97 ±0.22	99	BALDINI	98 FENI	e^+e^-
1.91 ±0.04 ±0.30		PALLIN	87 DM2	e^+e^-

2.16 ±0.07 ±0.15	1420	EATON	84	MRK2	e^+e^-
2.5 ±0.4	133	BRANDELIK	79C	DASP	e^+e^-
2.0 ±0.5		BESCH	78	BONA	e^+e^-
2.2 ±0.2	331	² PERUZZI	78	MRK1	e^+e^-
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
2.0 ±0.3	48	ANTONELLI	93	SPEC	e^+e^-

¹WU 06 reports $[\Gamma(J/\psi(1S) \rightarrow p\bar{p})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow J/\psi(1S)K^+)] = (2.21 \pm 0.13 \pm 0.10) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow J/\psi(1S)K^+) = (1.010 \pm 0.029) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

²Assuming angular distribution $(1+\cos^2\theta)$.


$\Gamma(p\bar{p}\pi^0)/\Gamma_{\text{total}}$ **Γ_{153}/Γ**

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.19±0.08 OUR AVERAGE	Error includes scale factor of 1.1.			
1.33±0.02±0.11	11k	ABLIKIM	09B	BES2 e^+e^-
1.13±0.09±0.09	685	EATON	84	MRK2 e^+e^-
1.4 ±0.4		BRANDELIK	79C	DASP e^+e^-
1.00±0.15	109	PERUZZI	78	MRK1 e^+e^-

$\Gamma(p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$ **Γ_{154}/Γ**

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
6.0 ±0.5 OUR AVERAGE	Error includes scale factor of 1.3. See the ideogram below.			
6.46±0.17±0.43	1435	EATON	84	MRK2 e^+e^-
3.8 ±1.6	48	BESCH	81	BONA e^+e^-
5.5 ±0.6	533	PERUZZI	78	MRK1 e^+e^-

WEIGHTED AVERAGE
6.0±0.5 (Error scaled by 1.3)



$\Gamma(\rho\bar{p}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{155}/Γ

Including $\rho\bar{p}\pi^+\pi^-\gamma$ and excluding ω, η, η'

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.3 ± 0.9 OUR AVERAGE		Error includes scale factor of 1.9.		
3.36 ± 0.65 ± 0.28	364	EATON	84	MRK2 e^+e^-
1.6 ± 0.6	39	PERUZZI	78	MRK1 e^+e^-

$\Gamma(\rho\bar{p}\eta)/\Gamma_{\text{total}}$ Γ_{156}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.00 ± 0.12 OUR AVERAGE				
1.91 ± 0.02 ± 0.17	13k	¹ ABLIKIM	09	BES2 e^+e^-
2.03 ± 0.13 ± 0.15	826	EATON	84	MRK2 e^+e^-
2.5 ± 1.2		BRANDELIK	79C	DASP e^+e^-
2.3 ± 0.4	197	PERUZZI	78	MRK1 e^+e^-

¹ From the combination of $\rho\bar{p}\eta \rightarrow \rho\bar{p}\gamma\gamma$ and $\rho\bar{p}\eta \rightarrow \rho\bar{p}\pi^+\pi^-\pi^0$ channels.


$\Gamma(\rho\bar{p}\rho)/\Gamma_{\text{total}}$ Γ_{157}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
< 0.31	90	EATON	84	MRK2 $e^+e^- \rightarrow \text{hadrons}\gamma$

$\Gamma(\rho\bar{p}\omega)/\Gamma_{\text{total}}$ Γ_{158}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.98 ± 0.10 OUR AVERAGE		Error includes scale factor of 1.3. See the ideogram below.		
0.90 ± 0.02 ± 0.09	2670	ABLIKIM	13P	BES3 e^+e^-
0.98 ± 0.03 ± 0.14	2449	ABLIKIM	08	BES2 e^+e^-
1.10 ± 0.17 ± 0.18	486	EATON	84	MRK2 e^+e^-
1.6 ± 0.3	77	PERUZZI	78	MRK1 e^+e^-

WEIGHTED AVERAGE
0.98 ± 0.10 (Error scaled by 1.3)



$\Gamma(\rho\bar{p}\eta'(958))/\Gamma_{\text{total}}$ Γ_{159}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.21 ± 0.04 OUR AVERAGE				
0.200 ± 0.023 ± 0.028	265 ± 31	¹ ABLIKIM	09 BES2	e^+e^-
0.68 ± 0.23 ± 0.17	19	EATON	84 MRK2	e^+e^-
1.8 ± 0.6	19	PERUZZI	78 MRK1	e^+e^-

¹From the combination of $p\bar{p}\eta' \rightarrow p\bar{p}\pi^+\pi^-\eta$ and $p\bar{p}\eta' \rightarrow p\bar{p}\gamma\rho^0$ channels.

 $\Gamma(\rho\bar{p}a_0(980) \rightarrow \rho\bar{p}\pi^0\eta)/\Gamma_{\text{total}}$ Γ_{160}/Γ

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
6.8 ± 1.2 ± 1.3	ABLIKIM 14N	BES3	$e^+e^- \rightarrow J/\psi$

 $\Gamma(\rho\bar{p}\phi)/\Gamma_{\text{total}}$ Γ_{161}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.519 ± 0.033 OUR AVERAGE				
0.523 ± 0.006 ± 0.033	14K	ABLIKIM 16K	BES3	$J/\psi \rightarrow \rho\bar{p}K_S^0 K_L^0,$ $\rho\bar{p}K^+ K^-$
0.45 ± 0.13 ± 0.07		FALVARD 88	DM2	$J/\psi \rightarrow \text{hadrons}$

 $\Gamma(n\bar{n})/\Gamma_{\text{total}}$ Γ_{162}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.09 ± 0.16 OUR AVERAGE				
2.07 ± 0.01 ± 0.17	36k	ABLIKIM 12C	BES3	e^+e^-
2.31 ± 0.49	79	BALDINI 98	FENI	e^+e^-
1.8 ± 0.9		BESCH 78	BONA	e^+e^-
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.90 ± 0.55	40	ANTONELLI 93	SPEC	e^+e^-

 $\Gamma(n\bar{n}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{163}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.8 ± 3.6	5	BESCH 81	BONA	e^+e^-

 $\Gamma(\Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}$ Γ_{164}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.50 ± 0.10 ± 0.22	399	ABLIKIM 08O	BES2	$e^+e^- \rightarrow J/\psi$

 $\Gamma(\Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}$ Γ_{165}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.172 ± 0.031 OUR AVERAGE				
Error includes scale factor of 1.4.				
1.164 ± 0.004 ± 0.023	111k	ABLIKIM 17L	BES3	$J/\psi \rightarrow \Sigma^0\bar{\Sigma}^0$
1.15 ± 0.24 ± 0.03		¹ AUBERT 07BD	BABR	10.6 $e^+e^- \rightarrow \Sigma^0\bar{\Sigma}^0\gamma$
1.33 ± 0.04 ± 0.11	1.7k	ABLIKIM 06	BES2	$J/\psi \rightarrow \Sigma^0\bar{\Sigma}^0$
1.06 ± 0.04 ± 0.23	884	PALLIN 87	DM2	$e^+e^- \rightarrow \Sigma^0\bar{\Sigma}^0$
1.58 ± 0.16 ± 0.25	90	EATON 84	MRK2	$e^+e^- \rightarrow \Sigma^0\bar{\Sigma}^0$
1.3 ± 0.4	52	PERUZZI 78	MRK1	$e^+e^- \rightarrow \Sigma^0\bar{\Sigma}^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.4 ± 2.6	3	BESCH 81	BONA	$e^+e^- \rightarrow \Sigma^+\bar{\Sigma}^-$

¹ AUBERT 07BD reports $[\Gamma(J/\psi(1S) \rightarrow \Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = (6.4 \pm 1.2 \pm 0.6) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2(\pi^+ \pi^-) K^+ K^-)/\Gamma_{\text{total}}$ Γ_{166}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
47 ± 7 OUR AVERAGE		Error includes scale factor of 1.3.		
49.8 ± 4.2 ± 3.4	205	¹ AUBERT	06D BABR	10.6 $e^+ e^- \rightarrow \omega K^+ K^- 2(\pi^+ \pi^-) \gamma$
31 ± 13	30	VANNUCCI	77 MRK1	$e^+ e^-$


¹ Using $\Gamma(J/\psi \rightarrow e^+ e^-) = 5.52 \pm 0.14 \pm 0.04$ keV.

$\Gamma(\rho \bar{n} \pi^-)/\Gamma_{\text{total}}$ Γ_{167}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.12 ± 0.09 OUR AVERAGE				
2.36 ± 0.02 ± 0.21	59k	ABLIKIM	06K BES2	$J/\psi \rightarrow \rho \pi^- \bar{n}$
2.47 ± 0.02 ± 0.24	55k	ABLIKIM	06K BES2	$J/\psi \rightarrow \bar{\rho} \pi^+ n$
2.02 ± 0.07 ± 0.16	1288	EATON	84 MRK2	$e^+ e^- \rightarrow \rho \pi^-$
1.93 ± 0.07 ± 0.16	1191	EATON	84 MRK2	$e^+ e^- \rightarrow \bar{\rho} \pi^+$
1.7 ± 0.7	32	BESCH	81 BONA	$e^+ e^- \rightarrow \rho \pi^-$
1.6 ± 1.2	5	BESCH	81 BONA	$e^+ e^- \rightarrow \bar{\rho} \pi^+$
2.16 ± 0.29	194	PERUZZI	78 MRK1	$e^+ e^- \rightarrow \rho \pi^-$
2.04 ± 0.27	204	PERUZZI	78 MRK1	$e^+ e^- \rightarrow \bar{\rho} \pi^+$

$\Gamma(\Xi^- \bar{\Xi}^+)/\Gamma_{\text{total}}$ Γ_{171}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.97 ± 0.08 OUR AVERAGE		Error includes scale factor of 1.4. See the ideogram below.		
1.040 ± 0.006 ± 0.074	43k	ABLIKIM	16L BES3	$J/\psi \rightarrow \Xi^- \bar{\Xi}^+$
0.90 ± 0.03 ± 0.18	961	ABLIKIM	12P BES2	$J/\psi \rightarrow \Xi^- \bar{\Xi}^+$
0.70 ± 0.06 ± 0.12	132	HENRARD	87 DM2	$e^+ e^- \rightarrow \Xi^- \bar{\Xi}^+$
1.14 ± 0.08 ± 0.20	194	EATON	84 MRK2	$e^+ e^- \rightarrow \Xi^- \bar{\Xi}^+$
1.4 ± 0.5	51	PERUZZI	78 MRK1	$e^+ e^- \rightarrow \Xi^- \bar{\Xi}^+$




$\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$					Γ_{172}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
1.89 ± 0.08	OUR AVERAGE	Error includes scale factor of 2.5. See the ideogram below.			
$1.943 \pm 0.003 \pm 0.033$	441k	ABLIKIM	17L BES3	e^+e^-	
$1.93 \pm 0.21 \pm 0.05$		¹ AUBERT	07BD BABR	$10.6 e^+e^- \rightarrow \Lambda\bar{\Lambda}\gamma$	
$2.03 \pm 0.03 \pm 0.15$	8887	ABLIKIM	06 BES2	$J/\psi \rightarrow \Lambda\bar{\Lambda}$	
$2.0^{+0.5}_{-0.4} \pm 0.1$	46	² WU	06 BELL	$B^+ \rightarrow \Lambda\bar{\Lambda}K^+$	
$1.08 \pm 0.06 \pm 0.24$	631	BAI	98G BES	e^+e^-	
$1.38 \pm 0.05 \pm 0.20$	1847	PALLIN	87 DM2	e^+e^-	
$1.58 \pm 0.08 \pm 0.19$	365	EATON	84 MRK2	e^+e^-	
2.6 ± 1.6	5	BESCH	81 BONA	e^+e^-	
1.1 ± 0.2	196	PERUZZI	78 MRK1	e^+e^-	

¹AUBERT 07BD reports $[\Gamma(J/\psi(1S) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+e^-)] = (10.7 \pm 0.9 \pm 0.7) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

²WU 06 reports $[\Gamma(J/\psi(1S) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow J/\psi(1S)K^+)] = (2.00^{+0.34}_{-0.29} \pm 0.34) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow J/\psi(1S)K^+) = (1.010 \pm 0.029) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

WEIGHTED AVERAGE
 1.89 ± 0.08 (Error scaled by 2.5)



$\Gamma(\Lambda\bar{\Lambda})/\Gamma(p\bar{p})$

$\Gamma_{172}/\Gamma_{152}$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.90^{+0.15}_{-0.14} \pm 0.10$	¹ WU	06 BELL	$B^+ \rightarrow p\bar{p}K^+, \Lambda\bar{\Lambda}K^+$

¹ Not independent of other $J/\psi \rightarrow \Lambda\bar{\Lambda}, p\bar{p}$ branching ratios reported by WU 06.

$\Gamma(\Lambda\bar{\Sigma}^- \pi^+ \text{ (or c.c.)})/\Gamma_{\text{total}}$

Γ_{173}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.83 ± 0.07 OUR AVERAGE				Error includes scale factor of 1.2.
$0.770 \pm 0.051 \pm 0.083$	335	¹ ABLIKIM	07H BES2	$e^+e^- \rightarrow \bar{\Lambda}\Sigma^+\pi^-$
$0.747 \pm 0.056 \pm 0.076$	254	¹ ABLIKIM	07H BES2	$e^+e^- \rightarrow \Lambda\bar{\Sigma}^-\pi^+$
$0.90 \pm 0.06 \pm 0.16$	225 ± 15	HENRARD	87 DM2	$e^+e^- \rightarrow \bar{\Lambda}\Sigma^+\pi^-$
$1.11 \pm 0.06 \pm 0.20$	342 ± 18	HENRARD	87 DM2	$e^+e^- \rightarrow \Lambda\bar{\Sigma}^-\pi^+$
$1.53 \pm 0.17 \pm 0.38$	135	EATON	84 MRK2	$e^+e^- \rightarrow \bar{\Lambda}\Sigma^+\pi^-$
$1.38 \pm 0.21 \pm 0.35$	118	EATON	84 MRK2	$e^+e^- \rightarrow \Lambda\bar{\Sigma}^-\pi^+$

¹ Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\Sigma^+ \rightarrow \pi^0 p) = 51.6\%$.

$\Gamma(pK^-\bar{\Lambda})/\Gamma_{\text{total}}$

Γ_{174}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$0.89 \pm 0.07 \pm 0.14$	307	EATON	84 MRK2	e^+e^-

$\Gamma(2(K^+ K^-))/\Gamma_{\text{total}}$ Γ_{175}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.74±0.07 OUR AVERAGE				
0.72±0.06±0.05	287 ± 24	LEES	12F	BABR 10.6 $e^+e^- \rightarrow 2(K^+ K^-)\gamma$
1.4 $^{+0.5}_{-0.4}$ ±0.2	11.0 $^{+4.3}_{-3.5}$	¹ HUANG	03	BELL $B^+ \rightarrow 2(K^+ K^-) K^+$
0.7 ±0.3		VANNUCCI	77	MRK1 e^+e^-
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.74±0.09±0.02	156 ± 15	² AUBERT	07AK	BABR 10.6 $e^+e^- \rightarrow 2(K^+ K^-)\gamma$
0.72±0.17±0.02	38	³ AUBERT	05D	BABR 10.6 $e^+e^- \rightarrow 2(K^+ K^-)\gamma$

¹ Using $B(B^+ \rightarrow J/\psi K^+) = (1.01 \pm 0.05) \times 10^{-3}$.

² Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow 2(K^+ K^-))/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+e^-)] = (4.11 \pm 0.39 \pm 0.30) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Superseded by AUBERT 07AK. AUBERT 05D reports $[\Gamma(J/\psi(1S) \rightarrow 2(K^+ K^-))/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+e^-)] = (4.0 \pm 0.7 \pm 0.6) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\rho K^- \bar{\Sigma}^0)/\Gamma_{\text{total}}$ Γ_{176}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.29±0.06±0.05	90	EATON	84	MRK2 e^+e^-

$\Gamma(K^+ K^-)/\Gamma_{\text{total}}$ Γ_{177}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.86±0.09±0.19	1k	¹ METREVELI	12	$\psi(2S) \rightarrow \pi^+ \pi^- K^+ K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
3.22±0.20±0.12	462	^{2,3} LEES	15J	BABR $e^+e^- \rightarrow K^+ K^- \gamma$
3.50±0.20±0.12	462	^{3,4} LEES	15J	BABR $e^+e^- \rightarrow K^+ K^- \gamma$
2.39±0.24±0.22	107	⁵ BALTRUSAIT..85D	MRK3	e^+e^-
2.2 ±0.9	6	⁵ BRANDELIK	79C	DASP e^+e^-

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

² $\sin\phi > 0$.

³ Using $\Gamma(J/\psi \rightarrow e^+e^-) = (5.55 \pm 0.14)$ keV.

⁴ $\sin\phi < 0$.


⁵ Interference with non-resonant $K^+ K^-$ production not taken into account.

$\Gamma(K_S^0 K_L^0)/\Gamma_{\text{total}}$ Γ_{178}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.95±0.11 OUR AVERAGE				Error includes scale factor of 2.4. See the ideogram below.
1.93±0.01±0.05	110K	ABLIKIM	17AH	BES3 $J/\psi \rightarrow K_S^0 K_L^0 \rightarrow \pi^+ \pi^- X$
2.62±0.15±0.14	0.3k	¹ METREVELI	12	$\psi(2S) \rightarrow \pi^+ \pi^- K_S^0 K_L^0$
1.82±0.04±0.13	2.1k	² BAI	04A	BES2 $J/\psi \rightarrow K_S^0 K_L^0 \rightarrow \pi^+ \pi^- X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.18±0.12±0.18		JOUSSET	90	DM2 $J/\psi \rightarrow$ hadrons
1.01±0.16±0.09	74	BALTRUSAIT..85D	MRK3	e^+e^-

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

² Using $B(K_S^0 \rightarrow \pi^+ \pi^-) = 0.6868 \pm 0.0027$.



$\Gamma(\Lambda \bar{\Lambda} \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{179}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
4.30±0.13±0.99	2.4k	ABLIKIM	12P BES2	J/ψ

$\Gamma(\Lambda \bar{\Lambda} \eta)/\Gamma_{\text{total}}$ Γ_{180}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
16.2±1.7 OUR AVERAGE				
15.7±0.80±1.54	454	¹ ABLIKIM	13F BES3	$J/\psi \rightarrow p \bar{p} \pi^+ \pi^- \gamma \gamma$
26.2±6.0 ±4.4	44	² ABLIKIM	07H BES2	$e^+ e^- \rightarrow \psi(2S)$

¹ Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\eta \rightarrow \gamma \gamma) = 39.31\%$.

² Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\eta \rightarrow \gamma \gamma) = 39.4\%$.

$\Gamma(\Lambda \bar{\Lambda} \pi^0)/\Gamma_{\text{total}}$ Γ_{181}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
3.78±0.27±0.30		323	¹ ABLIKIM	13F BES3	$J/\psi \rightarrow p \bar{p} \pi^+ \pi^- \gamma \gamma$

••• We do not use the following data for averages, fits, limits, etc. •••

< 6.4	90	² ABLIKIM	07H BES2	$e^+ e^- \rightarrow \psi(2S)$
23 ±7 ±8	11	BAI	98G BES	$e^+ e^-$
22 ±5 ±5	19	HENRARD	87 DM2	$e^+ e^-$

¹ Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\pi^0 \rightarrow \gamma \gamma) = 98.8\%$.

² Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$.

$\Gamma(\bar{\Lambda}nK_S^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{182}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
6.46 ± 0.20 ± 1.07	1058	¹ ABLIKIM	08C BES2	$e^+e^- \rightarrow J/\psi$

¹ Using $B(\bar{\Lambda} \rightarrow \bar{p}\pi^+) = 63.9\%$ and $B(K_S^0 \rightarrow \pi^+\pi^-) = 69.2\%$.

$\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{183}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.47 ± 0.14 OUR AVERAGE				
1.47 ± 0.13 ± 0.13	140	¹ METREVELI	12	$\psi(2S) \rightarrow 2(\pi^+\pi^-)$
1.58 ± 0.20 ± 0.15	84	BALTRUSAIT..85D	MRK3	e^+e^-
1.0 ± 0.5	5	BRANDELIK	78B DASP	e^+e^-
1.6 ± 1.6	1	VANNUCCI	77 MRK1	e^+e^-

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\Lambda\bar{\Sigma} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{184}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.83 ± 0.23 OUR AVERAGE					
2.74 ± 0.24 ± 0.22		234 ± 21	¹ ABLIKIM	12B BES3	$J/\psi \rightarrow \Lambda\bar{\Sigma}^0$
2.92 ± 0.22 ± 0.24		308 ± 24	² ABLIKIM	12B BES3	$J/\psi \rightarrow \bar{\Lambda}\Sigma^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<15	90	PERUZZI	78 MRK1	$e^+e^- \rightarrow \Lambda X$
-----	----	---------	---------	--------------------------------

¹ ABLIKIM 12B quotes $B(J/\psi \rightarrow \Lambda\bar{\Sigma}^0)$ which we multiply by 2.

² ABLIKIM 12B quotes $B(J/\psi \rightarrow \bar{\Lambda}\Sigma^0)$ which we multiply by 2.

$\Gamma(K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{185}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.4 × 10⁻⁸	95	¹ ABLIKIM	17AH BES3	$J/\psi \rightarrow K_S^0 K_S^0 \rightarrow \pi^+\pi^-\pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1 × 10 ⁻⁶	95	¹ BAI	04D BES	e^+e^-
-----------------------	----	------------------	---------	----------

<5.2 × 10 ⁻⁶	90	¹ BALTRUSAIT..85C	MRK3	e^+e^-
-------------------------	----	------------------------------	------	----------

¹ Forbidden by CP.

————— RADIATIVE DECAYS —————

$\Gamma(3\gamma)/\Gamma_{\text{total}}$ Γ_{186}/Γ

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
11.6 ± 2.2 OUR AVERAGE					
11.3 ± 1.8 ± 2.0		113 ± 18	ABLIKIM	13I BES3	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$
12 ± 3 ± 2		24.2 ^{+7.2} _{-6.0}	ADAMS	08 CLEO	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<55	90	PARTRIDGE	80 CBAL	e^+e^-
-----	----	-----------	---------	----------

$\Gamma(4\gamma)/\Gamma_{\text{total}}$ Γ_{187}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<9	90	ADAMS	08 CLEO	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$

$\Gamma(5\gamma)/\Gamma_{\text{total}}$ Γ_{188}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<15	90	ADAMS	08	CLEO $\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$

 $\Gamma(\gamma\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_{189}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
1.15 ± 0.05	¹ ABLIKIM	15AE BES3	$J/\psi \rightarrow \gamma\pi^0\pi^0$

¹ The uncertainty is systematic as statistical is negligible.

 $\Gamma(\gamma\eta\pi^0)/\Gamma_{\text{total}}$ Γ_{190}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
21.4 ± 1.8 ± 2.5	596	ABLIKIM	16P	BES3 $J/\psi \rightarrow 5\gamma$

 $\Gamma(\gamma a_0(980)^0 \rightarrow \gamma\eta\pi^0)/\Gamma_{\text{total}}$ Γ_{191}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2.5 × 10⁻⁶	95	ABLIKIM	16P	BES3 $J/\psi \rightarrow 5\gamma$

 $\Gamma(\gamma a_2(1320)^0 \rightarrow \gamma\eta\pi^0)/\Gamma_{\text{total}}$ Γ_{192}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<6.6 × 10⁻⁶	95	ABLIKIM	16P	BES3 $J/\psi \rightarrow 5\gamma$

 $\Gamma(\gamma\eta_c(1S))/\Gamma_{\text{total}}$ Γ_{193}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.7 ± 0.4 OUR AVERAGE		Error includes scale factor of 1.5.		
2.00 ± 0.31 ± 0.02		¹ MITCHELL	09	CLEO $e^+ e^- \rightarrow \gamma X$
1.27 ± 0.36		GAISER	86	CBAL $J/\psi \rightarrow \gamma X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
seen		ANASHIN	14	KEDR $J/\psi \rightarrow \gamma\eta_c$
0.79 ± 0.20	273 ± 43	² AUBERT	06E	BABR $B^\pm \rightarrow K^\pm X_{c\bar{c}}$
seen	16	BALTRUSAITIS	84	MRK3 $J/\psi \rightarrow 2\phi\gamma$

¹ MITCHELL 09 reports $(1.98 \pm 0.09 \pm 0.30) \times 10^{-2}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)]$ assuming $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (35.04 \pm 0.07 \pm 0.77) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.67 \pm 0.30) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Calculated by the authors using an average of $B(J/\psi \rightarrow \gamma\eta_c) \times B(\eta_c \rightarrow K\bar{K}\pi)$ from BALTRUSAITIS 86, BISELLO 91, BAI 04 and $B(\eta_c \rightarrow K\bar{K}\pi) = (8.5 \pm 1.8)\%$ from AUBERT 06E.

 $\Gamma(\gamma\eta_c(1S) \rightarrow 3\gamma)/\Gamma_{\text{total}}$ Γ_{194}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
3.8^{+1.3}_{-1.0} OUR AVERAGE		Error includes scale factor of 1.1.		
4.5 ± 1.2 ± 0.6	33 ± 9	ABLIKIM	13I	BES3 $\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$
1.2 ^{+2.7} _{-1.1} ± 0.3	1.2 ^{+2.8} _{-1.1}	ADAMS	08	CLEO $\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$

$\Gamma(\gamma\pi^+\pi^-2\pi^0)/\Gamma_{\text{total}}$ Γ_{195}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$8.3 \pm 0.2 \pm 3.1$	¹ BALTRUSAIT ..86B	MRK3	$J/\psi \rightarrow 4\pi\gamma$
¹ 4π mass less than 2.0 GeV.			

$\Gamma(\gamma\eta\pi\pi)/\Gamma_{\text{total}}$ Γ_{196}/Γ


VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
6.1 ± 1.0 OUR AVERAGE			
$5.85 \pm 0.3 \pm 1.05$	¹ EDWARDS	83B	CBAL $J/\psi \rightarrow \eta\pi^+\pi^-$
$7.8 \pm 1.2 \pm 2.4$	¹ EDWARDS	83B	CBAL $J/\psi \rightarrow \eta 2\pi^0$
¹ Broad enhancement at 1700 MeV.			

$\Gamma(\gamma\eta_2(1870) \rightarrow \gamma\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{197}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$6.2 \pm 2.2 \pm 0.9$	BAI	99	BES $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$

$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma K\bar{K}\pi)/\Gamma_{\text{total}}$ Γ_{198}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
2.8 ± 0.6 OUR AVERAGE	Error includes scale factor of 1.6. See the ideogram below.		
$1.66 \pm 0.1 \pm 0.58$	^{1,2} BAI	00D	BES $J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$
$3.8 \pm 0.3 \pm 0.6$	³ AUGUSTIN	90	DM2 $J/\psi \rightarrow \gamma K\bar{K}\pi$
$4.0 \pm 0.7 \pm 1.0$	³ EDWARDS	82E	CBAL $J/\psi \rightarrow K^+ K^- \pi^0 \gamma$
4.3 ± 1.7	^{3,4} SCHARRE	80	MRK2 $e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$1.78 \pm 0.21 \pm 0.33$	^{3,5,6} AUGUSTIN	92	DM2 $J/\psi \rightarrow \gamma K\bar{K}\pi$
$0.83 \pm 0.13 \pm 0.18$	^{3,7,8} AUGUSTIN	92	DM2 $J/\psi \rightarrow \gamma K\bar{K}\pi$
$0.66^{+0.17+0.24}_{-0.16-0.15}$	^{3,6,9} BAI	90C	MRK3 $J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
$1.03^{+0.21+0.26}_{-0.18-0.19}$	^{3,8,10} BAI	90C	MRK3 $J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$



- ¹ Interference with the $J/\psi(1S)$ radiative transition to the broad $K\bar{K}\pi$ pseudoscalar state around 1800 is $(0.15 \pm 0.01 \pm 0.05) \times 10^{-3}$.
² Interference with $J/\psi \rightarrow \gamma f_1(1420)$ is $(-0.03 \pm 0.01 \pm 0.01) \times 10^{-3}$.
³ Includes unknown branching fraction $\eta(1405) \rightarrow K\bar{K}\pi$.
⁴ Corrected for spin-zero hypothesis for $\eta(1405)$.
⁵ From fit to the $a_0(980)\pi 0^{-+}$ partial wave.
⁶ $a_0(980)\pi$ mode.
⁷ From fit to the $K^*(892)K 0^{-+}$ partial wave.
⁸ K^*K mode.
⁹ From $a_0(980)\pi$ final state.
¹⁰ From $K^*(890)K$ final state.

$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\gamma\rho^0)/\Gamma_{\text{total}}$ Γ_{199}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
0.78 ± 0.20 OUR AVERAGE	Error includes scale factor of 1.8.		
1.07 ± 0.17 ± 0.11	¹ BAI	04J	BES2 $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
0.64 ± 0.12 ± 0.07	¹ COFFMAN	90	MRK3 $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
¹ Includes unknown branching fraction $\eta(1405) \rightarrow \gamma\rho^0$.			

$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{200}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.0 ± 0.5 OUR AVERAGE				
2.6 ± 0.7 ± 0.4		BAI	99	BES $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
3.38 ± 0.33 ± 0.64		¹ BOLTON	92B	MRK3 $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
7.0 ± 0.6 ± 1.1	261	² AUGUSTIN	90	DM2 $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
¹ Via $a_0(980)\pi$.				
² Includes unknown branching fraction to $\eta\pi^+\pi^-$.				

$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\gamma\phi)/\Gamma_{\text{total}}$ Γ_{201}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.82	95	BAI	04J	BES2 $J/\psi \rightarrow \gamma\gamma K^+K^-$

$\Gamma(\gamma\rho\rho)/\Gamma_{\text{total}}$ Γ_{202}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
4.5 ± 0.8 OUR AVERAGE				
4.7 ± 0.3 ± 0.9		¹ BALTRUSAIT..	86B	MRK3 $J/\psi \rightarrow 4\pi\gamma$
3.75 ± 1.05 ± 1.20		² BURKE	82	MRK2 $J/\psi \rightarrow 4\pi\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.09	90	³ BISELLO	89B	$J/\psi \rightarrow 4\pi\gamma$
¹ 4π mass less than 2.0 GeV.				
² 4π mass less than 2.0 GeV. We have multiplied $2\rho^0$ measurement by 3 to obtain 2ρ .				
³ 4π mass in the range 2.0–25 GeV.				


$\Gamma(\gamma\rho\omega)/\Gamma_{\text{total}}$ Γ_{203}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<5.4	90	ABLIKIM	08A	BES2 $e^+e^- \rightarrow J/\psi$

$\Gamma(\gamma\rho\phi)/\Gamma_{\text{total}}$		Γ_{204}/Γ			
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	
<8.8	90	ABLIKIM	08A	BES2	$e^+e^- \rightarrow J/\psi$

$\Gamma(\gamma\eta'(958))/\Gamma_{\text{total}}$		Γ_{205}/Γ			
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
5.13±0.17 OUR AVERAGE	Error	includes scale factor	of 1.3.	See the ideogram below.	
4.78±0.22±0.08		¹ ABLIKIM	11	BES3	$J/\psi \rightarrow \eta'\gamma$
5.24±0.12±0.11		PEDLAR	09	CLE3	$J/\psi \rightarrow \eta'\gamma$
5.55±0.44	35k	ABLIKIM	06E	BES2	$J/\psi \rightarrow \eta'\gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
4.50±0.14±0.53		BOLTON	92B	MRK3	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta, \eta \rightarrow \gamma\gamma$
4.30±0.31±0.71		BOLTON	92B	MRK3	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta, \eta \rightarrow \pi^+\pi^-\pi^0$
4.04±0.16±0.85	622	AUGUSTIN	90	DM2	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
4.39±0.09±0.66	2420	AUGUSTIN	90	DM2	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
4.1 ±0.3 ±0.6		BLOOM	83	CBAL	$e^+e^- \rightarrow 3\gamma + \text{hadrons}$
2.9 ±1.1	6	BRANDELIK	79c	DASP	$e^+e^- \rightarrow 3\gamma$
2.4 ±0.7	57	BARTEL	76	CNTR	$e^+e^- \rightarrow 2\gamma\rho$


¹ ABLIKIM 11 reports $(4.84 \pm 0.03 \pm 0.24) \times 10^{-3}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta'(958))/\Gamma_{\text{total}}] / [B(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)]$ assuming $B(\eta'(958) \rightarrow \pi^+\pi^-\eta) = (43.2 \pm 0.7) \times 10^{-2}$, $B(\eta \rightarrow 2\gamma) = (39.31 \pm 0.20) \times 10^{-2}$, which we rescale to our best values $B(\eta'(958) \rightarrow \pi^+\pi^-\eta) = (42.6 \pm 0.7) \times 10^{-2}$, $B(\eta \rightarrow 2\gamma) = (39.41 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.



$\Gamma(\gamma 2\pi^+ 2\pi^-)/\Gamma_{\text{total}}$ Γ_{206}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
2.8 ± 0.5 OUR AVERAGE	Error includes scale factor of 1.9. See the ideogram below.		
4.32 ± 0.14 ± 0.73	¹ BISELLO	89B DM2	$J/\psi \rightarrow 4\pi\gamma$
2.08 ± 0.13 ± 0.35	² BISELLO	89B DM2	$J/\psi \rightarrow 4\pi\gamma$
3.05 ± 0.08 ± 0.45	² BALTRUSAIT..86B	MRK3	$J/\psi \rightarrow 4\pi\gamma$
4.85 ± 0.45 ± 1.20	³ BURKE	82 MRK2	e^+e^-

- ¹ 4π mass less than 3.0 GeV.
- ² 4π mass less than 2.0 GeV.
- ³ 4π mass less than 2.5 GeV.



$\Gamma(\gamma f_2(1270) f_2(1270))/\Gamma_{\text{total}}$ Γ_{207}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
9.5 ± 0.7 ± 1.6	646 ± 45	ABLIKIM	04M BES	$J/\psi \rightarrow \gamma 2\pi^+ 2\pi^-$

$\Gamma(\gamma f_2(1270) f_2(1270) \text{ (non resonant)})/\Gamma_{\text{total}}$ Γ_{208}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
8.2 ± 0.8 ± 1.7	¹ ABLIKIM	04M BES	$J/\psi \rightarrow \gamma 2\pi^+ 2\pi^-$

¹ Subtracting contribution from intermediate $\eta_c(1S)$ decays.

$\Gamma(\gamma K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{209}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.1 ± 0.1 ± 0.6	1516	BAI	00B BES	$J/\psi \rightarrow \gamma K^+ K^0 \pi^+ \pi^-$

$\Gamma(\gamma f_4(2050))/\Gamma_{\text{total}}$ Γ_{210}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
2.7 ± 0.5 ± 0.5	¹ BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^-$

¹ Assuming branching fraction $f_4(2050) \rightarrow \pi \pi / \text{total} = 0.167$.

$\Gamma(\gamma \omega \omega)/\Gamma_{\text{total}}$ Γ_{211}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.61 ± 0.33 OUR AVERAGE				
6.0 ± 4.8 ± 1.8		ABLIKIM	08A BES2	$J/\psi \rightarrow \gamma \omega \pi^+ \pi^-$
1.41 ± 0.2 ± 0.42	120 ± 17	BISELLO	87 SPEC	$e^+ e^-$, hadrons γ
1.76 ± 0.09 ± 0.45		BALTRUSAIT..85C	MRK3	$e^+ e^- \rightarrow \text{hadrons } \gamma$

$\Gamma(\gamma \eta(1405/1475) \rightarrow \gamma \rho^0 \rho^0)/\Gamma_{\text{total}}$ Γ_{212}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
1.7 ± 0.4 OUR AVERAGE Error includes scale factor of 1.3.			
2.1 ± 0.4	BUGG	95 MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$
1.36 ± 0.38	^{1,2} BISELLO	89B DM2	$J/\psi \rightarrow 4\pi \gamma$

¹ Estimated by us from various fits.

² Includes unknown branching fraction to $\rho^0 \rho^0$.

$\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$ Γ_{213}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.64 ± 0.12 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.				
2.07 ± 0.16 ^{+0.02} / _{-0.07}	2.4k	^{1,2} DOBBS	15	$J/\psi \rightarrow \gamma \pi \pi$
1.63 ± 0.26 ^{+0.02} / _{-0.06}		³ ABLIKIM	06V BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^+ \pi^-$
1.42 ± 0.21 ^{+0.01} / _{-0.05}		⁴ ABLIKIM	06V BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^0 \pi^0$
1.33 ± 0.05 ± 0.20		⁵ AUGUSTIN	87 DM2	$J/\psi \rightarrow \gamma \pi^+ \pi^-$
1.36 ± 0.09 ± 0.23		⁵ BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^-$
1.48 ± 0.25 ± 0.30	178	EDWARDS	82B CBAL	$e^+ e^- \rightarrow 2\pi^0 \gamma$
2.0 ± 0.7	35	ALEXANDER	78 PLUT	$e^+ e^-$
1.2 ± 0.6	30	⁶ BRANDELIK	78B DASP	$e^+ e^- \rightarrow \pi^+ \pi^- \gamma$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.


² DOBBS 15 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma f_2(1270))/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi \pi)] = (1.744 \pm 0.052 \pm 0.122) \times 10^{-3}$ which we divide by our best value $B(f_2(1270) \rightarrow \pi \pi) = (84.2^{+2.9}_{-0.9}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ ABLIKIM 06V reports $[\Gamma(J/\psi(1S) \rightarrow \gamma f_2(1270))/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi \pi)] = (1.371 \pm 0.010 \pm 0.222) \times 10^{-3}$ which we divide by our best value $B(f_2(1270) \rightarrow \pi \pi) = (84.2^{+2.9}_{-0.9}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ ABLIKIM 06V reports $[\Gamma(J/\psi(1S) \rightarrow \gamma f_2(1270))/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi \pi)] = (1.200 \pm 0.027 \pm 0.174) \times 10^{-3}$ which we divide by our best value $B(f_2(1270) \rightarrow \pi \pi) = (84.2^{+2.9}_{-0.9}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁵ Estimated using $B(f_2(1270) \rightarrow \pi \pi) = 0.843 \pm 0.012$. The errors do not contain the uncertainty in the $f_2(1270)$ decay.

⁶ Restated by us to take account of spread of E1, M2, E3 transitions.



$\Gamma(\gamma f_0(1370) \rightarrow \gamma K \bar{K})/\Gamma_{\text{total}}$ Γ_{214}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	COMMENT
4.19±0.73±1.34	478	¹ DOBBS 15	$J/\psi \rightarrow \gamma K \bar{K}$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\gamma f_0(1710) \rightarrow \gamma K \bar{K})/\Gamma_{\text{total}}$ Γ_{215}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------


10.0 \pm 1.1 \pm 0.9 OUR AVERAGE Error includes scale factor of 1.5. See the ideogram below.

11.76±	0.54±0.94	1.2k	¹ DOBBS	15	$J/\psi \rightarrow \gamma K \bar{K}$
9.62±0.29	+3.51 -1.86		² BAI	03G BES	$J/\psi \rightarrow \gamma K \bar{K}$
5.0 ± 0.8	+1.8 -0.4		^{3,4} BAI	96C BES	$J/\psi \rightarrow \gamma K^+ K^-$
9.2 ± 1.4	±1.4		⁴ AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K^+ K^-$
10.4 ± 1.2	±1.6		⁴ AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
9.6 ± 1.2	±1.8		⁴ BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.6 ± 0.2	+0.6 -0.2		^{4,5} BAI	96C BES	$J/\psi \rightarrow \gamma K^+ K^-$
< 0.8		90	⁶ BISELLO	89B	$J/\psi \rightarrow 4\pi\gamma$
1.6 ± 0.4	±0.3		⁷ BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^-$
3.8 ± 1.6			⁸ EDWARDS	82D CBAL	$e^+ e^- \rightarrow \eta \eta \gamma$

- ¹ Using CLEO-c data but not authored by the CLEO Collaboration.
- ² Includes unknown branching ratio to $K^+ K^-$ or $K_S^0 K_S^0$.
- ³ Assuming $J^P = 2^+$ for $f_0(1710)$.
- ⁴ Includes unknown branching fraction to $K^+ K^-$ or $K_S^0 K_S^0$. We have multiplied $K^+ K^-$ measurement by 2, and $K_S^0 K_S^0$ by 4 to obtain $K\bar{K}$ result.
- ⁵ Assuming $J^P = 0^+$ for $f_0(1710)$.
- ⁶ Includes unknown branching fraction to $\rho^0 \rho^0$.
- ⁷ Includes unknown branching fraction to $\pi^+ \pi^-$.
- ⁸ Includes unknown branching fraction to $\eta\eta$.



$\Gamma(\gamma f_0(1710) \rightarrow \gamma \pi \pi) / \Gamma_{\text{total}}$ Γ_{216} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.8 ± 0.5				OUR AVERAGE
3.72 ± 0.30 ± 0.43	483	¹ DOBBS	15	$J/\psi \rightarrow \gamma \pi \pi$
3.96 ± 0.06 ± 1.12		² ABLIKIM	06V BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^+ \pi^-$
3.99 ± 0.15 ± 2.64		² ABLIKIM	06V BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^0 \pi^0$
• • •				We do not use the following data for averages, fits, limits, etc. • • •
2.5 ± 1.6 ± 0.8		BAI	98H BES	$J/\psi \rightarrow \gamma \pi^0 \pi^0$

- ¹ Using CLEO-c data but not authored by the CLEO Collaboration.
- ² Including unknown branching fraction to $\pi\pi$.

$\Gamma(\gamma f_0(1710) \rightarrow \gamma \omega \omega) / \Gamma_{\text{total}}$ Γ_{217} / Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.31 ± 0.06 ± 0.08	180	ABLIKIM	06H BES	$J/\psi \rightarrow \gamma \omega \omega$

$\Gamma(\gamma f_0(1710) \rightarrow \gamma \eta \eta) / \Gamma_{\text{total}}$ Γ_{218} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.35^{+0.13+1.24}_{-0.11-0.74}$	5.5k	¹ ABLIKIM	13N BES3	$J/\psi \rightarrow \gamma \eta \eta$

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

 $\Gamma(\gamma \eta) / \Gamma_{\text{total}}$ Γ_{219} / Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.104 ± 0.034 OUR AVERAGE				

$1.101 \pm 0.029 \pm 0.022$		PEDLAR	09 CLE3	$J/\psi \rightarrow \eta \gamma$
1.123 ± 0.089	11k	ABLIKIM	06E BES2	$J/\psi \rightarrow \eta \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.88 \pm 0.08 \pm 0.11$		BLOOM	83 CBAL	$e^+ e^-$
0.82 ± 0.10		BRANDELIK	79C DASP	$e^+ e^-$
1.3 ± 0.4	21	BARTEL	77 CNTR	$e^+ e^-$

 $\Gamma(\gamma f_1(1420) \rightarrow \gamma K \bar{K} \pi) / \Gamma_{\text{total}}$ Γ_{220} / Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.79 ± 0.13 OUR AVERAGE			

$0.68 \pm 0.04 \pm 0.24$	BAI	00D BES	$J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$
$0.76 \pm 0.15 \pm 0.21$	^{1,2} AUGUSTIN	92 DM2	$J/\psi \rightarrow \gamma K \bar{K} \pi$
$0.87 \pm 0.14^{+0.14}_{-0.11}$	¹ BAI	90C MRK3	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$

¹ Included unknown branching fraction $f_1(1420) \rightarrow K \bar{K} \pi$.

² From fit to the $K^*(892) K 1^{++}$ partial wave.

 $\Gamma(\gamma f_1(1285)) / \Gamma_{\text{total}}$ Γ_{221} / Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.61 ± 0.08 OUR AVERAGE			

$0.69 \pm 0.16 \pm 0.20$	¹ BAI	04J BES2	$J/\psi \rightarrow \gamma \gamma \rho^0$
$0.61 \pm 0.04 \pm 0.21$	² BAI	00D BES	$J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$
$0.45 \pm 0.09 \pm 0.17$	³ BAI	99 BES	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
$0.625 \pm 0.063 \pm 0.103$	⁴ BOLTON	92 MRK3	$J/\psi \rightarrow \gamma f_1(1285)$
$0.70 \pm 0.08 \pm 0.16$	⁵ BOLTON	92B MRK3	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$

¹ Assuming $B(f_1(1285) \rightarrow \rho^0 \gamma) = 0.055 \pm 0.013$.

² Assuming $\Gamma(f_1(1285) \rightarrow K \bar{K} \pi) / \Gamma_{\text{total}} = 0.090 \pm 0.004$.

³ Assuming $\Gamma(f_1(1285) \rightarrow \eta \pi \pi) / \Gamma_{\text{total}} = 0.5 \pm 0.18$.

⁴ Obtained summing the sequential decay channels

$$B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow \pi \pi \pi \pi) = (1.44 \pm 0.39 \pm 0.27) \times 10^{-4};$$

$$B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow a_0(980) \pi, a_0(980) \rightarrow \eta \pi) = (3.90 \pm 0.42 \pm 0.87) \times 10^{-4};$$

$$B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow a_0(980) \pi, a_0(980) \rightarrow K \bar{K}) = (0.66 \pm 0.26 \pm 0.29) \times 10^{-4};$$

$$B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow \gamma \rho^0) = (0.25 \pm 0.07 \pm 0.03) \times 10^{-4}.$$

⁵ Using $B(f_1(1285) \rightarrow a_0(980) \pi) = 0.37$, and including unknown branching ratio for $a_0(980) \rightarrow \eta \pi$.

$\Gamma(\gamma f_1(1510) \rightarrow \gamma \eta \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{222} / Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$4.5 \pm 1.0 \pm 0.7$	BAI	99	BES $J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$

$\Gamma(\gamma f'_2(1525)) / \Gamma_{\text{total}}$ Γ_{223} / Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------


$5.7^{+0.8}_{-0.5}$ OUR AVERAGE Error includes scale factor of 1.5. See the ideogram below.

$8.0 \pm 0.9 \pm 0.2$	750	1,2	DOBBS	15	$J/\psi \rightarrow \gamma K \bar{K}$
$3.85 \pm 0.17^{+1.91}_{-0.73}$		3	BAI	03G	BES $J/\psi \rightarrow \gamma K \bar{K}$
$3.6 \pm 0.4^{+1.4}_{-0.4}$		3	BAI	96C	BES $J/\psi \rightarrow \gamma K^+ K^-$
$5.6 \pm 1.4 \pm 0.9$		3	AUGUSTIN	88	DM2 $J/\psi \rightarrow \gamma K^+ K^-$
$4.5 \pm 0.4 \pm 0.9$		3	AUGUSTIN	88	DM2 $J/\psi \rightarrow \gamma K_S^0 K_S^0$
$6.8 \pm 1.6 \pm 1.4$		3	BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<3.4	90	4	4	BRANDELIK	79C	DASP	$e^+ e^- \rightarrow \pi^+ \pi^- \gamma$
<2.3	90	3		ALEXANDER	78	PLUT	$e^+ e^- \rightarrow K^+ K^- \gamma$

WEIGHTED AVERAGE
 $5.7 \pm 0.8 \pm 0.5$ (Error scaled by 1.5)



¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² DOBBS 15 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma f'_2(1525)) / \Gamma_{\text{total}}] \times [B(f'_2(1525) \rightarrow K \bar{K})] = (7.09 \pm 0.46 \pm 0.67) \times 10^{-4}$ which we divide by our best value $B(f'_2(1525) \rightarrow K \bar{K}) = (88.7 \pm 2.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Using $B(f'_2(1525) \rightarrow K\bar{K}) = 0.888$.

⁴ Assuming isotropic production and decay of the $f'_2(1525)$ and isospin.

$\Gamma(\gamma f'_2(1525) \rightarrow \gamma\eta\eta)/\Gamma_{\text{total}}$					Γ_{224}/Γ
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	
$3.42^{+0.43+1.37}_{-0.51-1.30}$	5.5k	¹ ABLIKIM	13N BES3	$J/\psi \rightarrow \gamma\eta\eta$	

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

$\Gamma(\gamma f_2(1640) \rightarrow \gamma\omega\omega)/\Gamma_{\text{total}}$					Γ_{225}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
$0.28 \pm 0.05 \pm 0.17$	141	ABLIKIM	06H BES	$J/\psi \rightarrow \gamma\omega\omega$	

$\Gamma(\gamma f_2(1910) \rightarrow \gamma\omega\omega)/\Gamma_{\text{total}}$					Γ_{226}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
$0.20 \pm 0.04 \pm 0.13$	151	ABLIKIM	06H BES	$J/\psi \rightarrow \gamma\omega\omega$	

$\Gamma(\gamma f_0(1800) \rightarrow \gamma\omega\phi)/\Gamma_{\text{total}}$					Γ_{227}/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
2.5 \pm 0.6 OUR AVERAGE					
$2.00 \pm 0.08^{+1.38}_{-1.64}$	1.3k	ABLIKIM	13J BES3	$J/\psi \rightarrow \gamma\omega\phi$	
$2.61 \pm 0.27 \pm 0.65$	95	ABLIKIM	06J BES2	$J/\psi \rightarrow \gamma\omega\phi$	

$\Gamma(\gamma f_2(1810) \rightarrow \gamma\eta\eta)/\Gamma_{\text{total}}$					Γ_{228}/Γ
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	COMMENT		
$5.40^{+0.60+3.42}_{-0.67-2.35}$	5.5k	¹ ABLIKIM	13N	$J/\psi \rightarrow \gamma\eta\eta$	

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.


$\Gamma(\gamma f_2(1950) \rightarrow \gamma K^*(892)\bar{K}^*(892))/\Gamma_{\text{total}}$					Γ_{229}/Γ
VALUE (units 10^{-3})		DOCUMENT ID	TECN	COMMENT	
$0.7 \pm 0.1 \pm 0.2$		BAI	00B BES	$J/\psi \rightarrow \gamma K^+ K^0 \pi^+ \pi^-$	

$\Gamma(\gamma K^*(892)\bar{K}^*(892))/\Gamma_{\text{total}}$					Γ_{230}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
$4.0 \pm 0.3 \pm 1.3$	320	¹ BAI	00B BES	$J/\psi \rightarrow \gamma K^+ K^0 \pi^+ \pi^-$	

¹ Summed over all charges.

$\Gamma(\gamma\phi\phi)/\Gamma_{\text{total}}$					Γ_{231}/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
4.0 \pm 1.2 OUR AVERAGE				Error includes scale factor of 2.1. See the ideogram below.	
$7.5 \pm 0.6 \pm 1.2$	168	BAI	90B MRK3	$J/\psi \rightarrow \gamma 4K$	
$3.4 \pm 0.8 \pm 0.6$	33 ± 7	¹ BISELLO	90 DM2	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$	
$3.1 \pm 0.7 \pm 0.4$		¹ BISELLO	86B DM2	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$	

¹ $\phi\phi$ mass less than 2.9 GeV, η_c excluded.



$\Gamma(\gamma\rho\rho)/\Gamma_{\text{total}}$					Γ_{232}/Γ
VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$0.38 \pm 0.07 \pm 0.07$		49	EATON	84	MRK2 e^+e^-
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.11	90		PERUZZI	78	MRK1 e^+e^-

$\Gamma(\gamma\eta(2225))/\Gamma_{\text{total}}$					Γ_{233}/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
$3.14^{+0.50}_{-0.19}$ OUR AVERAGE					
$2.40 \pm 0.10^{+2.47}_{-0.18}$	1,2	ABLIKIM	16N BES3	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$	
$4.4 \pm 0.4 \pm 0.8$	196	2 ABLIKIM	08I BES	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$	
$3.3 \pm 0.8 \pm 0.5$		2 BAI	90B MRK3	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$	
$2.7 \pm 0.6 \pm 0.6$		2 BAI	90B MRK3	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$	
$2.4^{+1.5}_{-1.0}$		3,4 BISELLO	89B DM2	$J/\psi \rightarrow 4\pi\gamma$	

¹ From a partial wave analysis of $J/\psi \rightarrow \gamma\phi\phi$ that also finds significant signals for for $\eta(2100)$, 0^-+ phase space, $f_0(2100)$, $f_2(2010)$, $f_2(2300)$, $f_2(2340)$, and a previously unseen 0^-+ state $X(2500)$ ($M = 2470^{+15+101}_{-19-23}$ MeV, $\Gamma = 230^{+64+56}_{-35-33}$ MeV).

² Includes unknown branching fraction to $\phi\phi$.

³ Estimated by us from various fits.

⁴ Includes unknown branching fraction to $\rho^0\rho^0$.

$$\Gamma(\gamma\eta(1760) \rightarrow \gamma\rho^0\rho^0)/\Gamma_{\text{total}} \quad \Gamma_{234}/\Gamma$$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.13±0.09	^{1,2} BISELLO	89B	DM2 $J/\psi \rightarrow 4\pi\gamma$

¹ Estimated by us from various fits.

² Includes unknown branching fraction to $\rho^0\rho^0$.

$$\Gamma(\gamma\eta(1760) \rightarrow \gamma\omega\omega)/\Gamma_{\text{total}} \quad \Gamma_{235}/\Gamma$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.98±0.08±0.32	1045	ABLIKIM	06H	BES $J/\psi \rightarrow \gamma\omega\omega$

$$\Gamma(\gamma X(1835) \rightarrow \gamma\pi^+\pi^-\eta')/\Gamma_{\text{total}} \quad \Gamma_{236}/\Gamma$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.77^{+0.34}_{-0.40} OUR AVERAGE				Error includes scale factor of 1.1.

3.93±0.38^{+0.31}_{-0.84} ¹ ABLIKIM 16J BES3 $J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$

2.87±0.09^{+0.49}_{-0.52} 4265 ² ABLIKIM 11C BES3 $J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$

2.2 ±0.4 ±0.4 264 ABLIKIM 05R BES2 $J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$

¹ From a fit of the measured $\pi^+\pi^-\eta'$ lineshape that accounts for the abrupt distortion observed at the $p\bar{p}$ threshold with a Flatte formula in addition to known backgrounds and contributors, as well as an *ad hoc* Breit-Wigner ($M \approx 1919$ MeV; $\Gamma \approx 51$ MeV) that is required for a good fit. Another explanation for the distortion provided by ABLIKIM 16J is that a second resonance near 1870 MeV interferes with the X(1835); fits to this possibility yield product branching fraction values compatible with that shown within the respective systematic uncertainties.

² From a fit of the $\pi^+\pi^-\eta'$ mass distribution to a combination of $\gamma f_1(1510)$, $\gamma X(1835)$, and two unconfirmed states $\gamma X(2120)$, and $\gamma X(2370)$, for $M(p\bar{p}) < 2.8$ GeV, and accounting for backgrounds from non- η' events and $J/\psi \rightarrow \pi^0\pi^+\pi^-\eta'$.

$$\Gamma(\gamma X(1835) \rightarrow \gamma p\bar{p})/\Gamma_{\text{total}} \quad \Gamma_{237}/\Gamma$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.77^{+0.15}_{-0.09} OUR AVERAGE				

0.90^{+0.04+0.27}_{-0.11-0.55} ¹ ABLIKIM 12D BES3 $J/\psi \rightarrow \gamma p\bar{p}$

1.14^{+0.43+0.42}_{-0.30-0.26} 231 ² ALEXANDER 10 CLEO $J/\psi \rightarrow \gamma p\bar{p}$

0.70±0.04^{+0.19}_{-0.08} BAI 03F BES2 $J/\psi \rightarrow \gamma p\bar{p}$

¹ From the fit including final state interaction effects in isospin 0 S-wave according to SIBIRTSEV 05A.

² From a fit of the $p\bar{p}$ mass distribution to a combination of $\gamma X(1835)$, γR with $M(R) = 2100$ MeV and $\Gamma(R) = 160$ MeV, and $\gamma p\bar{p}$ phase space, for $M(p\bar{p}) < 2.85$ GeV.

$$\Gamma(\gamma X(1835) \rightarrow \gamma K_S^0 K_S^0 \eta)/\Gamma_{\text{total}} \quad \Gamma_{238}/\Gamma$$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
3.31^{+0.33+1.96}_{-0.30-1.29}	ABLIKIM	15T	BES3 $J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta$

$\Gamma(\gamma X(1840) \rightarrow \gamma 3(\pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{239}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.44 \pm 0.36^{+0.60}_{-0.74}$	0.6k	ABLIKIM	13U BES3	$J/\psi \rightarrow \gamma 3(\pi^+ \pi^-)$

 $\Gamma(\gamma(K\bar{K}\pi) [J^{PC} = 0^{-+}])/ \Gamma_{\text{total}}$ Γ_{240}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.7 ± 0.4 OUR AVERAGE	Error includes scale factor of 2.1.		
$0.58 \pm 0.03 \pm 0.20$	¹ BAI	00D BES	$J/\psi \rightarrow \gamma K^{\pm} K_S^0 \pi^{\mp}$
$2.1 \pm 0.1 \pm 0.7$	² BAI	00D BES	$J/\psi \rightarrow \gamma K^{\pm} K_S^0 \pi^{\mp}$

¹ For a broad structure around 1800 MeV.² For a broad structure around 2040 MeV. $\Gamma(\gamma\pi^0)/\Gamma_{\text{total}}$ Γ_{241}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$3.49^{+0.33}_{-0.30}$ OUR AVERAGE				

 $3.63 \pm 0.36 \pm 0.13$ PEDLAR 09 CLE3 $J/\psi \rightarrow \pi^0 \gamma$ $3.13^{+0.65}_{-0.47}$ 586 ABLIKIM 06E BES2 $J/\psi \rightarrow \pi^0 \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $3.6 \pm 1.1 \pm 0.7$ BLOOM 83 CBAL $e^+ e^-$ 7.3 ± 4.7 10 BRANDELIK 79C DASP $e^+ e^-$ $\Gamma(\gamma\rho\bar{\rho}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{242}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.79	90	EATON	84 MRK2	$e^+ e^-$

 $\Gamma(\gamma\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$ Γ_{243}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.13	90	HENRARD	87 DM2	$e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.16 90 BAI 98G BES $e^+ e^-$ $\Gamma(\gamma f_0(2100) \rightarrow \gamma\eta\eta)/\Gamma_{\text{total}}$ Γ_{244}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.13^{+0.09+0.64}_{-0.10-0.28}$	5.5k	¹ ABLIKIM	13N BES3	$J/\psi \rightarrow \gamma\eta\eta$

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances. $\Gamma(\gamma f_0(2100) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$ Γ_{245}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	COMMENT
$6.24 \pm 0.48 \pm 0.87$	744	¹ DOBBS	15 $J/\psi \rightarrow \gamma\pi\pi$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\gamma f_0(2200))/\Gamma_{\text{total}}$ Γ_{246}/Γ

VALUE (units 10^{-4})		DOCUMENT ID	TECN	COMMENT
--------------------------	--	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.5		¹ AUGUSTIN	88	DM2 $J/\psi \rightarrow \gamma K_S^0 K_S^0$
-----	--	-----------------------	----	---

¹ Includes unknown branching fraction to $K_S^0 K_S^0$.

$\Gamma(\gamma f_0(2200) \rightarrow \gamma K \bar{K})/\Gamma_{\text{total}}$ Γ_{247}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

5.86 ± 0.49 ± 1.20	490	¹ DOBBS	15	$J/\psi \rightarrow \gamma K \bar{K}$
---------------------------	-----	--------------------	----	---------------------------------------

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\gamma f_J(2220))/\Gamma_{\text{total}}$ Γ_{248}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

>300		¹ BAI	96B	BES	$e^+ e^- \rightarrow \gamma \bar{p} p, K \bar{K}$
>250	99.9	² HASAN	96	SPEC	$\bar{p} p \rightarrow \pi^+ \pi^-$
< 2.3	95	³ AUGUSTIN	88	DM2	$J/\psi \rightarrow \gamma K^+ K^-$
< 1.6	95	³ AUGUSTIN	88	DM2	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
$12.4^{+6.4}_{-5.2} \pm 2.8$		³ BALTRUSAIT..86D	MRK3		$J/\psi \rightarrow \gamma K_S^0 K_S^0$
$8.4^{+3.4}_{-2.8} \pm 1.6$		³ BALTRUSAIT..86D	MRK3		$J/\psi \rightarrow \gamma K^+ K^-$

¹ Using BARNES 93.

² Using BAI 96B.

³ Includes unknown branching fraction to $K^+ K^-$ or $K_S^0 K_S^0$.

$\Gamma(\gamma f_J(2220) \rightarrow \gamma \pi \pi)/\Gamma_{\text{total}}$ Γ_{249}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

< 3.9	90	^{1,2} DOBBS	15	$J/\psi \rightarrow \gamma \pi \pi$
-----------------	----	----------------------	----	-------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$14 \pm 8 \pm 4$		BAI	98H	BES	$J/\psi \rightarrow \gamma \pi^0 \pi^0$
$8.4 \pm 2.6 \pm 3.0$		BAI	96B	BES	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^+ \pi^-$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² For $\Gamma = 20/50$ MeV, the 90% CL upper limits for $\pi^+ \pi^-$ and $\pi^0 \pi^0$ are $2.6/5.2 \times 10^{-5}$ and $1.3/1.9 \times 10^{-5}$, respectively.

$\Gamma(\gamma f_J(2220) \rightarrow \gamma K \bar{K})/\Gamma_{\text{total}}$ Γ_{250}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

< 4.1	90	^{1,2} DOBBS	15	$J/\psi \rightarrow \gamma K \bar{K}$
-----------------	----	----------------------	----	---------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 3.6		³ DEL-AMO-SA..100	BABR	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma K^+ K^-$	
< 2.9		³ DEL-AMO-SA..100	BABR	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma K_S^0 K_S^0$	
$6.6 \pm 2.9 \pm 2.4$		BAI	96B	BES	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma K^+ K^-$
$10.8 \pm 4.0 \pm 3.2$		BAI	96B	BES	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma K_S^0 K_S^0$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² For $\Gamma = 20/50$ MeV, the 90% CL upper limits for $K^+ K^-$ and $K_S^0 K_S^0$ are $1.7/3.1 \times 10^{-5}$ and $1.2/2.0 \times 10^{-5}$, respectively.

³ For spin 2 and helicity 0; other combinations lead to more stringent upper limits.

$\Gamma(\gamma f_J(2220) \rightarrow \gamma p \bar{p})/\Gamma_{\text{total}}$ Γ_{251}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$1.5 \pm 0.6 \pm 0.5$	BAI	96B	BES $e^+ e^- \rightarrow J/\psi \rightarrow \gamma p \bar{p}$

 $\Gamma(\gamma f_2(2340) \rightarrow \gamma \eta \eta)/\Gamma_{\text{total}}$ Γ_{252}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$5.60^{+0.62+2.37}_{-0.65-2.07}$	5.5k	¹ ABLIKIM	13N	BES3 $J/\psi \rightarrow \gamma \eta \eta$

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

 $\Gamma(\gamma f_0(1500) \rightarrow \gamma \pi \pi)/\Gamma_{\text{total}}$ Γ_{253}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.09 ± 0.24 OUR AVERAGE				
$1.21 \pm 0.29 \pm 0.24$	174	¹ DOBBS	15	$J/\psi \rightarrow \gamma \pi \pi$
$1.00 \pm 0.03 \pm 0.45$		² ABLIKIM	06V	BES2 $e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^+ \pi^-$
$1.02 \pm 0.09 \pm 0.45$		² ABLIKIM	06V	BES2 $e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^0 \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.7 ± 0.8		^{3,4} BUGG	95	MRK3 $J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$
---------------	--	---------------------	----	--

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² Including unknown branching fraction to $\pi \pi$.

³ Including unknown branching ratio for $f_0(1500) \rightarrow \pi^+ \pi^- \pi^+ \pi^-$.

⁴ Assuming that $f_0(1500)$ decays only to two S -wave dipions.

 $\Gamma(\gamma f_0(1500) \rightarrow \gamma \eta \eta)/\Gamma_{\text{total}}$ Γ_{254}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.65^{+0.26+0.51}_{-0.31-1.40}$	5.5k	¹ ABLIKIM	13N	BES3 $J/\psi \rightarrow \gamma \eta \eta$

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

 $\Gamma(\gamma A \rightarrow \gamma \text{invisible})/\Gamma_{\text{total}}$ Γ_{255}/Γ
(narrow state A with $m_A < 960$ MeV)

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 6.3	90	¹ INSLER	10	CLEO $e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$

¹ The limit varies with mass m_A of a narrow state A and is 4.3×10^{-6} for $m_A = 0$ MeV, reaches its largest value of 6.3×10^{-6} at $m_A = 500$ MeV, and is 3.6×10^{-6} at $m_A = 960$ MeV.

 $\Gamma(\gamma A^0 \rightarrow \gamma \mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{256}/Γ
(narrow state A^0 with $0.2 \text{ GeV} < m_{A^0} < 3 \text{ GeV}$)

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
< 0.5	90	¹ ABLIKIM	16E	BES3 $J/\psi \rightarrow \gamma \mu^+ \mu^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 2.1	90	² ABLIKIM	12	BES3 $J/\psi \rightarrow \gamma \mu^+ \mu^-$
---------	----	----------------------	----	--

¹ For a narrow scalar or pseudoscalar, A^0 , with a mass in the range 0.212–3 GeV. The measured 90% CL limit as a function of m_{A^0} is in the range $(2.8\text{--}495.3) \times 10^{-8}$.

² For a narrow scalar or pseudoscalar, A^0 , with a mass in the range 0.21–3.00 GeV. The measured 90% CL limit as a function of m_{A^0} ranges from 4×10^{-7} to 2.1×10^{-5} .

DALITZ DECAYS

 $\Gamma(\pi^0 e^+ e^-)/\Gamma_{\text{total}}$ Γ_{257}/Γ

<u>VALUE (units 10^{-7})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$7.56 \pm 1.32 \pm 0.50$	39	ABLIKIM 14I	BES3	$J/\psi \rightarrow \pi^0 e^+ e^-$

 $\Gamma(\eta e^+ e^-)/\Gamma_{\text{total}}$ Γ_{258}/Γ

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.16 \pm 0.07 \pm 0.06$	320	¹ ABLIKIM 14I	BES3	$J/\psi \rightarrow \eta e^+ e^-$

¹ Using both $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow \pi^+ \pi^- \pi^0$ decays.

 $\Gamma(\eta'(958) e^+ e^-)/\Gamma_{\text{total}}$ Γ_{259}/Γ

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$5.81 \pm 0.16 \pm 0.31$	1.4k	¹ ABLIKIM 14I	BES3	$J/\psi \rightarrow \eta' e^+ e^-$

¹ Using both $\eta' \rightarrow \gamma\pi^+ \pi^-$ and $\eta' \rightarrow \pi^+ \pi^- \eta$ decays.

WEAK DECAYS

 $\Gamma(D^- e^+ \nu_e + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{260}/Γ

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 1.2	90	ABLIKIM 06M	BES2	$e^+ e^- \rightarrow J/\psi$

 $\Gamma(\bar{D}^0 e^+ e^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{261}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 8.5 \times 10^{-8}$	90	¹ ABLIKIM 17AF	BES3	$e^+ e^- \rightarrow J/\psi$

- • • We do not use the following data for averages, fits, limits, etc. • • •

$< 1.1 \times 10^{-5}$	90	ABLIKIM 06M	BES2	$e^+ e^- \rightarrow J/\psi$
------------------------	----	-------------	------	------------------------------

¹ Using D^0 decays to $K^- \pi^+$, $K^- \pi^+ \pi^0$, and $K^- \pi^+ \pi^+ \pi^-$.

 $\Gamma(D_s^- e^+ \nu_e + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{262}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 1.3	90	ABLIKIM 14R	BES3	$e^+ e^- \rightarrow J/\psi$

- • • We do not use the following data for averages, fits, limits, etc. • • •

< 36	90	¹ ABLIKIM 06M	BES2	$e^+ e^- \rightarrow J/\psi$
--------	----	--------------------------	------	------------------------------

¹ Using $B(D_s^- \rightarrow \phi \pi^-) = 4.4 \pm 0.5\%$.

 $\Gamma(D_s^{*-} e^+ \nu_e + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{263}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 1.8 \times 10^{-6}$	90	ABLIKIM 14R	BES3	$e^+ e^- \rightarrow J/\psi$

 $\Gamma(D^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{264}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 7.5 \times 10^{-5}$	90	ABLIKIM 08J	BES2	$e^+ e^- \rightarrow J/\psi$

 $\Gamma(\bar{D}^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{265}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$< 1.7 \times 10^{-4}$	90	ABLIKIM 08J	BES2	$e^+ e^- \rightarrow J/\psi$

$\Gamma(\overline{D}^0 \overline{K}^{*0} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{266}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 2.5 \times 10^{-6}$	90	ABLIKIM 14K	BES3	$e^+ e^- \rightarrow J/\psi$

$\Gamma(D_s^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{267}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.3 \times 10^{-4}$	90	ABLIKIM 08J	BES2	$e^+ e^- \rightarrow J/\psi$

$\Gamma(D_s^- \rho^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{268}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.3 \times 10^{-5}$	90	ABLIKIM 14K	BES3	$e^+ e^- \rightarrow J/\psi$

———— CHARGE CONJUGATION (C), PARITY (P), ————

———— LEPTON FAMILY NUMBER (LF) VIOLATING MODES ————

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{269}/Γ

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
< 2.7	90	ABLIKIM 14Q	BES3	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
< 50	90	ADAMS 08	CLEO	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$
< 1600	90	¹ WICHT 08	BELL	$B^\pm \rightarrow K^\pm \gamma\gamma$
< 220	90	ABLIKIM 07J	BES2	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
< 5000	90	BARTEL 77	CNTR	$e^+ e^-$

¹ WICHT 08 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow J/\psi(1S) K^+)] < 0.16 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow J/\psi(1S) K^+) = 1.010 \times 10^{-3}$.

$\Gamma(\gamma\phi)/\Gamma_{\text{total}}$ Γ_{270}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.4 \times 10^{-6}$	90	ABLIKIM 14Q	BES3	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$

$\Gamma(e^\pm \mu^\mp)/\Gamma_{\text{total}}$ Γ_{271}/Γ

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
< 1.6	90	ABLIKIM 13L	BES3	$e^+ e^- \rightarrow J/\psi$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
< 11	90	BAI 03D	BES	$e^+ e^- \rightarrow J/\psi$

$\Gamma(e^\pm \tau^\mp)/\Gamma_{\text{total}}$ Γ_{272}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 8.3	90	ABLIKIM 04	BES	$e^+ e^- \rightarrow J/\psi$

$\Gamma(\mu^\pm \tau^\mp)/\Gamma_{\text{total}}$ Γ_{273}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 2.0	90	ABLIKIM 04	BES	$e^+ e^- \rightarrow J/\psi$

OTHER DECAYS

$\Gamma(\text{invisible})/\Gamma(e^+e^-)$					Γ_{274}/Γ_5
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<6.6 \times 10^{-2}$	90	LEES	13I	BABR	$B \rightarrow K^{(*)} J/\psi$

$\Gamma(\text{invisible})/\Gamma(\mu^+\mu^-)$					Γ_{274}/Γ_7
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.2 \times 10^{-2}$	90	ABLIKIM	08G	BES2	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$

 $J/\psi(1S)$ REFERENCES

ABLIKIM	17AF	PR D96 111101	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	17AH	PR D96 112001	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	17AK	PR D96 112012	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	17E	PL B770 217	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	17L	PR D95 052003	M. Ablikim <i>et al.</i>	(BES III Collab.)
LEES	17A	PR D95 052001	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	17C	PR D95 072007	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	17D	PR D95 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	16E	PR D93 052005	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	16J	PRL 117 042002	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	16K	PR D93 052010	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	16L	PR D93 072003	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	16M	PR D93 072008	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	16N	PR D93 112011	M. Ablikim	(BES III Collab.)
ABLIKIM	16P	PR D94 072005	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	16Q	PL B761 98	M. Ablikim <i>et al.</i>	(BES III Collab.)
AAIJ	15BI	EPJ C75 311	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	15AE	PR D92 052003	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	15H	PR D91 052017	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	15K	PR D91 112001	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	15P	PR D92 012007	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	15T	PRL 115 091803	M. Ablikim <i>et al.</i>	(BES III Collab.)
ANASHIN	15	PL B749 50	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)
LEES	15J	PR D92 072008	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	14I	PR D89 092008	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	14K	PR D89 071101	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	14N	PR D90 052009	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	14Q	PR D90 092002	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	14R	PR D90 112014	M. Ablikim <i>et al.</i>	(BES III Collab.)
ANASHIN	14	PL B738 391	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
AULCHENKO	14	PL B731 227	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)
LEES	14H	PR D89 092002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	13F	PR D87 052007	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	13I	PR D87 032003	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	13J	PR D87 032008	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	13L	PR D87 112007	Ablikim M. <i>et al.</i>	(BES III Collab.)
ABLIKIM	13N	PR D87 092009	Ablikim M. <i>et al.</i>	(BES III Collab.)
ABLIKIM	13P	PR D87 112004	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	13R	PR D88 032007	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	13U	PR D88 091502	M. Ablikim <i>et al.</i>	(BES III Collab.)
LEES	13I	PR D87 112005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	13O	PR D87 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	13Q	PR D88 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	13Y	PR D88 072009	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	12	PR D85 092012	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	12B	PR D86 032008	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	12C	PR D86 032014	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	12D	PRL 108 112003	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	12H	PL B710 594	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	12P	CP C36 1031	M. Ablikim <i>et al.</i>	(BES II Collab.)
LEES	12E	PR D85 112009	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	12F	PR D86 012008	J.P. Lees <i>et al.</i>	(BABAR Collab.)
METREVELI	12	PR D85 092007	Z. Metreveli <i>et al.</i>	(NWES, FLOR, WAYN+)

ABLIKIM	11	PR D83 012003	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	11C	PRL 106 072002	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	11D	PR D83 032003	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	10C	PL B685 27	M. Ablikim <i>et al.</i>	(BES II Collab.)
ABLIKIM	10E	PL B693 88	M. Ablikim <i>et al.</i>	(BES II Collab.)
ALEXANDER	10	PR D82 092002	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
ANASHIN	10	PL B685 134	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
DEL-AMO-SA...	100	PRL 105 172001	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
INSLER	10	PR D81 091101	J. Insler <i>et al.</i>	(CLEO Collab.)
ABLIKIM	09	PL B676 25	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	09B	PR D80 052004	M. Ablikim <i>et al.</i>	(BES II Collab.)
MITCHELL	09	PRL 102 011801	R.E. Mitchell <i>et al.</i>	(CLEO Collab.)
PEDLAR	09	PR D79 111101	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)
SHEN	09	PR D80 031101	C.P. Shen <i>et al.</i>	(BELLE Collab.)
ABLIKIM	08	EPJ C53 15	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08A	PR D77 012001	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08C	PL B659 789	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08E	PR D77 032005	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08F	PRL 100 102003	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08G	PRL 100 192001	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08I	PL B662 330	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08J	PL B663 297	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08O	PR D78 092005	M. Ablikim <i>et al.</i>	(BES Collab.)
ADAMS	08	PRL 101 101801	G.S. Adams <i>et al.</i>	(CLEO Collab.)
AUBERT	08S	PR D77 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)
BESSION	08	PR D78 032012	D. Besson <i>et al.</i>	(CLEO Collab.)
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)
WICHT	08	PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)
ABLIKIM	07H	PR D76 092003	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	07J	PR D76 117101	M. Ablikim <i>et al.</i>	(BES Collab.)
ANDREOTTI	07	PL B654 74	M. Andreotti <i>et al.</i>	(Femilab E835 Collab.)
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)
Also		PR D77 119902E (errat.)	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07BD	PR D76 092006	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABLIKIM	06	PL B632 181	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06C	PL B633 681	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06E	PR D73 052008	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06F	PR D73 052007	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06H	PR D73 112007	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06J	PRL 96 162002	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06K	PRL 97 062001	M. Ablikim <i>et al.</i>	(BES II Collab.)
ABLIKIM	06M	PL B639 418	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06V	PL B642 441	M. Ablikim <i>et al.</i>	(BES Collab.)
ADAMS	06A	PR D73 051103	G.S. Adams <i>et al.</i>	(CLEO Collab.)
AUBERT	06B	PR D73 012005	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	06D	PR D73 052003	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	06E	PRL 96 052002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT, BE	06D	PR D74 091103	B. Aubert <i>et al.</i>	(BABAR Collab.)
WU	06	PRL 97 162003	C.-H. Wu <i>et al.</i>	(BELLE Collab.)
ABLIKIM	05	PL B607 243	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05B	PR D71 032003	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05C	PL B610 192	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05H	PR D72 012002	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05R	PRL 95 262001	M. Ablikim <i>et al.</i>	(BES Collab.)
AUBERT	05D	PR D71 052001	B. Aubert <i>et al.</i>	(BABAR Collab.)
LI	05C	PR D71 111103	Z. Li <i>et al.</i>	(CLEO Collab.)
SIBIRTSEV	05A	PR D71 054010	A. Sibirtsev, J. Haidenbauer	
ABLIKIM	04	PL B598 172	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	04M	PR D70 112008	M. Ablikim <i>et al.</i>	(BES Collab.)
AUBERT	04	PR D69 011103	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT, B	04N	PR D70 072004	B. Aubert <i>et al.</i>	(BABAR Collab.)
BAI	04	PL B578 16	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04A	PR D69 012003	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04D	PL B589 7	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04E	PL B591 42	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04G	PR D70 012004	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04H	PR D70 012005	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04J	PL B594 47	J.Z. Bai <i>et al.</i>	(BES Collab.)
SETH	04	PR D69 097503	K.K. Seth	
AULCHENKO	03	PL B573 63	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)

BAI	03D	PL B561 49	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	03F	PRL 91 022001	J.Z. Bai <i>et al.</i>	(BES II Collab.)
BAI	03G	PR D68 052003	J.Z. Bai <i>et al.</i>	(BES Collab.)
HUANG	03	PRL 91 241802	H.-C. Huang <i>et al.</i>	(BELLE Collab.)
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>	
BAI	00	PRL 84 594	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	00B	PL B472 200	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	00D	PL B476 25	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	99	PL B446 356	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	99C	PRL 83 1918	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98D	PR D58 092006	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98G	PL B424 213	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98H	PRL 81 1179	J.Z. Bai <i>et al.</i>	(BES Collab.)
BALDINI	98	PL B444 111	R. Baldini <i>et al.</i>	(FENICE Collab.)
ARMSTRONG	96	PR D54 7067	T.A. Armstrong <i>et al.</i>	(E760 Collab.)
BAI	96B	PRL 76 3502	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	96C	PRL 77 3959	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	96D	PR D54 1221	J.Z. Bai <i>et al.</i>	(BES Collab.)
GRIBUSHIN	96	PR D53 4723	A. Gribushin <i>et al.</i>	(E672 Collab., E706 Collab.)
HASAN	96	PL B388 376	A. Hasan, D.V. Bugg	(BRUN, LOQM)
BAI	95B	PL B355 374	J.Z. Bai <i>et al.</i>	(BES Collab.)
BUGG	95	PL B353 378	D.V. Bugg <i>et al.</i>	(LOQM, PNPI, WASH)
ANTONELLI	93	PL B301 317	A. Antonelli <i>et al.</i>	(FENICE Collab.)
ARMSTRONG	93B	PR D47 772	T.A. Armstrong <i>et al.</i>	(FNAL E760 Collab.)
BARNES	93	PL B309 469	P.D. Barnes <i>et al.</i>	(PS185 Collab.)
AUGUSTIN	92	PR D46 1951	J.E. Augustin, G. Cosme	(DM2 Collab.)
BOLTON	92	PL B278 495	T. Bolton <i>et al.</i>	(Mark III Collab.)
BOLTON	92B	PRL 69 1328	T. Bolton <i>et al.</i>	(Mark III Collab.)
COFFMAN	92	PRL 68 282	D.M. Coffman <i>et al.</i>	(Mark III Collab.)
HSUEH	92	PR D45 2181	S. Hsueh, S. Palestini	(FNAL, TORI)
BISELLO	91	NP B350 1	D. Bisello <i>et al.</i>	(DM2 Collab.)
AUGUSTIN	90	PR D42 10	J.E. Augustin <i>et al.</i>	(DM2 Collab.)
BAI	90B	PRL 65 1309	Z. Bai <i>et al.</i>	(Mark III Collab.)
BAI	90C	PRL 65 2507	Z. Bai <i>et al.</i>	(Mark III Collab.)
BISELLO	90	PL B241 617	D. Bisello <i>et al.</i>	(DM2 Collab.)
COFFMAN	90	PR D41 1410	D.M. Coffman <i>et al.</i>	(Mark III Collab.)
JOUSSET	90	PR D41 1389	J. Jousset <i>et al.</i>	(DM2 Collab.)
ALEXANDER	89	NP B320 45	J.P. Alexander <i>et al.</i>	(LBL, MICH, SLAC)
AUGUSTIN	89	NP B320 1	J.E. Augustin, G. Cosme	(DM2 Collab.)
BISELLO	89B	PR D39 701	G. Busetto <i>et al.</i>	(DM2 Collab.)
AUGUSTIN	88	PRL 60 2238	J.E. Augustin <i>et al.</i>	(DM2 Collab.)
COFFMAN	88	PR D38 2695	D.M. Coffman <i>et al.</i>	(Mark III Collab.)
FALVARD	88	PR D38 2706	A. Falvard <i>et al.</i>	(CLER, FRAS, LALO+)
AUGUSTIN	87	ZPHY C36 369	J.E. Augustin <i>et al.</i>	(LALO, CLER, FRAS+)
BAGLIN	87	NP B286 592	C. Baglin <i>et al.</i>	(LAPP, CERN, GENO, LYON+)
BALTRUSAITIS...	87	PR D35 2077	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)
BECKER	87	PRL 59 186	J.J. Becker <i>et al.</i>	(Mark III Collab.)
BISELLO	87	PL B192 239	D. Bisello <i>et al.</i>	(PADO, CLER, FRAS+)
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)
HENRARD	87	NP B292 670	P. Henrard <i>et al.</i>	(CLER, FRAS, LALO+)
PALLIN	87	NP B292 653	D. Pallin <i>et al.</i>	(CLER, FRAS, LALO, PADO)
BALTRUSAITIS...	86	PR D33 629	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)
BALTRUSAITIS...	86B	PR D33 1222	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)
BALTRUSAITIS...	86D	PRL 56 107	R.M. Baltrusaitis	(CIT, UCSC, ILL, SLAC+)
BISELLO	86B	PL B179 294	D. Bisello <i>et al.</i>	(DM2 Collab.)
GAISER	86	PR D34 711	J. Gaiser <i>et al.</i>	(Crystal Ball Collab.)
BALTRUSAITIS...	85C	PRL 55 1723	R.M. Baltrusaitis <i>et al.</i>	(CIT, UCSC+)
BALTRUSAITIS...	85D	PR D32 566	R.M. Baltrusaitis <i>et al.</i>	(CIT, UCSC+)
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)
BALTRUSAITIS...	84	PRL 52 2126	Translated from YAF 41 733.	
EATON	84	PR D29 804	R.M. Baltrusaitis <i>et al.</i>	(CIT, UCSC+)
BLOOM	83	ARNS 33 143	M.W. Eaton <i>et al.</i>	(LBL, SLAC)
EDWARDS	83B	PRL 51 859	E.D. Bloom, C. Peck	(SLAC, CIT)
FRANKLIN	83	PRL 51 963	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
BURKE	82	PRL 49 632	M.E.B. Franklin <i>et al.</i>	(LBL, SLAC)
EDWARDS	82B	PR D25 3065	D.L. Burke <i>et al.</i>	(LBL, SLAC)
EDWARDS	82D	PRL 48 458	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
Also		ARNS 33 143	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
EDWARDS	82E	PRL 49 259	E.D. Bloom, C. Peck	(SLAC, CIT)
			C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)

LEMOIGNE	82	PL 113B 509	Y. Lemoigne <i>et al.</i>	(SACL, LOIC, SHMP+)
BESCH	81	ZPHY C8 1	H.J. Besch <i>et al.</i>	(BONN, DESY, MANZ)
GIDAL	81	PL 107B 153	G. Gidal <i>et al.</i>	(SLAC, LBL)
PARTRIDGE	80	PRL 44 712	R. Partridge <i>et al.</i>	(CIT, HARV, PRIN+)
SCHARRE	80	PL 97B 329	D.L. Scharre <i>et al.</i>	(SLAC, LBL)
ZHOLENTZ	80	PL 96B 214	A.A. Zholents <i>et al.</i>	(NOVO)
Also		SJNP 34 814	A.A. Zholents <i>et al.</i>	(NOVO)
		Translated from YAF 34	1471.	
BRANDELIK	79C	ZPHY C1 233	R. Brandelik <i>et al.</i>	(DASP Collab.)
ALEXANDER	78	PL 72B 493	G. Alexander <i>et al.</i>	(DESY, HAMB, SIEG+)
BESCH	78	PL 78B 347	H.J. Besch <i>et al.</i>	(BONN, DESY, MANZ)
BRANDELIK	78B	PL 74B 292	R. Brandelik <i>et al.</i>	(DASP Collab.)
PERUZZI	78	PR D17 2901	I. Peruzzi <i>et al.</i>	(SLAC, LBL)
BARTEL	77	PL 66B 489	W. Bartel <i>et al.</i>	(DESY, HEIDP)
BURMESTER	77D	PL 72B 135	J. Burmester <i>et al.</i>	(DESY, HAMB, SIEG+)
FELDMAN	77	PRPL 33C 285	G.J. Feldman, M.L. Perl	(LBL, SLAC)
VANNUCCI	77	PR D15 1814	F. Vannucci <i>et al.</i>	(SLAC, LBL)
BARTEL	76	PL 64B 483	W. Bartel <i>et al.</i>	(DESY, HEIDP)
BRAUNSCH...	76	PL 63B 487	W. Braunschweig <i>et al.</i>	(DASP Collab.)
JEAN-MARIE	76	PRL 36 291	B. Jean-Marie <i>et al.</i>	(SLAC, LBL) IG
BALDINI-...	75	PL 58B 471	R. Baldini-Celio <i>et al.</i>	(FRAS, ROMA)
BOYARSKI	75	PRL 34 1357	A.M. Boyarski <i>et al.</i>	(SLAC, LBL) JPC
DASP	75	PL 56B 491	W. Braunschweig <i>et al.</i>	(DASP Collab.)
ESPOSITO	75B	LNC 14 73	B. Esposito <i>et al.</i>	(FRAS, NAPL, PADO+)
FORD	75	PRL 34 604	R.L. Ford <i>et al.</i>	(SLAC, PENN)
