

$K^*(1680)$

$$I(J^P) = \frac{1}{2}(1^-)$$

 $K^*(1680)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1718±18 OUR AVERAGE					
1722±20 ⁺³³ ₋₁₀₉	4289	¹ AAIJ	17C	LHCB	$B^+ \rightarrow J/\psi \phi K^+$
1677±10±32		ASTON	88	LASS 0	11 $K^- p \rightarrow$ $K^- \pi^+ n$
1735±10±20		ASTON	87	LASS 0	11 $K^- p \rightarrow$ $\bar{K}^0 \pi^+ \pi^- n$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
1678±64		BIRD	89	LASS -	11 $K^- p \rightarrow \bar{K}^0 \pi^- p$
1800±70		ETKIN	80	MPS 0	6 $K^- p \rightarrow$ $\bar{K}^0 \pi^+ \pi^- n$
~ 1650		ESTABROOKS 78	ASPK 0	13	$K^\pm p \rightarrow$ $K^\pm \pi^\pm n$

¹ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 8.5 σ . **$K^*(1680)$ WIDTH**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
322±110 OUR AVERAGE					
Error includes scale factor of 4.2.					
354±75 ⁺¹⁴⁰ ₋₁₈₁	4289	² AAIJ	17C	LHCB	$B^+ \rightarrow J/\psi \phi K^+$
205±16±34		ASTON	88	LASS 0	11 $K^- p \rightarrow$ $K^- \pi^+ n$
423±18±30		ASTON	87	LASS 0	11 $K^- p \rightarrow$ $\bar{K}^0 \pi^+ \pi^- n$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
454±270		BIRD	89	LASS -	11 $K^- p \rightarrow \bar{K}^0 \pi^- p$
170±30		ETKIN	80	MPS 0	6 $K^- p \rightarrow$ $\bar{K}^0 \pi^+ \pi^- n$
250 to 300		ESTABROOKS 78	ASPK 0	13	$K^\pm p \rightarrow$ $K^\pm \pi^\pm n$

² From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 8.5 σ . **$K^*(1680)$ DECAY MODES**

Mode	Fraction (Γ_i/Γ)
Γ_1 $K\pi$	(38.7±2.5) %
Γ_2 $K\rho$	(31.4 ^{+5.0} _{-2.1}) %
Γ_3 $K^*(892)\pi$	(29.9 ^{+2.2} _{-5.0}) %
Γ_4 $K\phi$	seen

CONSTRAINED FIT INFORMATION

An overall fit to 4 branching ratios uses 4 measurements and one constraint to determine 3 parameters. The overall fit has a $\chi^2 = 2.9$ for 2 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	-36	
x_3	-39	-72
	x_1	x_2

 $K^*(1680)$ BRANCHING RATIOS

$\Gamma(K\pi)/\Gamma_{\text{total}}$ **Γ_1/Γ**

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
0.387 ± 0.026 OUR FIT				
$0.388 \pm 0.014 \pm 0.022$	ASTON	88	LASS	0 11 $K^- p \rightarrow K^- \pi^+ n$

$\Gamma(K\pi)/\Gamma(K^*(892)\pi)$ **Γ_1/Γ_3**

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
$1.30^{+0.23}_{-0.14}$ OUR FIT				
2.8 ± 1.1	ASTON	84	LASS	0 11 $K^- p \rightarrow \bar{K}^0 2\pi n$

$\Gamma(K\rho)/\Gamma(K\pi)$ **Γ_2/Γ_1**

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
$0.81^{+0.14}_{-0.09}$ OUR FIT				
1.2 ± 0.4	ASTON	84	LASS	0 11 $K^- p \rightarrow \bar{K}^0 2\pi n$

$\Gamma(K\rho)/\Gamma(K^*(892)\pi)$ **Γ_2/Γ_3**

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
$1.05^{+0.27}_{-0.11}$ OUR FIT				
$0.97 \pm 0.09^{+0.30}_{-0.10}$	ASTON	87	LASS	0 11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$

$\Gamma(K\phi)/\Gamma_{\text{total}}$ **Γ_4/Γ**

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
seen	4289	³ AAIJ	17C	LHCB $B^+ \rightarrow J/\psi \phi K^+$

³From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 8.5σ .

$K^*(1680)$ REFERENCES

AAIJ	17C	PRL 118 022003	R. Aaij <i>et al.</i>	(LHCb Collab.)
Also		PR D95 012002	R. Aaij <i>et al.</i>	(LHCb Collab.)
BIRD	89	SLAC-332	P.F. Bird	(SLAC)
ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
ASTON	87	NP B292 693	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
ASTON	84	PL 149B 258	D. Aston <i>et al.</i>	(SLAC, CARL, OTTA) JP
ETKIN	80	PR D22 42	A. Etkin <i>et al.</i>	(BNL, CUNY) JP
ESTABROOKS	78	NP B133 490	P.G. Estabrooks <i>et al.</i>	(MCGI, CARL, DURH+) JP
