

$\eta_c(1S)$

$$J^{PC} = 0^+(0^-+)$$

$\eta_c(1S)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2983.9 ± 0.5 OUR AVERAGE		Error includes scale factor of 1.3. See the ideogram below.		
2986.7 ± 0.5 ± 0.9	11K	¹ AAIJ	17AD LHCb	$pp \rightarrow B^+ X \rightarrow p\bar{p}K^+ X$
2982.8 ± 1.0 ± 0.5	6.4k	² AAIJ	17BB LHCb	$pp \rightarrow b\bar{b}X \rightarrow 2(K^+K^-)X$
2982.2 ± 1.5 ± 0.1	2.0k	³ AAIJ	15BI LHCb	$pp \rightarrow \eta_c(1S)X$
2983.5 ± 1.4 $\begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 1.6 \\ 3.6 \end{smallmatrix}$		⁴ ANASHIN	14 KEDR	$J/\psi \rightarrow \gamma\eta_c$
2979.8 ± 0.8 ± 3.5	4.5k	^{5,6} LEES	14E BABR	$\gamma\gamma \rightarrow K^+K^-\pi^0$
2984.1 ± 1.1 ± 2.1	900	^{5,6,7} LEES	14E BABR	$\gamma\gamma \rightarrow K^+K^-\eta$
2984.3 ± 0.6 ± 0.6		^{8,9} ABLIKIM	12F BES3	$\psi(2S) \rightarrow \gamma\eta_c$
2984.49 ± 1.16 ± 0.52	832	⁵ ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0\gamma \text{ hadrons}$
2982.7 ± 1.8 ± 2.2	486	ZHANG	12A BELL	$e^+e^- \rightarrow e^+e^-\eta'/\pi^+\pi^-$
2984.5 ± 0.8 ± 3.1	11k	DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K^+K^-\pi^+\pi^-\pi^0$
2985.4 ± 1.5 $\begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 0.5 \\ 2.0 \end{smallmatrix}$	920	⁹ VINOKUROVA	11 BELL	$B^\pm \rightarrow K^\pm(K_S^0 K^\pm\pi^\mp)$
2982.2 ± 0.4 ± 1.6	14k	¹⁰ LEES	10 BABR	$10.6 e^+e^- \rightarrow e^+e^-K_S^0 K^\pm\pi^\mp$
2985.8 ± 1.5 ± 3.1	0.9k	AUBERT	08AB BABR	$B \rightarrow \eta_c(1S)K^{(*)} \rightarrow K\bar{K}\pi K^{(*)}$
2986.1 ± 1.0 ± 2.5	7.5k	UEHARA	08 BELL	$\gamma\gamma \rightarrow \eta_c \rightarrow \text{hadrons}$
2970 ± 5 ± 6	501	¹¹ ABE	07 BELL	$e^+e^- \rightarrow J/\psi(c\bar{c})$
2971 ± 3 $\begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 2 \\ 1 \end{smallmatrix}$	195	WU	06 BELL	$B^+ \rightarrow p\bar{p}K^+$
2974 ± 7 $\begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 2 \\ 1 \end{smallmatrix}$	20	WU	06 BELL	$B^+ \rightarrow \Lambda\bar{\Lambda}K^+$
2981.8 ± 1.3 ± 1.5	592	ASNER	04 CLEO	$\gamma\gamma \rightarrow \eta_c \rightarrow K_S^0 K^\pm\pi^\mp$
2984.1 ± 2.1 ± 1.0	190	¹² AMBROGIANI	03 E835	$\bar{p}p \rightarrow \eta_c \rightarrow \gamma\gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
2982.5 ± 0.4 ± 1.4	12k	¹³ DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K_S^0 K^\pm\pi^\mp$
2982.2 ± 0.6		¹⁴ MITCHELL	09 CLEO	$e^+e^- \rightarrow \gamma X$
2982 ± 5	270	¹⁵ AUBERT	06E BABR	$B^\pm \rightarrow K^\pm X_c\bar{c}$
2982.5 ± 1.1 ± 0.9	2.5k	¹⁶ AUBERT	04D BABR	$\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K\bar{K}\pi$
2977.5 ± 1.0 ± 1.2		^{14,17} BAI	03 BES	$J/\psi \rightarrow \gamma\eta_c$
2979.6 ± 2.3 ± 1.6	180	¹⁸ FANG	03 BELL	$B \rightarrow \eta_c K$
2976.3 ± 2.3 ± 1.2		^{14,19} BAI	00F BES	$J/\psi, \psi(2S) \rightarrow \gamma\eta_c$
2976.6 ± 2.9 ± 1.3	140	^{14,20} BAI	00F BES	$J/\psi \rightarrow \gamma\eta_c$
2980.4 ± 2.3 ± 0.6		²¹ BRANDENB...	00B CLE2	$\gamma\gamma \rightarrow \eta_c \rightarrow K^\pm K_S^0\pi^\mp$

2975.8 ± 3.9 ± 1.2		20	BAI	99B	BES	Sup. by BAI 00F
2999 ± 8	25		ABREU	980	DLPH	$e^+e^- \rightarrow e^+e^- + \text{hadrons}$
2988.3 + 3.3 - 3.1			ARMSTRONG	95F	E760	$\bar{p}p \rightarrow \gamma\gamma$
2974.4 ± 1.9	14,22		BISELLO	91	DM2	$J/\psi \rightarrow \eta_c \gamma$
2969 ± 4 ± 4	80	14	BAI	90B	MRK3	$J/\psi \rightarrow$ $\gamma K^+ K^- K^+ K^-$
2956 ± 12 ± 12		14	BAI	90B	MRK3	$J/\psi \rightarrow$ $\gamma K^+ K^- K_S^0 K_L^0$
2982.6 + 2.7 - 2.3	12		BAGLIN	87B	SPEC	$\bar{p}p \rightarrow \gamma\gamma$
2980.2 ± 1.6	14,22		BALTRUSAIT..	86	MRK3	$J/\psi \rightarrow \eta_c \gamma$
2984 ± 2.3 ± 4.0	14		GAISER	86	CBAL	$J/\psi \rightarrow \gamma X, \psi(2S) \rightarrow$ γX
2976 ± 8	14,23		BALTRUSAIT..	84	MRK3	$J/\psi \rightarrow 2\phi\gamma$
2982 ± 8	18	24	HIMEL	80B	MRK2	e^+e^-
2980 ± 9		24	PARTRIDGE	80B	CBAL	e^+e^-

¹ AAIJ 17AD report $m_{J/\psi} - m_{\eta_c(1S)} = 110.2 \pm 0.5 \pm 0.9$ MeV. We use the current value $m_{J/\psi} = 3096.900 \pm 0.006$ MeV to obtain the quoted mass.

² From a fit of the $\phi\phi$ invariant mass with the mass and width of $\eta_c(1S)$ as free parameters.

³ AAIJ 15BI reports $m_{J/\psi} - m_{\eta_c(1S)} = 114.7 \pm 1.5 \pm 0.1$ MeV from a sample of $\eta_c(1S)$ and J/ψ produced in b -hadron decays. We have used current value of $m_{J/\psi} = 3096.900 \pm 0.006$ MeV to arrive at the quoted $m_{\eta_c(1S)}$ result.

⁴ Taking into account an asymmetric photon lineshape.

⁵ With floating width.

⁶ Ignoring possible interference with the non-resonant 0^- amplitude.

⁷ Using both, $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow \pi^+\pi^-\pi^0$ decays.

⁸ From a simultaneous fit to six decay modes of the η_c .

⁹ Accounts for interference with non-resonant continuum.

¹⁰ Taking into account interference with the non-resonant $J^P = 0^-$ amplitude.

¹¹ From a fit of the J/ψ recoil mass spectrum. Supersedes ABE,K 02 and ABE 04G.

¹² Using mass of $\psi(2S) = 3686.00$ MeV.

¹³ Not independent from the measurements reported by LEES 10.

¹⁴ MITCHELL 09 observes a significant asymmetry in the lineshapes of $\psi(2S) \rightarrow \gamma\eta_c$ and $J/\psi \rightarrow \gamma\eta_c$ transitions. If ignored, this asymmetry could lead to significant bias whenever the mass and width are measured in $\psi(2S)$ or J/ψ radiative decays.

¹⁵ From the fit of the kaon momentum spectrum. Systematic errors not evaluated.

¹⁶ Superseded by LEES 10.

¹⁷ From a simultaneous fit of five decay modes of the η_c .

¹⁸ Superseded by VINOKUROVA 11.

¹⁹ Weighted average of the $\psi(2S)$ and $J/\psi(1S)$ samples. Using an η_c width of 13.2 MeV.

²⁰ Average of several decay modes. Using an η_c width of 13.2 MeV.


²¹ Superseded by ASNER 04.

²² Average of several decay modes.

²³ $\eta_c \rightarrow \phi\phi$.

²⁴ Mass adjusted by us to correspond to $J/\psi(1S)$ mass = 3097 MeV.

WEIGHTED AVERAGE
 2983.9 ± 0.5 (Error scaled by 1.3)



$\eta_c(1S)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
32.0 ± 0.8 OUR FIT				
32.1 ± 0.9 OUR AVERAGE				Error includes scale factor of 1.1.
$34.0 \pm 1.9 \pm 1.3$	11K	AAIJ	17AD LHCb	$pp \rightarrow B^+ X \rightarrow p\bar{p}K^+ X$
$31.4 \pm 3.5 \pm 2.0$	6.4k	1 AAIJ	17BB LHCb	$pp \rightarrow b\bar{b}X \rightarrow 2(K^+K^-)X$
$27.2 \pm 3.1^{+5.4}_{-2.6}$		2 ANASHIN	14 KEDR	$J/\psi \rightarrow \gamma\eta_c$
$25.2 \pm 2.6 \pm 2.4$	4.5k	3,4 LEES	14E BABR	$\gamma\gamma \rightarrow K^+K^-\pi^0$
$34.8 \pm 3.1 \pm 4.0$	900	3,4,5 LEES	14E BABR	$\gamma\gamma \rightarrow K^+K^-\eta$
$32.0 \pm 1.2 \pm 1.0$		6,7 ABLIKIM	12F BES3	$\psi(2S) \rightarrow \gamma\eta_c$
$36.4 \pm 3.2 \pm 1.7$	832	3 ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0\gamma$ hadrons
$37.8^{+5.8}_{-5.3} \pm 3.1$	486	ZHANG	12A BELL	$e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$
$36.2 \pm 2.8 \pm 3.0$	11k	DEL-AMO-SA..	11M BABR	$\gamma\gamma \rightarrow K^+K^-\pi^+\pi^-\pi^0$
$35.1 \pm 3.1^{+1.0}_{-1.6}$	920	7 VINOKUROVA	11 BELL	$B^\pm \rightarrow K^\pm(K_S^0 K^\pm\pi^\mp)$
$31.7 \pm 1.2 \pm 0.8$	14k	8 LEES	10 BABR	$10.6 e^+e^- \rightarrow e^+e^-K_S^0 K^\pm\pi^\mp$
$36.3^{+3.7}_{-3.6} \pm 4.4$	0.9k	AUBERT	08AB BABR	$B \rightarrow \eta_c(1S)K^{(*)} \rightarrow K\bar{K}\pi K^{(*)}$

$28.1 \pm 3.2 \pm 2.2$	7.5k	UEHARA	08	BELL	$\gamma\gamma \rightarrow \eta_c \rightarrow \text{hadrons}$
$48 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 8 \\ 7 \end{smallmatrix} \pm 5$	195	WU	06	BELL	$B^+ \rightarrow p \bar{p} K^+$
$40 \pm 19 \pm 5$	20	WU	06	BELL	$B^+ \rightarrow \Lambda \bar{\Lambda} K^+$
$24.8 \pm 3.4 \pm 3.5$	592	ASNER	04	CLEO	$\gamma\gamma \rightarrow \eta_c \rightarrow K_S^0 K^\pm \pi^\mp$
$20.4 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 7.7 \\ 6.7 \end{smallmatrix} \pm 2.0$	190	AMBROGIANI	03	E835	$\bar{p} p \rightarrow \eta_c \rightarrow \gamma\gamma$
$23.9 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 12.6 \\ 7.1 \end{smallmatrix}$		ARMSTRONG	95F	E760	$\bar{p} p \rightarrow \gamma\gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
$32.1 \pm 1.1 \pm 1.3$	12k	⁹ DEL-AMO-SA..	11M	BABR	$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$
$34.3 \pm 2.3 \pm 0.9$	2.5k	¹⁰ AUBERT	04D	BABR	$\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K \bar{K} \pi$
$17.0 \pm 3.7 \pm 7.4$		¹¹ BAI	03	BES	$J/\psi \rightarrow \gamma \eta_c$
$29 \pm 8 \pm 6$	180	¹² FANG	03	BELL	$B \rightarrow \eta_c K$
$11.0 \pm 8.1 \pm 4.1$		¹³ BAI	00F	BES	$J/\psi \rightarrow \gamma \eta_c$ and $\psi(2S) \rightarrow \gamma \eta_c$
$27.0 \pm 5.8 \pm 1.4$		¹⁴ BRANDENB...	00B	CLE2	$\gamma\gamma \rightarrow \eta_c \rightarrow K^\pm K_S^0 \pi^\mp$
$7.0 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 7.5 \\ 7.0 \end{smallmatrix}$	12	BAGLIN	87B	SPEC	$\bar{p} p \rightarrow \gamma\gamma$
$10.1 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 33.0 \\ 8.2 \end{smallmatrix}$	23	¹⁵ BALTRUSAIT..	86	MRK3	$J/\psi \rightarrow \gamma p \bar{p}$
11.5 ± 4.5		GAISER	86	CBAL	$J/\psi \rightarrow \gamma X, \psi(2S) \rightarrow \gamma X$
< 40 90% CL	18	HIMEL	80B	MRK2	$e^+ e^-$
< 20 90% CL		PARTRIDGE	80B	CBAL	$e^+ e^-$

¹ From a fit of the $\phi\phi$ invariant mass with the mass and width of $\eta_c(1S)$ as free parameters.

² Taking into account an asymmetric photon lineshape.

³ With floating mass.

⁴ Ignoring possible interference with the non-resonant 0^- amplitude.

⁵ Using both, $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow \pi^+ \pi^- \pi^0$ decays.

⁶ From a simultaneous fit to six decay modes of the η_c .

⁷ Accounts for interference with non-resonant continuum.

⁸ Taking into account interference with the non-resonant $J^P = 0^-$ amplitude.

⁹ Not independent from the measurements reported by LEES 10.

¹⁰ Superseded by LEES 10.

¹¹ From a simultaneous fit of five decay modes of the η_c .

¹² Superseded by VINOKUROVA 11.

¹³ From a fit to the 4-prong invariant mass in $\psi(2S) \rightarrow \gamma \eta_c$ and $J/\psi(1S) \rightarrow \gamma \eta_c$ decays.

¹⁴ Superseded by ASNER 04.

¹⁵ Positive and negative errors correspond to 90% confidence level.

$\eta_c(1S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
Decays involving hadronic resonances		
Γ_1 $\eta'(958) \pi \pi$	(4.1 \pm 1.7) %	
Γ_2 $\rho \rho$	(1.8 \pm 0.5) %	
Γ_3 $K^*(892)^0 K^- \pi^+ + \text{c.c.}$	(2.0 \pm 0.7) %	
Γ_4 $K^*(892) \bar{K}^*(892)$	(7.1 \pm 1.3) $\times 10^{-3}$	
Γ_5 $K^*(892)^0 \bar{K}^*(892)^0 \pi^+ \pi^-$	(1.1 \pm 0.5) %	

Γ_6	$\phi K^+ K^-$	$(2.9 \pm 1.4) \times 10^{-3}$	
Γ_7	$\phi\phi$	$(1.79 \pm 0.20) \times 10^{-3}$	
Γ_8	$\phi 2(\pi^+ \pi^-)$	$< 4 \times 10^{-3}$	90%
Γ_9	$a_0(980)\pi$	< 2 %	90%
Γ_{10}	$a_2(1320)\pi$	< 2 %	90%
Γ_{11}	$K^*(892)\bar{K} + \text{c.c.}$	< 1.28 %	90%
Γ_{12}	$f_2(1270)\eta$	< 1.1 %	90%
Γ_{13}	$\omega\omega$	$< 3.1 \times 10^{-3}$	90%
Γ_{14}	$\omega\phi$	$< 2.5 \times 10^{-4}$	90%
Γ_{15}	$f_2(1270) f_2(1270)$	$(9.8 \pm 2.5) \times 10^{-3}$	
Γ_{16}	$f_2(1270) f_2'(1525)$	$(9.8 \pm 3.2) \times 10^{-3}$	
Γ_{17}	$f_0(980)\eta$	seen	
Γ_{18}	$f_0(1500)\eta$	seen	
Γ_{19}	$f_0(2200)\eta$	seen	
Γ_{20}	$a_0(980)\pi$	seen	
Γ_{21}	$a_0(1320)\pi$	seen	
Γ_{22}	$a_0(1450)\pi$	seen	
Γ_{23}	$a_0(1950)\pi$	seen	
Γ_{24}	$a_2(1950)\pi$	not seen	
Γ_{25}	$K_0^*(1430)\bar{K}$	seen	
Γ_{26}	$K_2^*(1430)\bar{K}$	seen	
Γ_{27}	$K_0^*(1950)\bar{K}$	seen	

Decays into stable hadrons

Γ_{28}	$K\bar{K}\pi$	$(7.3 \pm 0.5) \%$	
Γ_{29}	$K\bar{K}\eta$	$(1.36 \pm 0.16) \%$	
Γ_{30}	$\eta\pi^+\pi^-$	$(1.7 \pm 0.5) \%$	
Γ_{31}	$\eta 2(\pi^+ \pi^-)$	$(4.4 \pm 1.3) \%$	
Γ_{32}	$K^+ K^- \pi^+ \pi^-$	$(6.9 \pm 1.1) \times 10^{-3}$	
Γ_{33}	$K^+ K^- \pi^+ \pi^- \pi^0$	$(3.5 \pm 0.6) \%$	
Γ_{34}	$K^0 K^- \pi^+ \pi^- \pi^+ + \text{c.c.}$	$(5.6 \pm 1.5) \%$	
Γ_{35}	$K^+ K^- 2(\pi^+ \pi^-)$	$(7.5 \pm 2.4) \times 10^{-3}$	
Γ_{36}	$2(K^+ K^-)$	$(1.47 \pm 0.31) \times 10^{-3}$	
Γ_{37}	$\pi^+ \pi^- \pi^0$	$< 5 \times 10^{-4}$	90%
Γ_{38}	$\pi^+ \pi^- \pi^0 \pi^0$	$(4.7 \pm 1.0) \%$	
Γ_{39}	$2(\pi^+ \pi^-)$	$(9.7 \pm 1.2) \times 10^{-3}$	
Γ_{40}	$2(\pi^+ \pi^- \pi^0)$	$(17.4 \pm 3.3) \%$	
Γ_{41}	$3(\pi^+ \pi^-)$	$(1.8 \pm 0.4) \%$	
Γ_{42}	$p\bar{p}$	$(1.52 \pm 0.16) \times 10^{-3}$	
Γ_{43}	$p\bar{p}\pi^0$	$(3.6 \pm 1.3) \times 10^{-3}$	
Γ_{44}	$\Lambda\bar{\Lambda}$	$(1.09 \pm 0.24) \times 10^{-3}$	
Γ_{45}	$\Sigma^+ \bar{\Sigma}^-$	$(2.1 \pm 0.6) \times 10^{-3}$	
Γ_{46}	$\Xi^- \bar{\Xi}^+$	$(9.0 \pm 2.6) \times 10^{-4}$	
Γ_{47}	$\pi^+ \pi^- p\bar{p}$	$(5.3 \pm 1.8) \times 10^{-3}$	

Radiative decays

$$\Gamma_{48} \quad \gamma\gamma \quad (1.57 \pm 0.12) \times 10^{-4}$$

Charge conjugation (C), Parity (P), Lepton family number (LF) violating modes

Γ_{49}	$\pi^+ \pi^-$	$P, CP < 1.1$	$\times 10^{-4}$	90%
Γ_{50}	$\pi^0 \pi^0$	$P, CP < 4$	$\times 10^{-5}$	90%
Γ_{51}	$K^+ K^-$	$P, CP < 6$	$\times 10^{-4}$	90%
Γ_{52}	$K_S^0 K_S^0$	$P, CP < 3.1$	$\times 10^{-4}$	90%

CONSTRAINED FIT INFORMATION

An overall fit to the total width, 8 combinations of partial widths obtained from integrated cross section, and 19 branching ratios uses 88 measurements and one constraint to determine 13 parameters. The overall fit has a $\chi^2 = 119.9$ for 76 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_7	18									
x_{15}	3	6								
x_{28}	21	41	7							
x_{29}	11	22	4	54						
x_{32}	11	20	4	24	13					
x_{36}	9	16	3	25	13	10				
x_{39}	13	25	5	30	16	15	12			
x_{42}	14	25	5	35	19	16	13	19		
x_{44}	3	6	1	9	5	4	3	5	25	
x_{48}	-29	-54	-10	-65	-35	-34	-26	-41	-45	-11
Γ	-1	-3	-1	-3	-2	-2	-1	-2	6	2
	x_4	x_7	x_{15}	x_{28}	x_{29}	x_{32}	x_{36}	x_{39}	x_{42}	x_{44}
Γ	-26									
	x_{48}									

	Mode	Rate (MeV)
Γ_4	$K^*(892) \bar{K}^*(892)$	0.23 ± 0.04
Γ_7	$\phi\phi$	0.057 ± 0.006
Γ_{15}	$f_2(1270) f_2(1270)$	0.31 ± 0.08
Γ_{28}	$K \bar{K} \pi$	2.34 ± 0.16
Γ_{29}	$K \bar{K} \eta$	0.43 ± 0.05

Γ_{32}	$K^+ K^- \pi^+ \pi^-$	0.222 ± 0.034
Γ_{36}	$2(K^+ K^-)$	0.047 ± 0.010
Γ_{39}	$2(\pi^+ \pi^-)$	0.31 ± 0.04
Γ_{42}	$p\bar{p}$	0.048 ± 0.005
Γ_{44}	$\Lambda\bar{\Lambda}$	0.035 ± 0.008
Γ_{48}	$\gamma\gamma$	0.0050 ± 0.0004

$\eta_c(1S)$ PARTIAL WIDTHS

$\Gamma(\gamma\gamma)$					Γ_{48}
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT	
5.0 ± 0.4 OUR FIT					
• • • We do not use the following data for averages, fits, limits, etc. • • •					
5.8 ± 1.1	486	¹ ZHANG	12A BELL	$e^+ e^- \rightarrow e^+ e^- \eta' \pi^+ \pi^-$	
5.2 ± 1.2	273 ± 43	^{2,3} AUBERT	06E BABR	$B^\pm \rightarrow K^\pm X_{c\bar{c}}$	
$5.5 \pm 1.2 \pm 1.8$	157 ± 33	⁴ KUO	05 BELL	$\gamma\gamma \rightarrow p\bar{p}$	
$7.4 \pm 0.4 \pm 2.3$		⁵ ASNER	04 CLEO	$\gamma\gamma \rightarrow \eta_c \rightarrow K_S^0 K^\pm \pi^\mp$	
$13.9 \pm 2.0 \pm 3.0$	41	⁶ ABDALLAH	03J DLPH	$\gamma\gamma \rightarrow \eta_c$	
$3.8^+_{-1.0} \pm 1.1^+_{-1.0}$	190	⁷ AMBROGIANI	03 E835	$\bar{p}p \rightarrow \eta_c \rightarrow \gamma\gamma$	
$7.6 \pm 0.8 \pm 2.3$		^{5,8} BRANDENB...	00B CLE2	$\gamma\gamma \rightarrow \eta_c \rightarrow K^\pm K_S^0 \pi^\mp$	
$6.9 \pm 1.7 \pm 2.1$	76	⁹ ACCIARRI	99T L3	$e^+ e^- \rightarrow e^+ e^- \eta_c$	
$27 \pm 16 \pm 10$	5	⁵ SHIRAI	98 AMY	$58 e^+ e^-$	
$6.7^+_{-1.7} \pm 2.4 \pm 2.3$		⁴ ARMSTRONG	95F E760	$\bar{p}p \rightarrow \gamma\gamma$	
11.3 ± 4.2		¹⁰ ALBRECHT	94H ARG	$e^+ e^- \rightarrow e^+ e^- \eta_c$	
$8.0 \pm 2.3 \pm 2.4$	17	¹¹ ADRIANI	93N L3	$e^+ e^- \rightarrow e^+ e^- \eta_c$	
$5.9^+_{-1.8} \pm 2.1 \pm 1.9$		⁷ CHEN	90B CLEO	$e^+ e^- \rightarrow e^+ e^- \eta_c$	
$6.4^+_{-3.4} \pm 5.0 \pm 3.4$		¹² AIHARA	88D TPC	$e^+ e^- \rightarrow e^+ e^- X$	
$4.3^+_{-3.7} \pm 3.4 \pm 2.4$		⁴ BAGLIN	87B SPEC	$\bar{p}p \rightarrow \gamma\gamma$	
28 ± 15		^{5,13} BERGER	86 PLUT	$\gamma\gamma \rightarrow K\bar{K}\pi$	

¹ Assuming there is no interference with the non-resonant background.

² Calculated by us using $\Gamma(\eta_c \rightarrow K\bar{K}\pi) \times \Gamma(\eta_c \rightarrow \gamma\gamma) / \Gamma = 0.44 \pm 0.05$ keV from PDG 06 and $B(\eta_c \rightarrow K\bar{K}\pi) = (8.5 \pm 1.8)\%$ from AUBERT 06E.

³ Systematic errors not evaluated.

⁴ Normalized to $B(\eta_c \rightarrow p\bar{p}) = (1.3 \pm 0.4) \times 10^{-3}$.

⁵ Normalized to $B(\eta_c \rightarrow K^\pm K_S^0 \pi^\mp)$.

⁶ Average of $K_S^0 K^\pm \pi^\mp$, $\pi^+ \pi^- K^+ K^-$, and $2(K^+ K^-)$ decay modes.

⁷ Normalized to the sum of $B(\eta_c \rightarrow K^\pm K_S^0 \pi^\mp)$, $B(\eta_c \rightarrow K^+ K^- \pi^+ \pi^-)$, and $B(\eta_c \rightarrow 2\pi^+ 2\pi^-)$.

⁸ Superseded by ASNER 04.

⁹ Normalized to the sum of 9 branching ratios.

¹⁰ Normalized to the sum of $B(\eta_c \rightarrow K^\pm K_S^0 \pi^\mp)$, $B(\eta_c \rightarrow \phi\phi)$, $B(\eta_c \rightarrow K^+ K^- \pi^+ \pi^-)$, and $B(\eta_c \rightarrow 2\pi^+ 2\pi^-)$.

¹¹ Superseded by ACCIARRI 99T.

¹²Normalized to the sum of $B(\eta_c \rightarrow K^\pm K_S^0 \pi^\mp)$, $B(\eta_c \rightarrow 2K^+ 2K^-)$, $B(\eta_c \rightarrow K^+ K^- \pi^+ \pi^-)$, and $B(\eta_c \rightarrow 2\pi^+ 2\pi^-)$.

¹³Re-evaluated by AIHARA 88D.

$\eta_c(1S) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(\eta'(958)\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_1\Gamma_{48}/\Gamma$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
75.8^{+6.3}_{-6.2} ± 8.4	486	¹ ZHANG	12A BELL	$e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$

¹ Assuming there is no interference with the non-resonant background.

$\Gamma(\rho\rho) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_2\Gamma_{48}/\Gamma$

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
------------	-----	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<39 90 < 1556 UEHARA 08 BELL $\gamma\gamma \rightarrow 2(\pi^+\pi^-)$

$\Gamma(K^*(892)\bar{K}^*(892)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_4\Gamma_{48}/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

35 ± 6 OUR FIT

32.4 ± 4.2 ± 5.8 882 ± 115 UEHARA 08 BELL $\gamma\gamma \rightarrow \pi^+\pi^-K^+K^-$

$\Gamma(\phi\phi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_7\Gamma_{48}/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

9.0 ± 0.8 OUR FIT

7.75 ± 0.66 ± 0.62 386 ± 31 ¹ LIU 12B BELL $\gamma\gamma \rightarrow 2(K^+K^-)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.8 ± 1.2 ± 1.3 132 ± 23 UEHARA 08 BELL $\gamma\gamma \rightarrow 2(K^+K^-)$

¹ Supersedes UEHARA 08. Using $B(\phi \rightarrow K^+K^-) = (48.9 \pm 0.5)\%$.

$\Gamma(\omega\omega) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{13}\Gamma_{48}/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

8.67 ± 2.86 ± 0.96 85 ± 29 ¹ LIU 12B BELL $\gamma\gamma \rightarrow 2(\pi^+\pi^-\pi^0)$

¹ Using $B(\omega \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7)\%$.

$\Gamma(\omega\phi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{14}\Gamma_{48}/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.49 90 ¹ LIU 12B BELL $\gamma\gamma \rightarrow K^+K^-\pi^+\pi^-\pi^0$

¹ Using $B(\phi \rightarrow K^+K^-) = (48.9 \pm 0.5)\%$ and $B(\omega \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7)\%$.

$\Gamma(f_2(1270)f_2(1270)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{15}\Gamma_{48}/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

49 ± 13 OUR FIT

69 ± 17 ± 12 3182 ± 766 UEHARA 08 BELL $\gamma\gamma \rightarrow 2(\pi^+\pi^-)$

$\Gamma(f_2(1270)f'_2(1525)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{16}\Gamma_{48}/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
49±9±13	1128±206	UEHARA	08 BELL	$\gamma\gamma \rightarrow \pi^+\pi^-K^+K^-$

$\Gamma(K\bar{K}\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{28}\Gamma_{48}/\Gamma$

VALUE (keV)	CL% EVTS	DOCUMENT ID	TECN	COMMENT
0.367±0.021 OUR FIT				
0.407±0.027 OUR AVERAGE				Error includes scale factor of 1.2.
0.374±0.009±0.031	14k	¹ LEES	10 BABR	$10.6 e^+e^- \rightarrow e^+e^-K_S^0K^\pm\pi^\mp$
0.407±0.022±0.028		^{2,3} ASNER	04 CLEO	$\gamma\gamma \rightarrow \eta_c \rightarrow K_S^0K^\pm\pi^\mp$
0.60 ±0.12 ±0.09	41	^{3,4} ABDALLAH	03J DLPH	$\gamma\gamma \rightarrow K_S^0K^\pm\pi^\mp$
1.47 ±0.87 ±0.27		³ SHIRAI	98 AMY	$\gamma\gamma \rightarrow \eta_c \rightarrow K^\pm K_S^0\pi^\mp$
0.84 ±0.21		³ ALBRECHT	94H ARG	$\gamma\gamma \rightarrow K^\pm K_S^0\pi^\mp$
0.60 ^{+0.23} _{-0.20}		³ CHEN	90B CLEO	$\gamma\gamma \rightarrow \eta_c K^\pm K_S^0\pi^\mp$
1.06 ±0.41 ±0.27	11	³ BRAUNSCH...	89 TASS	$\gamma\gamma \rightarrow K\bar{K}\pi$
1.5 ^{+0.60} _{-0.45} ±0.3	7	³ BERGER	86 PLUT	$\gamma\gamma \rightarrow K\bar{K}\pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.386±0.008±0.021	12k	⁵ DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K_S^0K^\pm\pi^\mp$
0.418±0.044±0.022		^{3,6} BRANDENB...	00B CLE2	$\gamma\gamma \rightarrow \eta_c \rightarrow K^\pm K_S^0\pi^\mp$
<0.63	95	³ BEHREND	89 CELL	$\gamma\gamma \rightarrow K_S^0K^\pm\pi^\mp$
<4.4	95	ALTHOFF	85B TASS	$\gamma\gamma \rightarrow K\bar{K}\pi$

¹ From the corrected and unfolded mass spectrum.

² Calculated by us from the value reported in ASNER 04 that assumes $B(\eta_c \rightarrow K\bar{K}\pi) = 5.5 \pm 1.7\%$

³ We have multiplied $K^\pm K_S^0\pi^\mp$ measurement by 3 to obtain $K\bar{K}\pi$.

⁴ Calculated by us from the value reported in ABDALLAH 03J, which uses $B(\eta_c \rightarrow K_S^0K^\pm\pi^\mp) = (1.5 \pm 0.4)\%$.

⁵ Not independent from the measurements reported by LEES 10.

⁶ Superseded by ASNER 04.

$\Gamma(K^+K^-\pi^+\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{32}\Gamma_{48}/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
35 ± 5 OUR FIT				
27 ± 6 OUR AVERAGE				
25.7± 3.2± 4.9	2019±248	UEHARA	08 BELL	$\gamma\gamma \rightarrow \pi^+\pi^-K^+K^-$
280 ±100 ±60	42	¹ ABDALLAH	03J DLPH	$\gamma\gamma \rightarrow \pi^+\pi^-K^+K^-$
170 ± 80 ±20	13.9 ± 6.6	ALBRECHT	94H ARG	$\gamma\gamma \rightarrow \pi^+\pi^-K^+K^-$

¹ Calculated by us from the value reported in ABDALLAH 03J, which uses $B(\eta_c \rightarrow \pi^+\pi^-K^+K^-) = (2.0 \pm 0.7)\%$.

$\Gamma(K^+K^-\pi^+\pi^-\pi^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{33}\Gamma_{48}/\Gamma$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.190 ± 0.006 ± 0.028 11k ¹ DEL-AMO-SA..11M BABR $\gamma\gamma \rightarrow K^+K^-\pi^+\pi^-\pi^0$

¹ Not independent from other measurements reported in DEL-AMO-SANCHEZ 11M.

 $\Gamma(2(K^+K^-)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{36}\Gamma_{48}/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

7.4 ± 1.5 OUR FIT

5.8 ± 1.9 OUR AVERAGE

5.6 ± 1.1 ± 1.6 216 ± 42 UEHARA 08 BELL $\gamma\gamma \rightarrow 2(K^+K^-)$

350 ± 90 ± 60 46 ¹ ABDALLAH 03J DLPH $\gamma\gamma \rightarrow 2(K^+K^-)$

231 ± 90 ± 23 9.1 ± 3.3 ² ALBRECHT 94H ARG $\gamma\gamma \rightarrow 2(K^+K^-)$

¹ Calculated by us from the value reported in ABDALLAH 03J, which uses $B(\eta_c \rightarrow)$

$2(K^+K^-) = (2.1 \pm 1.2)\%$.

² Includes all topological modes except $\eta_c \rightarrow \phi\phi$.

 $\Gamma(2(\pi^+\pi^-)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{39}\Gamma_{48}/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

49 ± 6 OUR FIT

42 ± 6 OUR AVERAGE

40.7 ± 3.7 ± 5.3 5381 ± 492 UEHARA 08 BELL $\gamma\gamma \rightarrow 2(\pi^+\pi^-)$

180 ± 70 ± 20 21.4 ± 8.6 ALBRECHT 94H ARG $\gamma\gamma \rightarrow 2(\pi^+\pi^-)$

 $\Gamma(p\bar{p}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{42}\Gamma_{48}/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

7.6 ± 0.7 OUR FIT

7.20 ± 1.53^{+0.67}_{-0.75} 157 ± 33 ¹ KUO 05 BELL $\gamma\gamma \rightarrow p\bar{p}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.6 ^{+1.3}_{-1.1} ± 0.4 190 ¹ AMBROGIANI 03 E835 $\bar{p}p \rightarrow \gamma\gamma$

8.1 ^{+2.9}_{-2.0} ¹ ARMSTRONG 95F E760 $\bar{p}p \rightarrow \gamma\gamma$

¹ Not independent from the $\Gamma_{\gamma\gamma}$ reported by the same experiment.

 $\Gamma(K_S^0K_S^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{52}\Gamma_{48}/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

<1.6 90 ¹ UEHARA 13 BELL $\gamma\gamma \rightarrow K_S^0K_S^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.29 90 ² UEHARA 13 BELL $\gamma\gamma \rightarrow K_S^0K_S^0$

¹ Taking into account interference with the non-resonant continuum.

² Neglecting interference with the non-resonant continuum.

$\eta_c(1S)$ BRANCHING RATIOS

HADRONIC DECAYS

 $\Gamma(\eta'(958)\pi\pi)/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.041±0.017	14	¹ BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c \gamma$

¹ The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 0.0127 \pm 0.0036$.

 $\Gamma(\rho\rho)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
18 ± 5 OUR AVERAGE					
12.6± 3.8±5.1		72	¹ ABLIKIM	05L BES2	$J/\psi \rightarrow \pi^+\pi^-\pi^+\pi^-\gamma$
26.0± 2.4±8.8		113	¹ BISELLO	91 DM2	$J/\psi \rightarrow \gamma\rho^0\rho^0$
23.6±10.6±8.2		32	¹ BISELLO	91 DM2	$J/\psi \rightarrow \gamma\rho^+\rho^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<14 90 ¹ BALTRUSAIT..86 MRK3 $J/\psi \rightarrow \eta_c \gamma$

¹ The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 0.0127 \pm 0.0036$. Where relevant, the error in this branching ratio is treated as a common systematic in computing averages.

 $\Gamma(K^*(892)^0 K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.02±0.007	63	^{1,2} BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c \gamma$

¹ BALTRUSAITIS 86 has an error according to Partridge.

² The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 0.0127 \pm 0.0036$.

 $\Gamma(K^*(892)\bar{K}^*(892))/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
71±13 OUR FIT				
91±26 OUR AVERAGE				
108±25±44	60	¹ ABLIKIM	05L BES2	$J/\psi \rightarrow K^+ K^- \pi^+ \pi^- \gamma$
82±28±27	14	¹ BISELLO	91 DM2	$e^+ e^- \rightarrow \gamma K^+ K^- \pi^+ \pi^-$
90±50	9	¹ BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c \gamma$

¹ The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 0.0127 \pm 0.0036$. Where relevant, the error in this branching ratio is treated as a common systematic in computing averages.

 $\Gamma(K^*(892)^0 \bar{K}^*(892)^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
113±47±24	45	¹ ABLIKIM	06A BES2	$J/\psi \rightarrow K^{*0} \bar{K}^{*0} \pi^+ \pi^- \gamma$

¹ ABLIKIM 06A reports $[\Gamma(\eta_c(1S) \rightarrow K^*(892)^0 \bar{K}^*(892)^0 \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] = (1.91 \pm 0.64 \pm 0.48) \times 10^{-4}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi K^+ K^-)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.9^{+0.9}_{-0.8} \pm 1.1$	$14.1^{+4.4}_{-3.7}$	¹ HUANG	03 BELL	$B^+ \rightarrow (\phi K^+ K^-) K^+$

¹ Using $B(B^+ \rightarrow \eta_c K^+) = (1.25 \pm 0.12^{+0.10}_{-0.12}) \times 10^{-3}$ from FANG 03 and $B(\eta_c \rightarrow K \bar{K} \pi) = (5.5 \pm 1.7) \times 10^{-2}$.

 $\Gamma(\phi\phi)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

17.9 ± 2.0 OUR FIT

28 ± 4 OUR AVERAGE

$26^{+4}_{-8} \pm 5$	1.2k	¹ ABLIKIM	17P BES3	$J/\psi \rightarrow K^+ K^- K^+ K^- \gamma$
$25.3 \pm 5.1 \pm 9.1$	72	² ABLIKIM	05L BES2	$J/\psi \rightarrow K^+ K^- K^+ K^- \gamma$
26 ± 9	357	² BAI	04 BES	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$
$31 \pm 7 \pm 10$	19	² BISELLO	91 DM2	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$
$30^{+18}_{-12} \pm 10$	5	² BISELLO	91 DM2	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$
$74 \pm 18 \pm 24$	80	² BAI	90B MRK3	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$
$67 \pm 21 \pm 24$		² BAI	90B MRK3	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$18^{+8}_{-6} \pm 7$	7	³ HUANG	03 BELL	$B^+ \rightarrow (\phi\phi) K^+$
----------------------	---	--------------------	---------	----------------------------------

¹ ABLIKIM 17P reports $[\Gamma(\eta_c(1S) \rightarrow \phi\phi)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] = (4.3 \pm 0.5^{+0.5}_{-1.2}) \times 10^{-5}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 0.0127 \pm 0.0036$. Where relevant, the error in this branching ratio is treated as a common systematic in computing averages.

³ Using $B(B^+ \rightarrow \eta_c K^+) = (1.25 \pm 0.12^{+0.10}_{-0.12}) \times 10^{-3}$ from FANG 03 and $B(\eta_c \rightarrow K \bar{K} \pi) = (5.5 \pm 1.7) \times 10^{-2}$.

 $\Gamma(\phi\phi)/\Gamma(K \bar{K} \pi)$ Γ_7/Γ_{28}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.0245 ± 0.0025 OUR FIT

0.044 $^{+0.012}_{-0.010}$ OUR AVERAGE

$0.055 \pm 0.014 \pm 0.005$		AUBERT,B	04B BABR	$B^\pm \rightarrow K^\pm \eta_c$
$0.032^{+0.014}_{-0.010} \pm 0.009$	7	¹ HUANG	03 BELL	$B^\pm \rightarrow K^\pm \phi\phi$

¹ Using $B(B^+ \rightarrow \eta_c K^+) = (1.25 \pm 0.12^{+0.10}_{-0.12}) \times 10^{-3}$ from FANG 03 and $B(\eta_c \rightarrow K \bar{K} \pi) = (5.5 \pm 1.7) \times 10^{-2}$.

 $\Gamma(\phi\phi)/\Gamma(p\bar{p})$ Γ_7/Γ_{42}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

1.79 ± 0.14 ± 0.32 6.4k ¹ AAIJ 17BB LHCB $pp \rightarrow b\bar{b}X \rightarrow 2(K^+ K^-)X$

¹ Using inputs from AAIJ 15AS and AAIJ 15BI and $\Gamma(b \rightarrow J/\psi(1S) \text{ anything})/\Gamma_{\text{total}} = (1.16 \pm 0.10)\%$ and $\Gamma(J/\psi(1S) \rightarrow p\bar{p})/\Gamma_{\text{total}} = (2.120 \pm 0.029) \times 10^{-3}$ from PDG 16.

$\Gamma(\phi 2(\pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<40	90	¹ ABLIKIM 06A	BES2	$J/\psi \rightarrow \phi 2(\pi^+ \pi^-) \gamma$
---------------	----	--------------------------	------	---

¹ ABLIKIM 06A reports $[\Gamma(\eta_c(1S) \rightarrow \phi 2(\pi^+ \pi^-))/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] < 0.603 \times 10^{-4}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 1.7 \times 10^{-2}$.

 $\Gamma(a_0(980)\pi)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<0.02	90	^{1,2} BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c \gamma$
-----------------	----	-------------------------------	------	------------------------------------

¹ The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 0.0127 \pm 0.0036$.

² We are assuming $B(a_0(980) \rightarrow \eta \pi) > 0.5$.

 $\Gamma(a_2(1320)\pi)/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<0.02	90	¹ BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c \gamma$
-----------------	----	-----------------------------	------	------------------------------------

¹ The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 0.0127 \pm 0.0036$.

 $\Gamma(K^*(892)\bar{K} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<0.0128	90	BISELLO 91	DM2	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
-------------------	----	------------	-----	---

<0.0132	90	¹ BISELLO 91	DM2	$J/\psi \rightarrow \gamma K^+ K^- \pi^0$
---------	----	-------------------------	-----	---

¹ The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 0.0127 \pm 0.0036$.

 $\Gamma(f_2(1270)\eta)/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<0.011	90	¹ BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c \gamma$
------------------	----	-----------------------------	------	------------------------------------

¹ The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 0.0127 \pm 0.0036$.

 $\Gamma(\omega\omega)/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<0.0031	90	¹ BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c \gamma$
-------------------	----	-----------------------------	------	------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0063	90	¹ ABLIKIM 05L	BES2	$J/\psi \rightarrow \pi^+ \pi^- \pi^0 \pi^+ \pi^- \pi^0 \gamma$
---------	----	--------------------------	------	---

<0.0063		¹ BISELLO 91	DM2	$J/\psi \rightarrow \gamma \omega \omega$
---------	--	-------------------------	-----	---

¹ The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 0.0127 \pm 0.0036$. Where relevant, the error in this branching ratio is treated as a common systematic in computing averages.

 $\Gamma(\omega\phi)/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

< 2.5×10^{-4}	90	¹ ABLIKIM 17P	BES3	$J/\psi \rightarrow \pi^+ \pi^- \pi^0 K^+ K^- \gamma$
---	----	--------------------------	------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 17×10^{-4}	90	² ABLIKIM 05L	BES2	$J/\psi \rightarrow \pi^+ \pi^- \pi^0 K^+ K^- \gamma$
-----------------------	----	--------------------------	------	---

¹ Using $B(J/\psi \rightarrow \gamma \eta_c) = 0.017 \pm 0.004$.

² The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 0.0127 \pm 0.0036$.

$\Gamma(f_2(1270)f_2(1270))/\Gamma_{\text{total}}$ Γ_{15}/Γ

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.98 ± 0.25 OUR FIT				
0.77^{+0.25}_{-0.30} ± 0.17	91.2 ± 19.8	¹ ABLIKIM	04M BES	$J/\psi \rightarrow \gamma 2\pi^+ 2\pi^-$

¹ ABLIKIM 04M reports $[\Gamma(\eta_c(1S) \rightarrow f_2(1270)f_2(1270))/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] = (1.3 \pm 0.3^{+0.3}_{-0.4}) \times 10^{-4}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(f_0(980)\eta)/\Gamma_{\text{total}}$ Γ_{17}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
seen	LEES	14E	BABR Dalitz anal. of $\eta_c \rightarrow K^+ K^- \eta$

 $\Gamma(f_0(1500)\eta)/\Gamma_{\text{total}}$ Γ_{18}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
seen	LEES	14E	BABR Dalitz anal. of $\eta_c \rightarrow K^+ K^- \eta$

 $\Gamma(f_0(2200)\eta)/\Gamma_{\text{total}}$ Γ_{19}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
seen	LEES	14E	BABR Dalitz anal. of $\eta_c \rightarrow K^+ K^- \eta$

 $\Gamma(a_0(980)\pi)/\Gamma_{\text{total}}$ Γ_{20}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
seen	LEES	14E	BABR Dalitz anal. of $\eta_c \rightarrow K^+ K^- \pi^0$

 $\Gamma(a_0(1320)\pi)/\Gamma_{\text{total}}$ Γ_{21}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
seen	LEES	14E	BABR Dalitz anal. of $\eta_c \rightarrow K^+ K^- \pi^0$

 $\Gamma(a_0(1450)\pi)/\Gamma_{\text{total}}$ Γ_{22}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
seen	LEES	14E	BABR Dalitz anal. of $\eta_c \rightarrow K^+ K^- \pi^0$

 $\Gamma(a_0(1950)\pi)/\Gamma_{\text{total}}$ Γ_{23}/Γ

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
seen	12k	¹ LEES	16A	BABR $\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K\bar{K}\pi$

¹ From a model-independent partial wave analysis.

 $\Gamma(a_2(1950)\pi)/\Gamma_{\text{total}}$ Γ_{24}/Γ

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
not seen	12k	¹ LEES	16A	BABR $\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K\bar{K}\pi$

¹ From a model-independent partial wave analysis assuming the existence of a hypothetical tensor isovector $a_2(1950)$.

$\Gamma(K_0^*(1430)\bar{K})/\Gamma_{\text{total}}$ Γ_{25}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	12k	¹ LEES	16A BABR	$\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K\bar{K}\pi$
seen		LEES	14E BABR	Dalitz anal. of $\eta_c \rightarrow K^+K^-\eta/\pi^0$

¹ From a model-independant partial wave analysis.

$\Gamma(K_2^*(1430)\bar{K})/\Gamma_{\text{total}}$ Γ_{26}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	LEES	14E BABR	Dalitz anal. of $\eta_c \rightarrow K^+K^-\pi^0$

$\Gamma(K_0^*(1950)\bar{K})/\Gamma_{\text{total}}$ Γ_{27}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	12K	¹ LEES	16A BABR	$\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K\bar{K}\pi$
seen		LEES	14E BABR	Dalitz anal. of $\eta_c \rightarrow K^+K^-\eta/\pi^0$

¹ From a Dalitz plot analysis using an isobar model.

$\Gamma(K\bar{K}\pi)/\Gamma_{\text{total}}$ Γ_{28}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
7.3 ± 0.5 OUR FIT				
6.5 ± 0.6 OUR AVERAGE				
6.3 ± 1.3 ± 0.6	55	1,2 ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0\gamma K^+K^-\pi^0$
7.9 ± 1.4 ± 0.7	107	3,4 ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0\gamma K_S^0 K^\mp\pi^\pm$
8.5 ± 1.8		5 AUBERT	06E BABR	$B^\pm \rightarrow K^\pm X_{c\bar{c}}$
5.1 ± 2.1	0.6k	6 BAI	04 BES	$J/\psi \rightarrow \gamma K^\pm\pi^\mp K_S^0$
6.90 ± 1.42 ± 1.32	33	6 BISELLO	91 DM2	$J/\psi \rightarrow \gamma K^+K^-\pi^0$
5.43 ± 0.94 ± 0.94	68	6 BISELLO	91 DM2	$J/\psi \rightarrow \gamma K^\pm\pi^\mp K_S^0$
4.8 ± 1.7	95	6,7 BALTRUSAIT..	86 MRK3	$J/\psi \rightarrow \eta_c\gamma$
16.1 ^{+9.2} / _{-7.3}		8,9 HIMEL	80B MRK2	$\psi(2S) \rightarrow \eta_c\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 10.7 90% CL ^{6,10} PARTRIDGE 80B CBAL $J/\psi \rightarrow \eta_c\gamma$

¹ ABLIKIM 12N quotes $B(\psi(2S) \rightarrow \pi^0 h_c) \cdot B(h_c \rightarrow \gamma\eta_c) \cdot B(\eta_c \rightarrow K^+K^-\pi^0) = (4.54 \pm 0.76 \pm 0.48) \times 10^{-6}$ which we multiply by 6 to account for isospin symmetry.

² ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow K\bar{K}\pi)/\Gamma_{\text{total}}] \times [\Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}}] = (27.24 \pm 4.56 \pm 2.88) \times 10^{-6}$ which we divide by our best value $\Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}} = (4.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ ABLIKIM 12N quotes $B(\psi(2S) \rightarrow \pi^0 h_c) \cdot B(h_c \rightarrow \gamma\eta_c) \cdot B(\eta_c \rightarrow K_S^0 K^\pm\pi^\mp) = (11.35 \pm 1.25 \pm 1.50) \times 10^{-6}$ which we multiply by 3 to account for isospin symmetry.

⁴ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow K\bar{K}\pi)/\Gamma_{\text{total}}] \times [\Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}}] = (34.05 \pm 3.75 \pm 4.50) \times 10^{-6}$ which we divide by our best value $\Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}} = (4.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁵ Determined from the ratio of $B(B^\pm \rightarrow K^\pm \eta_c) B(\eta_c \rightarrow K \bar{K} \pi) = (7.4 \pm 0.5 \pm 0.7) \times 10^{-5}$ reported in AUBERT, B 04B and $B(B^\pm \rightarrow K^\pm \eta_c) = (8.7 \pm 1.5) \times 10^{-3}$ reported in AUBERT 06E.

⁶ The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 0.0127 \pm 0.0036$. Where relevant, the error in this branching ratio is treated as a common systematic in computing averages.

⁷ Average from $K^+ K^- \pi^0$ and $K^\pm K_S^0 \pi^\mp$ decay channels.

⁸ $K^\pm K_S^0 \pi^\mp$ corrected to $K \bar{K} \pi$ by factor 3. KS, MR.

⁹ Estimated using $B(\psi(2S) \rightarrow \gamma \eta_c(1S)) = 0.0028 \pm 0.0006$.

¹⁰ $K^+ K^- \pi^0$ corrected to $K \bar{K} \pi$ by factor 6. KS, MR

$\Gamma(\phi K^+ K^-)/\Gamma(K \bar{K} \pi)$ Γ_6/Γ_{28}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$0.052^{+0.016}_{-0.014} \pm 0.014$	7	¹ HUANG	03 BELL	$B^\pm \rightarrow K^\pm \phi \phi$

¹ Using $B(B^+ \rightarrow \eta_c K^+) = (1.25 \pm 0.12^{+0.10}_{-0.12}) \times 10^{-3}$ from FANG 03 and $B(\eta_c \rightarrow K \bar{K} \pi) = (5.5 \pm 1.7) \times 10^{-2}$.

$\Gamma(K \bar{K} \eta)/\Gamma_{\text{total}}$ Γ_{29}/Γ

VALUE (units 10^{-2})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.36 ± 0.16 OUR FIT					

$1.0 \pm 0.5 \pm 0.2$ 7 ^{1,2} ABLIKIM 12N BES3 $\psi(2S) \rightarrow \pi^0 \gamma \eta K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<3.1 90 ³ BALTRUSAIT..86 MRK3 $J/\psi \rightarrow \eta_c \gamma$

¹ ABLIKIM 12N quotes $B(\psi(2S) \rightarrow \pi^0 h_c) \cdot B(h_c \rightarrow \gamma \eta_c) \cdot B(\eta_c \rightarrow K^+ K^- \eta) = (2.11 \pm 1.01 \pm 0.32) \times 10^{-6}$ which we multiply by 2 to account for isospin symmetry.

² ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow K \bar{K} \eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \pi^0 h_c(1P))] \times [B(h_c(1P) \rightarrow \gamma \eta_c(1S))] = (4.22 \pm 2.02 \pm 0.64) \times 10^{-6}$ which we divide by our best values $B(\psi(2S) \rightarrow \pi^0 h_c(1P)) = (8.6 \pm 1.3) \times 10^{-4}$, $B(h_c(1P) \rightarrow \gamma \eta_c(1S)) = (51 \pm 6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

³ The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 0.0127 \pm 0.0036$.

$\Gamma(K \bar{K} \eta)/\Gamma(K \bar{K} \pi)$ Γ_{29}/Γ_{28}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.186 ± 0.018 OUR FIT				

$0.190 \pm 0.008 \pm 0.017$ 5.4k ¹ LEES 14E BABR $\gamma \gamma \rightarrow K^+ K^- \eta/\pi^0$

¹ LEES 14E reports $B(\eta_c(1S) \rightarrow K^+ K^- \eta)/B(\eta_c(1S) \rightarrow K^+ K^- \pi^0) = 0.571 \pm 0.025 \pm 0.051$, which we divide by 3 to account for isospin symmetry. It uses both $\eta \rightarrow \gamma \gamma$ and $\eta \rightarrow \pi^+ \pi^- \pi^0$ decays.

$\Gamma(\eta \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{30}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.7 \pm 0.4 \pm 0.1$	33	¹ ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma \eta \pi^+ \pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.4 ± 2.0 75 ² BALTRUSAIT..86 MRK3 $J/\psi \rightarrow \eta_c \gamma$

$3.7 \pm 1.3 \pm 2.0$ 18 ² PARTRIDGE 80B CBAL $J/\psi \rightarrow \eta \pi^+ \pi^- \gamma$

¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow \eta \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [\Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}}] = (7.22 \pm 1.47 \pm 1.11) \times 10^{-6}$ which we divide by our best value $\Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}} = (4.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 0.0127 \pm 0.0036$. Where relevant, the error in this branching ratio is treated as a common systematic in computing averages.

$\Gamma(\eta 2(\pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{31}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
4.4 ± 1.2 ± 0.4	39	¹ ABLIKIM 12N	BES3	$\psi(2S) \rightarrow \pi^0 \gamma \eta 2(\pi^+ \pi^-)$

¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow \eta 2(\pi^+ \pi^-))/\Gamma_{\text{total}}] \times [\Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}}] = (19.17 \pm 3.77 \pm 3.72) \times 10^{-6}$ which we divide by our best value $\Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}} = (4.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{32}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

6.9 ± 1.1 OUR FIT

11.2 ± 1.9 OUR AVERAGE

9.7 ± 2.2 ± 0.9	38	¹ ABLIKIM 12N	BES3	$\psi(2S) \rightarrow \pi^0 \gamma K^+ K^- \pi^+ \pi^-$
12 ± 4	0.4k	² BAI 04	BES	$J/\psi \rightarrow \gamma K^+ K^- \pi^+ \pi^-$
21 ± 7	110	² BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c \gamma$
14 + ²² / ₉		³ HIMEL 80B	MRK2	$\psi(2S) \rightarrow \eta_c \gamma$

¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [\Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}}] = (4.16 \pm 0.76 \pm 0.59) \times 10^{-6}$ which we divide by our best value $\Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}} = (4.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 0.0127 \pm 0.0036$. Where relevant, the error in this branching ratio is treated as a common systematic in computing averages.

³ Estimated using $B(\psi(2S) \rightarrow \gamma \eta_c(1S)) = 0.0028 \pm 0.0006$.

$\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma(K \bar{K} \pi)$ Γ_{33}/Γ_{28}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.477 ± 0.017 ± 0.070	11k	¹ DEL-AMO-SA..11M	BABR	$\gamma \gamma \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

¹ We have multiplied the value of $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma(K_S^0 K^\pm \pi^\mp)$ reported in DEL-AMO-SANCHEZ 11M by a factor 1/3 to obtain $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma(K \bar{K} \pi)$. Not independent from other measurements reported in DEL-AMO-SANCHEZ 11M.

$\Gamma(K^0 K^- \pi^+ \pi^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{34}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
5.6 ± 1.4 ± 0.5	43	^{1,2} ABLIKIM 12N	BES3	$\psi(2S) \rightarrow \pi^0 \gamma K_S^0 K^\mp \pi^\mp 2\pi^\pm$

¹ ABLIKIM 12N quotes $B(\psi(2S) \rightarrow \pi^0 h_c) \cdot B(h_c \rightarrow \gamma \eta_c) \cdot B(\eta_c \rightarrow K_S^0 K^- \pi^- 2\pi^+)$
 $= (12.01 \pm 2.22 \pm 2.04) \times 10^{-6}$ which we multiply by 2 to take c.c. into account.

² ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow K^0 K^- \pi^+ \pi^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}] \times [\Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}}] = (24.02 \pm 4.44 \pm 4.08) \times 10^{-6}$
 which we divide by our best value $\Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}} = (4.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^+ K^- 2(\pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{35}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

7.5 ± 2.4 OUR AVERAGE

8 ± 4 ± 1	10	¹ ABLIKIM 12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma K^+ K^- 2(\pi^+ \pi^-)$
7.2 ± 2.4 ± 1.5	100	² ABLIKIM 06A BES2	$J/\psi \rightarrow K^+ K^- 2(\pi^+ \pi^-) \gamma$

¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow K^+ K^- 2(\pi^+ \pi^-))/\Gamma_{\text{total}}] \times [\Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}}] = (3.60 \pm 1.71 \pm 0.64) \times 10^{-6}$
 which we divide by our best value $\Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}} = (4.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ABLIKIM 06A reports $[\Gamma(\eta_c(1S) \rightarrow K^+ K^- 2(\pi^+ \pi^-))/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] = (1.21 \pm 0.32 \pm 0.24) \times 10^{-4}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2(K^+ K^-))/\Gamma_{\text{total}}$ Γ_{36}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

1.47 ± 0.31 OUR FIT

2.2 ± 0.9 ± 0.2	7	¹ ABLIKIM 12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma 2(K^+ K^-)$
------------------------	---	-------------------------------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.4 $\begin{smallmatrix} +0.5 \\ -0.4 \end{smallmatrix}$ ± 0.6	14.5 $\begin{smallmatrix} +4.6 \\ -3.0 \end{smallmatrix}$	² HUANG 03 BELL	$B^+ \rightarrow 2(K^+ K^-) K^+$
21 ± 10 ± 6		³ ALBRECHT 94H ARG	$\gamma \gamma \rightarrow K^+ K^- K^+ K^-$

¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow 2(K^+ K^-))/\Gamma_{\text{total}}] \times [\Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}}] = (0.94 \pm 0.37 \pm 0.14) \times 10^{-6}$ which we divide by our best value $\Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}} = (4.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Using $B(B^+ \rightarrow \eta_c K^+) = (1.25 \pm 0.12 \pm 0.10) \times 10^{-3}$ from FANG 03 and $B(\eta_c \rightarrow K \bar{K} \pi) = (5.5 \pm 1.7) \times 10^{-2}$.

³ Normalized to the sum of $B(\eta_c \rightarrow K^\pm K_S^0 \pi^\mp)$, $B(\eta_c \rightarrow \phi \phi)$, $B(\eta_c \rightarrow K^+ K^- \pi^+ \pi^-)$, and $B(\eta_c \rightarrow 2\pi^+ 2\pi^-)$.

$\Gamma(2(K^+ K^-))/\Gamma(K \bar{K} \pi)$ Γ_{36}/Γ_{28}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.020 ± 0.004 OUR FIT

0.024 ± 0.007 OUR AVERAGE

0.023 ± 0.007 ± 0.006		AUBERT,B 04B BABR	$B^\pm \rightarrow K^\pm \eta_c$
0.026 $\begin{smallmatrix} +0.009 \\ -0.007 \end{smallmatrix}$ ± 0.007	15	¹ HUANG 03 BELL	$B^\pm \rightarrow K^\pm (2K^+ 2K^-)$

¹ Using $B(B^+ \rightarrow \eta_c K^+) = (1.25 \pm 0.12_{-0.12}^{+0.10}) \times 10^{-3}$ from FANG 03 and $B(\eta_c \rightarrow K \bar{K} \pi) = (5.5 \pm 1.7) \times 10^{-2}$.

$\Gamma(\pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$ Γ_{37}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5 \times 10^{-4}$	90	¹ ABLIKIM	17AJ BES3	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- \pi^0$

¹ ABLIKIM 17AJ reports $[\Gamma(\eta_c(1S) \rightarrow \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \eta_c(1S))]$
 $< 1.6 \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \eta_c(1S)) = 3.4 \times 10^{-3}$.

$\Gamma(\pi^+ \pi^- \pi^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{38}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$4.7 \pm 0.9 \pm 0.4$	118	¹ ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma \pi^+ \pi^- 2\pi^0$

¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow \pi^+ \pi^- \pi^0 \pi^0)/\Gamma_{\text{total}}] \times [\Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}}] = (20.31 \pm 2.20 \pm 3.33) \times 10^{-6}$ which we divide by our best value $\Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}} = (4.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2(\pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{39}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
0.97 ± 0.12 OUR FIT				
1.35 ± 0.21 OUR AVERAGE				

1.74 \pm 0.32 \pm 0.15	100	¹ ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma 2(\pi^+ \pi^-)$
1.0 \pm 0.5	542 \pm 75	² BAI	04 BES	$J/\psi \rightarrow \gamma 2(\pi^+ \pi^-)$
1.05 \pm 0.17 \pm 0.34	137	² BISELLO	91 DM2	$J/\psi \rightarrow \gamma 2\pi^+ 2\pi^-$
1.3 \pm 0.6	25	² BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c \gamma$
2.0 $^{+1.5}_{-1.0}$		³ HIMEL	80B MRK2	$\psi(2S) \rightarrow \eta_c \gamma$

¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow 2(\pi^+ \pi^-))/\Gamma_{\text{total}}] \times [\Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}}] = (7.51 \pm 0.85 \pm 1.11) \times 10^{-6}$ which we divide by our best value $\Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}} = (4.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 0.0127 \pm 0.0036$. Where relevant, the error in this branching ratio is treated as a common systematic in computing averages.

³ Estimated using $B(\psi(2S) \rightarrow \gamma \eta_c(1S)) = 0.0028 \pm 0.0006$.

$\Gamma(2(\pi^+ \pi^- \pi^0))/\Gamma_{\text{total}}$ Γ_{40}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$17.4 \pm 2.9 \pm 1.5$	175	¹ ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma 2(\pi^+ \pi^- 2\pi^0)$

¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow 2(\pi^+ \pi^- \pi^0))/\Gamma_{\text{total}}] \times [\Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}}] = (75.13 \pm 7.42 \pm 9.99) \times 10^{-6}$ which we divide by our best value $\Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}} = (4.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(3(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{41}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
18 ± 4 OUR AVERAGE				
20 ± 5 ± 2	51	¹ ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma 3(\pi^+\pi^-)$
15.4 ± 3.4 ± 3.3	479	² ABLIKIM	06A BES2	$J/\psi \rightarrow 3(\pi^+\pi^-)\gamma$

¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow 3(\pi^+\pi^-))/\Gamma_{\text{total}}] \times [\Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}}] = (8.82 \pm 1.57 \pm 1.59) \times 10^{-6}$ which we divide by our best value $\Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}} = (4.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ABLIKIM 06A reports $[\Gamma(\eta_c(1S) \rightarrow 3(\pi^+\pi^-))/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] = (2.59 \pm 0.32 \pm 0.47) \times 10^{-4}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(p\bar{p})/\Gamma_{\text{total}}$ Γ_{42}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
15.2 ± 1.6 OUR FIT				
13.2 ± 2.7 OUR AVERAGE				
15 ± 5 ± 1	15	¹ ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma p\bar{p}$
15 ± 6	213 ± 33	² BAI	04 BES	$J/\psi \rightarrow \gamma p\bar{p}$
10 ± 3 ± 4	18	² BISELLO	91 DM2	$J/\psi \rightarrow \gamma p\bar{p}$
11 ± 6	23	² BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c \gamma$
29 ⁺²⁹ / ₋₁₅		³ HIMEL	80B MRK2	$\psi(2S) \rightarrow \eta_c \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

13.1 ^{+1.8+1.0} / _{-2.1-1.1}	195	⁴ WU	06 BELL	$B^+ \rightarrow p\bar{p}K^+$
--	-----	-----------------	---------	-------------------------------

¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow p\bar{p})/\Gamma_{\text{total}}] \times [\Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}}] = (0.65 \pm 0.19 \pm 0.10) \times 10^{-6}$ which we divide by our best value $\Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}} = (4.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 0.0127 \pm 0.0036$. Where relevant, the error in this branching ratio is treated as a common systematic in computing averages.

³ Estimated using $B(\psi(2S) \rightarrow \gamma\eta_c(1S)) = 0.0028 \pm 0.0006$.

⁴ WU 06 reports $[\Gamma(\eta_c(1S) \rightarrow p\bar{p})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \eta_c K^+)] = (1.42 \pm 0.11 ^{+0.16}/_{-0.20}) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \eta_c K^+) = (1.09 \pm 0.09) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(p\bar{p})/\Gamma(K\bar{K}\pi)$ Γ_{42}/Γ_{28}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0207 ± 0.0021 OUR FIT				
0.021 ± 0.002 ^{+0.004}/_{-0.006}	195	¹ WU	06 BELL	$B^\pm \rightarrow K^\pm p\bar{p}$

¹ Using $B(B^+ \rightarrow \eta_c K^+) = (1.25 \pm 0.12 ^{+0.10}/_{-0.12}) \times 10^{-3}$ from FANG 03 and $B(\eta_c \rightarrow K\bar{K}\pi) = (5.5 \pm 1.7) \times 10^{-2}$.

$\Gamma(\rho\bar{\rho})/\Gamma_{\text{total}} \times \Gamma(\phi\phi)/\Gamma_{\text{total}}$		$\Gamma_{42}/\Gamma \times \Gamma_7/\Gamma$		
VALUE (units 10^{-5})		DOCUMENT ID	TECN	COMMENT
0.27±0.05 OUR FIT				
4.0^{+3.5}_{-3.2}		BAGLIN	89	SPEC $\bar{p}p \rightarrow K^+ K^- K^+ K^-$

$\Gamma(\rho\bar{\rho}\pi^0)/\Gamma_{\text{total}}$		Γ_{43}/Γ		
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
0.36±0.13±0.03	14	¹ ABLIKIM	12N	BES3 $\psi(2S) \rightarrow \pi^0 \gamma \rho\bar{\rho}\pi^0$

¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow \rho\bar{\rho}\pi^0)/\Gamma_{\text{total}}] \times [\Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}}] = (1.53 \pm 0.49 \pm 0.23) \times 10^{-6}$ which we divide by our best value $\Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}} = (4.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$			Γ_{44}/Γ		
VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
10.9±2.4 OUR FIT					
11.7±2.3±2.5			¹ ABLIKIM	12B	BES3

• • • We do not use the following data for averages, fits, limits, etc. • • •

$8.8^{+2.4}_{-2.3} \pm 0.7$	20	² WU	06	BELL	$B^+ \rightarrow \Lambda\bar{\Lambda}K^+$
<20	90	³ BISELLO	91	DM2	$e^+e^- \rightarrow \gamma\Lambda\bar{\Lambda}$

¹ ABLIKIM 12B reports $[\Gamma(\eta_c(1S) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] = (0.198 \pm 0.021 \pm 0.032) \times 10^{-4}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² WU 06 reports $[\Gamma(\eta_c(1S) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \eta_c K^+)] = (0.95^{+0.25+0.08}_{-0.22-0.11}) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \eta_c K^+) = (1.09 \pm 0.09) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 0.0127 \pm 0.0036$.

$\Gamma(\Lambda\bar{\Lambda})/\Gamma(\rho\bar{\rho})$		Γ_{44}/Γ_{42}		
VALUE		DOCUMENT ID	TECN	COMMENT
0.72±0.16 OUR FIT				
0.67^{+0.19}_{-0.16} ± 0.12		¹ WU	06	BELL $B^+ \rightarrow \rho\bar{\rho}K^+, \Lambda\bar{\Lambda}K^+$

¹ Not independent from other $\eta_c \rightarrow \Lambda\bar{\Lambda}, \rho\bar{\rho}$ branching ratios reported by WU 06.

$\Gamma(\Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}$		Γ_{45}/Γ		
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.1±0.3±0.5	112	¹ ABLIKIM	13C	BES3 $J/\psi \rightarrow \gamma\rho\bar{\rho}\pi^0\pi^0$

¹ ABLIKIM 13C reports $[\Gamma(\eta_c(1S) \rightarrow \Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] = (3.60 \pm 0.48 \pm 0.31) \times 10^{-5}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\Xi^- \Xi^+)/\Gamma_{\text{total}} \qquad \Gamma_{46}/\Gamma$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.90±0.18±0.19	78	¹ ABLIKIM	13C BES3	$J/\psi \rightarrow \gamma \Lambda \bar{\Lambda} \pi^+ \pi^-$

¹ ABLIKIM 13C reports $[\Gamma(\eta_c(1S) \rightarrow \Xi^- \Xi^+)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] = (1.51 \pm 0.27 \pm 0.14) \times 10^{-5}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\pi^+ \pi^- \rho \bar{\rho})/\Gamma_{\text{total}} \qquad \Gamma_{47}/\Gamma$$

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
5.3±1.7±0.5		19	¹ ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma \rho \bar{\rho} \pi^+ \pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<12	90	HIMEL	80B MRK2	$\psi(2S) \rightarrow \eta_c \gamma$
-----	----	-------	----------	--------------------------------------

¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow \pi^+ \pi^- \rho \bar{\rho})/\Gamma_{\text{total}}] \times [\Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}}] = (2.30 \pm 0.65 \pm 0.36) \times 10^{-6}$ which we divide by our best value $\Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}} = (4.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

————— RADIATIVE DECAYS —————

$$\Gamma(\gamma\gamma)/\Gamma_{\text{total}} \qquad \Gamma_{48}/\Gamma$$

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.57±0.12 OUR FIT					

1.9 $\begin{smallmatrix} +0.7 \\ -0.6 \end{smallmatrix}$ OUR AVERAGE

2.7 ±0.8 ±0.6			¹ ABLIKIM	13I BES3	
1.4 $\begin{smallmatrix} +0.7 \\ -0.5 \end{smallmatrix}$ ±0.3		1.2 $\begin{smallmatrix} +2.8 \\ -1.1 \end{smallmatrix}$	² ADAMS	08 CLEO	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.0 $\begin{smallmatrix} +0.9 \\ -0.7 \end{smallmatrix}$ ±0.2		13	³ WICHT	08 BELL	$B^\pm \rightarrow K^\pm \gamma \gamma$
2.80 $\begin{smallmatrix} +0.67 \\ -0.58 \end{smallmatrix}$ ±1.0			⁴ ARMSTRONG	95F E760	$\bar{p} p \rightarrow \gamma \gamma$
< 9	90		⁵ BISELLO	91 DM2	$J/\psi \rightarrow \gamma \gamma \gamma$
6 $\begin{smallmatrix} +4 \\ -3 \end{smallmatrix}$ ±4			⁴ BAGLIN	87B SPEC	$\bar{p} p \rightarrow \gamma \gamma$
< 18	90		⁶ BLOOM	83 CBAL	$J/\psi \rightarrow \eta_c \gamma$

¹ ABLIKIM 13I reports $[\Gamma(\eta_c(1S) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] = (4.5 \pm 1.2 \pm 0.6) \times 10^{-6}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ADAMS 08 reports $[\Gamma(\eta_c(1S) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] = (2.4 $\begin{smallmatrix} +1.1 \\ -0.8 \end{smallmatrix}$ ± 0.3) $\times 10^{-6}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.$

³ WICHT 08 reports $[\Gamma(\eta_c(1S) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \eta_c K^+)] = (2.2 $\begin{smallmatrix} +0.9+0.4 \\ -0.7-0.2 \end{smallmatrix}$) $\times 10^{-7}$ which we divide by our best value $B(B^+ \rightarrow \eta_c K^+) =$$

$(1.09 \pm 0.09) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ Not independent from the values of the total and two-photon width quoted by the same experiment.

⁵ The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 0.0127 \pm 0.0036$.

⁶ Using $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 0.0127 \pm 0.0036$.

$\Gamma(\gamma\gamma)/\Gamma(K\bar{K}\pi)$					Γ_{48}/Γ_{28}
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
2.15±0.28 OUR FIT					
3.2 $\begin{smallmatrix} +1.3 \\ -1.0 \end{smallmatrix}$ $\begin{smallmatrix} +0.8 \\ -0.6 \end{smallmatrix}$	13	¹ WICHT	08	BELL	$B^\pm \rightarrow K^\pm \gamma\gamma$

¹ Using $B(B^+ \rightarrow \eta_c K^+) = (1.25 \pm 0.12^{+0.10}_{-0.12}) \times 10^{-3}$ from FANG 03 and $B(\eta_c \rightarrow K\bar{K}\pi) = (5.5 \pm 1.7) \times 10^{-2}$.

$\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_{42}/\Gamma \times \Gamma_{48}/\Gamma$
VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT	
0.238±0.023 OUR FIT					
0.26 ±0.05 OUR AVERAGE Error includes scale factor of 1.4.					
$0.224^{+0.038}_{-0.037} \pm 0.020$	190	AMBROGIANI 03	E835	$\bar{p}p \rightarrow \eta_c \rightarrow \gamma\gamma$	
$0.336^{+0.080}_{-0.070}$		ARMSTRONG 95F	E760	$\bar{p}p \rightarrow \gamma\gamma$	
$0.68^{+0.42}_{-0.31}$	12	BAGLIN	87B	SPEC	$\bar{p}p \rightarrow \gamma\gamma$

———— Charge conjugation (C), Parity (P), ————
 ———— Lepton family number (LF) violating modes ————

$\Gamma(\pi^+ \pi^-)/\Gamma_{\text{total}}$					Γ_{49}/Γ
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	
<11	90	¹ ABLIKIM	11G	BES3	$J/\psi \rightarrow \gamma\pi^+\pi^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
<70	90	² ABLIKIM	06B	BES2	$J/\psi \rightarrow \pi^+\pi^-\gamma$
¹ ABLIKIM 11G reports $[\Gamma(\eta_c(1S) \rightarrow \pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] < 1.82 \times 10^{-6}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 1.7 \times 10^{-2}$.					
² ABLIKIM 06B reports $[\Gamma(\eta_c(1S) \rightarrow \pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] < 1.1 \times 10^{-5}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 1.7 \times 10^{-2}$.					

$\Gamma(\pi^0 \pi^0)/\Gamma_{\text{total}}$					Γ_{50}/Γ
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	
< 4	90	¹ ABLIKIM	11G	BES3	$J/\psi \rightarrow \gamma\pi^0\pi^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
<40	90	² ABLIKIM	06B	BES2	$J/\psi \rightarrow \pi^0\pi^0\gamma$
¹ ABLIKIM 11G reports $[\Gamma(\eta_c(1S) \rightarrow \pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] < 6.0 \times 10^{-7}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 1.7 \times 10^{-2}$.					
² ABLIKIM 06B reports $[\Gamma(\eta_c(1S) \rightarrow \pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] < 0.71 \times 10^{-5}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 1.7 \times 10^{-2}$.					

$\Gamma(K^+K^-)/\Gamma_{\text{total}}$ Γ_{51}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<60	90	¹ ABLIKIM	06B BES2	$J/\psi \rightarrow K^+K^- \gamma$

¹ ABLIKIM 06B reports $[\Gamma(\eta_c(1S) \rightarrow K^+K^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))]$
 $< 0.96 \times 10^{-5}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 1.7 \times 10^{-2}$.

 $\Gamma(K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{52}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<31	90	¹ ABLIKIM	06B BES2	$J/\psi \rightarrow K_S^0 K_S^0 \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<32	90	² UEHARA	13 BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$
< 5.6	90	³ UEHARA	13 BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$

¹ ABLIKIM 06B reports $[\Gamma(\eta_c(1S) \rightarrow K_S^0 K_S^0)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))]$
 $< 0.53 \times 10^{-5}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 1.7 \times 10^{-2}$.

² Taking into account interference with the non-resonant continuum.

³ Neglecting interference with the non-resonant continuum.

 $\eta_c(1S)$ REFERENCES

AAIJ	17AD	PL B769 305	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17BB	EPJ C77 609	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	17AJ	PR D96 112008	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	17P	PR D95 092004	M. Ablikim <i>et al.</i>	(BES III Collab.)
LEES	16A	PR D93 012005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
PDG	16	CP C40 100001	C. Patrignani <i>et al.</i>	(PDG Collab.)
AAIJ	15AS	JHEP 1510 053	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	15BI	EPJ C75 311	R. Aaij <i>et al.</i>	(LHCb Collab.)
ANASHIN	14	PL B738 391	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
LEES	14E	PR D89 112004	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	13C	PR D87 012003	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	13I	PR D87 032003	M. Ablikim <i>et al.</i>	(BES III Collab.)
UEHARA	13	PTEP 2013 123C01	S. Uehara <i>et al.</i>	(BELLE Collab.)
ABLIKIM	12B	PR D86 032008	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	12F	PRL 108 222002	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	12N	PR D86 092009	M. Ablikim <i>et al.</i>	(BES III Collab.)
LIU	12B	PRL 108 232001	Z.Q. Liu <i>et al.</i>	(BELLE Collab.)
ZHANG	12A	PR D86 052002	C.C. Zhang <i>et al.</i>	(BELLE Collab.)
ABLIKIM	11G	PR D84 032006	M. Ablikim <i>et al.</i>	(BES III Collab.)
DEL-AMO-SA...	11M	PR D84 012004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
VINOKUROVA	11	PL B706 139	A. Vinokurova <i>et al.</i>	(BELLE Collab.)
LEES	10	PR D81 052010	J.P. Lees <i>et al.</i>	(BABAR Collab.)
MITCHELL	09	PRL 102 011801	R.E. Mitchell <i>et al.</i>	(CLEO Collab.)
ADAMS	08	PRL 101 101801	G.S. Adams <i>et al.</i>	(CLEO Collab.)
AUBERT	08AB	PR D78 012006	B. Aubert <i>et al.</i>	(BABAR Collab.)
UEHARA	08	EPJ C53 1	S. Uehara <i>et al.</i>	(BELLE Collab.)
WICHT	08	PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)
ABE	07	PRL 98 082001	K. Abe <i>et al.</i>	(BELLE Collab.)
ABLIKIM	06A	PL B633 19	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06B	EPJ C45 337	M. Ablikim <i>et al.</i>	(BES Collab.)
AUBERT	06E	PRL 96 052002	B. Aubert <i>et al.</i>	(BABAR Collab.)
PDG	06	JP G33 1	W.-M. Yao <i>et al.</i>	(PDG Collab.)
WU	06	PRL 97 162003	C.-H. Wu <i>et al.</i>	(BELLE Collab.)
ABLIKIM	05L	PR D72 072005	M. Ablikim <i>et al.</i>	(BES Collab.)
KUO	05	PL B621 41	C.C. Kuo <i>et al.</i>	(BELLE Collab.)
ABE	04G	PR D70 071102	K. Abe <i>et al.</i>	(BELLE Collab.)
ABLIKIM	04M	PR D70 112008	M. Ablikim <i>et al.</i>	(BES Collab.)
ASNER	04	PRL 92 142001	D.M. Asner <i>et al.</i>	(CLEO Collab.)
AUBERT	04D	PRL 92 142002	B. Aubert <i>et al.</i>	(BABAR Collab.)

AUBERT,B	04B	PR D70 011101	B. Aubert <i>et al.</i>	(BABAR Collab.)
BAI	04	PL B578 16	J.Z. Bai <i>et al.</i>	(BES Collab.)
ABDALLAH	03J	EPJ C31 481	J. Abdallah <i>et al.</i>	(DELPHI Collab.)
AMBROGIANI	03	PL B566 45	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)
BAI	03	PL B555 174	J.Z. Bai <i>et al.</i>	(BES Collab.)
FANG	03	PRL 90 071801	F. Fang <i>et al.</i>	(BELLE Collab.)
HUANG	03	PRL 91 241802	H.-C. Huang <i>et al.</i>	(BELLE Collab.)
ABE,K	02	PRL 89 142001	K. Abe <i>et al.</i>	(BELLE Collab.)
BAI	00F	PR D62 072001	J.Z. Bai <i>et al.</i>	(BES Collab.)
BRANDENB...	00B	PRL 85 3095	G. Brandenburg <i>et al.</i>	(CLEO Collab.)
ACCIARRI	99T	PL B461 155	M. Acciarri <i>et al.</i>	(L3 Collab.)
BAI	99B	PR D60 072001	J.Z. Bai <i>et al.</i>	(BES Collab.)
ABREU	98O	PL B441 479	P. Abreu <i>et al.</i>	(DELPHI Collab.)
SHIRAI	98	PL B424 405	M. Shirai <i>et al.</i>	(AMY Collab.)
ARMSTRONG	95F	PR D52 4839	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
ALBRECHT	94H	PL B338 390	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ADRIANI	93N	PL B318 575	O. Adriani <i>et al.</i>	(L3 Collab.)
BISELLO	91	NP B350 1	D. Bisello <i>et al.</i>	(DM2 Collab.)
BAI	90B	PRL 65 1309	Z. Bai <i>et al.</i>	(Mark III Collab.)
CHEN	90B	PL B243 169	W.Y. Chen <i>et al.</i>	(CLEO Collab.)
BAGLIN	89	PL B231 557	C. Baglin, S. Baird, G. Bassompierre	(R704 Collab.)
BEHREND	89	ZPHY C42 367	H.J. Behrend <i>et al.</i>	(CELLO Collab.)
BRAUNSCH...	89	ZPHY C41 533	W. Braunschweig <i>et al.</i>	(TASSO Collab.)
AIHARA	88D	PRL 60 2355	H. Aihara <i>et al.</i>	(TPC Collab.)
BAGLIN	87B	PL B187 191	C. Baglin <i>et al.</i>	(R704 Collab.)
BALTRUSAIT...	86	PR D33 629	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)
BERGER	86	PL 167B 120	C. Berger <i>et al.</i>	(PLUTO Collab.)
GAISER	86	PR D34 711	J. Gaiser <i>et al.</i>	(Crystal Ball Collab.)
ALTHOFF	85B	ZPHY C29 189	M. Althoff <i>et al.</i>	(TASSO Collab.)
BALTRUSAIT...	84	PRL 52 2126	R.M. Baltrusaitis <i>et al.</i>	(CIT, UCSC+) JP
BLOOM	83	ARNS 33 143	E.D. Bloom, C. Peck	(SLAC, CIT)
HIMEL	80B	PRL 45 1146	T.M. Himel <i>et al.</i>	(SLAC, LBL, UCB)
PARTRIDGE	80B	PRL 45 1150	R. Partridge <i>et al.</i>	(CIT, HARV, PRIN+)