

**$\psi(4230)$** 

$$I^G(J^{PC}) = 0^-(1^{--})$$

OMITTED FROM SUMMARY TABLE  
was  $X(4230)$

This state shows properties different from a conventional  $q\bar{q}$  state.  
A candidate for an exotic structure. See the review on non- $q\bar{q}$  states.

Enhancement reported by ABLIKIM 15C in  $e^+e^- \rightarrow \omega\chi_{c0}$  at  $\sqrt{s} = 4.23\text{--}4.26$  GeV at  $9\sigma$  significance. Lineshape found to be inconsistent with origination from  $\psi(4260)$ . Needs confirmation.

 **$\psi(4230)$  MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4218 <math>\begin{smallmatrix} +5 \\ -4 \end{smallmatrix}</math> OUR AVERAGE</b>		Error includes scale factor of 1.2.		
4218 $\begin{smallmatrix} +5.5 \\ -4.5 \end{smallmatrix} \pm 0.9$		ABLIKIM	17G BES3	$e^+e^- \rightarrow \pi^+\pi^-h_c$
4209.5 $\pm 7.4 \pm 1.4$		<sup>1</sup> ABLIKIM	17V BES3	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$
4230 $\pm 8 \pm 6$	180	<sup>2</sup> ABLIKIM	15C BES3	$e^+e^- \rightarrow \omega\chi_{c0}$

<sup>1</sup> From a fit to the cross section for  $e^+e^- \rightarrow \pi^+\pi^-\psi(2S) \rightarrow 2(\pi^+\pi^-)\ell^+\ell^-$  obtained from 16 center-of-mass energies between 4.008 and 4.600 GeV and comprising  $5.1 \text{ fb}^{-1}$ .

<sup>2</sup> From a 3-parameter fit of measured cross sections from  $\sqrt{s} = 4.21\text{--}4.42$  GeV to a phase-space modified Breit-Wigner function, using the decays  $\chi_{c0} \rightarrow \pi^+\pi^-$ ,  $\chi_{c0} \rightarrow K^+K^-$ , and  $\omega \rightarrow \pi^+\pi^-\pi^0$ .


 **$\psi(4230)$  WIDTH**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>59 <math>\begin{smallmatrix} +12 \\ -10 \end{smallmatrix}</math> OUR AVERAGE</b>		Error includes scale factor of 1.5. See the ideogram below.		
66.0 $\begin{smallmatrix} +12.3 \\ -8.3 \end{smallmatrix} \pm 0.4$		ABLIKIM	17G BES3	$e^+e^- \rightarrow \pi^+\pi^-h_c$
80.1 $\pm 24.6 \pm 2.9$		<sup>1</sup> ABLIKIM	17V BES3	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$
38 $\pm 12 \pm 2$	180	<sup>2</sup> ABLIKIM	15C BES3	$e^+e^- \rightarrow \omega\chi_{c0}$

<sup>1</sup> From a fit to the cross section for  $e^+e^- \rightarrow \pi^+\pi^-\psi(2S) \rightarrow 2(\pi^+\pi^-)\ell^+\ell^-$  obtained from 16 center-of-mass energies between 4.008 and 4.600 GeV and comprising  $5.1 \text{ fb}^{-1}$ .

<sup>2</sup> From a 3-parameter fit of measured cross sections from  $\sqrt{s} = 4.21\text{--}4.42$  GeV to a phase-space modified Breit-Wigner function, using the decays  $\chi_{c0} \rightarrow \pi^+\pi^-$ ,  $\chi_{c0} \rightarrow K^+K^-$ , and  $\omega \rightarrow \pi^+\pi^-\pi^0$ .

WEIGHTED AVERAGE  
59+12-10 (Error scaled by 1.5)



### $\psi(4230)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $e^+ e^-$	
$\Gamma_2$ $\omega \chi_{c0}$	seen
$\Gamma_3$ $\pi^+ \pi^- h_c$	seen
$\Gamma_4$ $\pi^+ \pi^- \psi(2S)$	seen

### $\psi(4230)$ $\Gamma(i)\Gamma(e^+ e^-)/\Gamma(\text{total})$

$\Gamma(\omega \chi_{c0}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_2\Gamma_1/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
<b><math>2.7 \pm 0.5 \pm 0.4</math></b>	180	<sup>1</sup> ABLIKIM	15C BES3	$e^+ e^- \rightarrow \omega \chi_{c0}$	

<sup>1</sup> From a 3-parameter fit of measured cross sections from  $\sqrt{s} = 4.21\text{--}4.42$  GeV to a phase-space modified Breit-Wigner function, using the decays  $\chi_{c0} \rightarrow \pi^+ \pi^-$ ,  $\chi_{c0} \rightarrow K^+ K^-$ , and  $\omega \rightarrow \pi^+ \pi^- \pi^0$ .

### $\psi(4230)$ BRANCHING RATIOS

$\Gamma(\omega \chi_{c0})/\Gamma_{\text{total}}$					$\Gamma_2/\Gamma$
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>seen</b>	180	<sup>1</sup> ABLIKIM	15C BES3	$e^+ e^- \rightarrow \omega \chi_{c0}$	

<sup>1</sup>From a 3-parameter fit of measured cross sections from  $\sqrt{s} = 4.21\text{--}4.42$  GeV to a phase-space modified Breit-Wigner function, using the decays  $\chi_{c0} \rightarrow \pi^+\pi^-$ ,  $\chi_{c0} \rightarrow K^+K^-$ , and  $\omega \rightarrow \pi^+\pi^-\pi^0$ .

$\Gamma(\pi^+\pi^-h_c)/\Gamma_{\text{total}}$				$\Gamma_3/\Gamma$
VALUE	DOCUMENT ID	TECN	COMMENT	
<b>seen</b>	ABLIKIM	17G	BES3	$e^+e^- \rightarrow \pi^+\pi^-h_c$

$\Gamma(\pi^+\pi^-\psi(2S))/\Gamma_{\text{total}}$				$\Gamma_4/\Gamma$
VALUE	DOCUMENT ID	TECN	COMMENT	
<b>seen</b>	<sup>1</sup> ABLIKIM	17V	BES3	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$

<sup>1</sup>From a fit to the cross section for  $e^+e^- \rightarrow \pi^+\pi^-\psi(2S) \rightarrow 2(\pi^+\pi^-)\ell^+\ell^-$  obtained from 16 center-of-mass energies between 4.008 and 4.600 GeV and comprising  $5.1\text{ fb}^{-1}$ .

### $\psi(4230)$ REFERENCES

ABLIKIM	17G	PRL 118 092002	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	17V	PR D96 032004	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	15C	PRL 114 092003	M. Ablikim <i>et al.</i>	(BES III Collab.)