

$\Sigma(1775) 5/2^-$ $I(J^P) = 1(\frac{5}{2}^-)$ Status: ****

Discovered by GALTIERI 63, this resonance plays the same role as cornerstone for isospin-1 analyses in this region as the $\Lambda(1820)F_{05}$ does in the isospin-0 channel.

For most results published before 1974 (they are now obsolete), see our 1982 edition Physics Letters **111B** 1 (1982).

$\Sigma(1775)$ POLE POSITION

REAL PART

| <u>VALUE (MeV)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|---------------------|-------------|-------------------|
| 1767_{-2}^{+2} | ¹ KAMANO | 15 | DPWA Multichannel |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| 1759 | ZHANG | 13A | DPWA Multichannel |
| ¹ From the preferred solution A in KAMANO 15. | | | |

-2×IMAGINARY PART

| <u>VALUE (MeV)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|---------------------|-------------|-------------------|
| 128_{-2}^{+4} | ¹ KAMANO | 15 | DPWA Multichannel |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| 118 | ZHANG | 13A | DPWA Multichannel |
| ¹ From the preferred solution A in KAMANO 15. | | | |

$\Sigma(1775)$ POLE RESIDUES

The normalized residue is the residue divided by $\Gamma_{pole}/2$.

Normalized residue in $N\bar{K} \rightarrow \Sigma(1775) \rightarrow N\bar{K}$

| <u>MODULUS</u> | <u>PHASE (°)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------------|---------------------|-------------|-------------------|
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| 0.371 | -32 | ¹ KAMANO | 15 | DPWA Multichannel |
| ¹ From the preferred solution A in KAMANO 15. | | | | |

Normalized residue in $N\bar{K} \rightarrow \Sigma(1775) \rightarrow \Sigma\pi$

| <u>MODULUS</u> | <u>PHASE (°)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------------|---------------------|-------------|-------------------|
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| 0.115 | -24 | ¹ KAMANO | 15 | DPWA Multichannel |
| ¹ From the preferred solution A in KAMANO 15. | | | | |

Normalized residue in $N\bar{K} \rightarrow \Sigma(1775) \rightarrow \Lambda\pi$

| <u>MODULUS</u> | <u>PHASE (°)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------------|---------------------|-------------|-------------------|
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| 0.325 | 157 | ¹ KAMANO | 15 | DPWA Multichannel |
| ¹ From the preferred solution A in KAMANO 15. | | | | |

Normalized residue in $N\bar{K} \rightarrow \Sigma(1775) \rightarrow \Sigma(1385)\pi$, D-wave

| <u>MODULUS</u> | <u>PHASE ($^\circ$)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------|------------------------------------|--------------------|-------------|----------------|
|----------------|------------------------------------|--------------------|-------------|----------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-------|-----|---------------------|----|-------------------|
| 0.391 | 137 | ¹ KAMANO | 15 | DPWA Multichannel |
|-------|-----|---------------------|----|-------------------|

¹From the preferred solution A in KAMANO 15.

Normalized residue in $N\bar{K} \rightarrow \Sigma(1775) \rightarrow \Sigma(1385)\pi$, G-wave

| <u>MODULUS</u> | <u>PHASE ($^\circ$)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------|------------------------------------|--------------------|-------------|----------------|
|----------------|------------------------------------|--------------------|-------------|----------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|--------|-----|---------------------|----|-------------------|
| 0.0129 | -58 | ¹ KAMANO | 15 | DPWA Multichannel |
|--------|-----|---------------------|----|-------------------|

¹From the preferred solution A in KAMANO 15.

 $\Sigma(1775)$ MASS

| <u>VALUE (MeV)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------|--------------------|-------------|----------------|
|--------------------|--------------------|-------------|----------------|

1770 to 1780 (≈ 1775) OUR ESTIMATE

| | | | |
|---------------|------------|-----|--|
| 1778 \pm 1 | ZHANG | 13A | DPWA Multichannel |
| 1778 \pm 5 | GOPAL | 80 | DPWA $\bar{K}N \rightarrow \bar{K}N$ |
| 1777 \pm 5 | ALSTON-... | 78 | DPWA $\bar{K}N \rightarrow \bar{K}N$ |
| 1774 \pm 5 | GOPAL | 77 | DPWA $\bar{K}N$ multichannel |
| 1775 \pm 10 | BAILLON | 75 | IPWA $\bar{K}N \rightarrow \Lambda\pi$ |
| 1774 \pm 10 | VANHORN | 75 | DPWA $K^-p \rightarrow \Lambda\pi^0$ |
| 1772 \pm 6 | KANE | 74 | DPWA $K^-p \rightarrow \Sigma\pi$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|--------------|---------------------|----|--------------------------------------|
| 1772 or 1777 | ¹ MARTIN | 77 | DPWA $\bar{K}N$ multichannel |
| 1765 | DEBELLEFON | 76 | IPWA $K^-p \rightarrow \Lambda\pi^0$ |

¹The two MARTIN 77 values are from a T-matrix pole and from a Breit-Wigner fit.

 $\Sigma(1775)$ WIDTH

| <u>VALUE (MeV)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------|--------------------|-------------|----------------|
|--------------------|--------------------|-------------|----------------|

105 to 135 (≈ 120) OUR ESTIMATE

| | | | |
|--------------|------------|-----|--|
| 131 \pm 3 | ZHANG | 13A | DPWA Multichannel |
| 137 \pm 10 | GOPAL | 80 | DPWA $\bar{K}N \rightarrow \bar{K}N$ |
| 116 \pm 10 | ALSTON-... | 78 | DPWA $\bar{K}N \rightarrow \bar{K}N$ |
| 130 \pm 10 | GOPAL | 77 | DPWA $\bar{K}N$ multichannel |
| 125 \pm 15 | BAILLON | 75 | IPWA $\bar{K}N \rightarrow \Lambda\pi$ |
| 146 \pm 18 | VANHORN | 75 | DPWA $K^-p \rightarrow \Lambda\pi^0$ |
| 154 \pm 10 | KANE | 74 | DPWA $K^-p \rightarrow \Sigma\pi$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|------------|---------------------|----|--------------------------------------|
| 102 or 103 | ¹ MARTIN | 77 | DPWA $\bar{K}N$ multichannel |
| 120 | DEBELLEFON | 76 | IPWA $K^-p \rightarrow \Lambda\pi^0$ |

¹The two MARTIN 77 values are from a T-matrix pole and from a Breit-Wigner fit.

$\Sigma(1775)$ DECAY MODES

| Mode | Fraction (Γ_i/Γ) |
|---|--------------------------------|
| Γ_1 $N\bar{K}$ | 37–43% |
| Γ_2 $\Lambda\pi$ | 14–20% |
| Γ_3 $\Sigma\pi$ | 2–5% |
| Γ_4 $\Sigma(1385)\pi$ | 8–12% |
| Γ_5 $\Sigma(1385)\pi$, <i>D</i> -wave | |
| Γ_6 $\Sigma(1385)\pi$, <i>D</i> -wave | |
| Γ_7 $\Sigma(1385)\pi$, <i>G</i> -wave | |
| Γ_8 $\Lambda(1520)\pi$, <i>P</i> -wave | 17–23% |
| Γ_9 $\Sigma\pi\pi$ | |
| Γ_{10} $\Delta(1232)\bar{K}$, <i>D</i> -wave | |
| Γ_{11} $N\bar{K}^*(892)$, <i>S</i> =1/2 | |
| Γ_{12} $N\bar{K}^*(892)$, <i>S</i> =1/2, <i>D</i> -wave | |
| Γ_{13} $N\bar{K}^*(892)$, <i>S</i> =3/2, <i>D</i> -wave | |
| Γ_{14} $N\bar{K}^*(892)$, <i>S</i> =3/2, <i>G</i> -wave | |

CONSTRAINED FIT INFORMATION

An overall fit to 7 branching ratios uses 18 measurements and one constraint to determine 5 parameters. The overall fit has a $\chi^2 = 363.4$ for 14 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

| | | | | |
|-------|-------|-------|-------|-------|
| x_2 | −44 | | | |
| x_3 | −23 | 10 | | |
| x_4 | −23 | −32 | −4 | |
| x_8 | −3 | 1 | 1 | −84 |
| | x_1 | x_2 | x_3 | x_4 |

$\Sigma(1775)$ BRANCHING RATIOS

See “Sign conventions for resonance couplings” in the Note on Λ and Σ Resonances. Also, the errors quoted do not include uncertainties due to the parametrization used in the partial-wave analyses and are thus too small.

 $\Gamma(N\bar{K})/\Gamma_{\text{total}}$ Γ_1/Γ

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------|--------------------|-------------|----------------|
|--------------|--------------------|-------------|----------------|

0.37 to 0.43 OUR ESTIMATE**0.421±0.020 OUR FIT** Error includes scale factor of 2.5.**0.398±0.009 OUR AVERAGE**

| | | | |
|---|---------------------|-----|--------------------------------------|
| 0.40 ±0.01 | ZHANG | 13A | DPWA Multichannel |
| 0.40 ±0.02 | GOPAL | 80 | DPWA $\bar{K}N \rightarrow \bar{K}N$ |
| 0.37 ±0.03 | ALSTON-... | 78 | DPWA $\bar{K}N \rightarrow \bar{K}N$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| 0.402 | ¹ KAMANO | 15 | DPWA Multichannel |
| 0.41 ±0.03 | GOPAL | 77 | DPWA See GOPAL 80 |
| 0.37 or 0.36 | ² MARTIN | 77 | DPWA $\bar{K}N$ multichannel |

¹ From the preferred solution A in KAMANO 15.² The two MARTIN 77 values are from a T-matrix pole and from a Breit-Wigner fit. **$\Gamma(\Lambda\pi)/\Gamma_{\text{total}}$ Γ_2/Γ**

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------|--------------------|-------------|----------------|
|--------------|--------------------|-------------|----------------|

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

0.244 ¹ KAMANO 15 DPWA Multichannel¹ From the preferred solution A in KAMANO 15. **$\Gamma(\Lambda\pi)/\Gamma(N\bar{K})$ Γ_2/Γ_1**

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------|--------------------|-------------|----------------|
|--------------|--------------------|-------------|----------------|

0.48±0.06 OUR FIT Error includes scale factor of 2.3.**0.33±0.05** UHLIG 67 HBC $K^- p$ 0.9 GeV/c **$\Gamma(\Sigma\pi)/\Gamma_{\text{total}}$ Γ_3/Γ**

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------|--------------------|-------------|----------------|
|--------------|--------------------|-------------|----------------|

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

0.042 ¹ KAMANO 15 DPWA Multichannel¹ From the preferred solution A in KAMANO 15. **$\Gamma(\Sigma(1385)\pi)/\Gamma(N\bar{K})$ Γ_4/Γ_1**

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------|--------------------|-------------|----------------|
|--------------|--------------------|-------------|----------------|

0.79±0.11 OUR FIT Error includes scale factor of 3.2.**0.25±0.09** UHLIG 67 HBC $K^- p$ 0.9 GeV/c **$\Gamma(\Sigma(1385)\pi, D\text{-wave})/\Gamma_{\text{total}}$ Γ_6/Γ**

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------|--------------------|-------------|----------------|
|--------------|--------------------|-------------|----------------|

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

0.309 ¹ KAMANO 15 DPWA Multichannel¹ From the preferred solution A in KAMANO 15.

$\Gamma(\Sigma(1385)\pi, G\text{-wave})/\Gamma_{\text{total}} \quad \Gamma_7/\Gamma$

VALUE DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen ¹KAMANO 15 DPWA Multichannel

¹From the preferred solution A in KAMANO 15.

$\Gamma(\Lambda(1520)\pi, P\text{-wave})/\Gamma(N\bar{K}) \quad \Gamma_8/\Gamma_1$

VALUE DOCUMENT ID TECN COMMENT

0.053^{+0.080}_{-0.035} OUR FIT Error includes scale factor of 11.8.

0.28 ± 0.05 UHLIG 67 HBC $K^- p$ 0.9 GeV/c

$\Gamma(\Sigma\pi\pi)/\Gamma_{\text{total}} \quad \Gamma_9/\Gamma$

VALUE DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.12 ¹ARMENTEROS68C HDBC $K^- N \rightarrow \Sigma\pi\pi$

¹For about 3/4 of this, the $\Sigma\pi$ system has $l = 0$ and is almost entirely $\Lambda(1520)$. For the rest, the $\Sigma\pi$ has $l = 1$, which is about what is expected from the known $\Sigma(1775) \rightarrow \Sigma(1385)\pi$ rate, as seen in $\Lambda\pi\pi$.

$\Gamma(N\bar{K}^*(892), S=1/2, D\text{-wave})/\Gamma_{\text{total}} \quad \Gamma_{12}/\Gamma$

VALUE DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen ¹KAMANO 15 DPWA Multichannel

¹From the preferred solution A in KAMANO 15.

$\Gamma(N\bar{K}^*(892), S=3/2, D\text{-wave})/\Gamma_{\text{total}} \quad \Gamma_{13}/\Gamma$

VALUE DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.003 ¹KAMANO 15 DPWA Multichannel

¹From the preferred solution A in KAMANO 15.

$\Gamma(N\bar{K}^*(892), S=3/2, G\text{-wave})/\Gamma_{\text{total}} \quad \Gamma_{14}/\Gamma$

VALUE DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen ¹KAMANO 15 DPWA Multichannel

¹From the preferred solution A in KAMANO 15.

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(1775) \rightarrow \Lambda\pi \quad (\Gamma_1\Gamma_2)^{1/2}/\Gamma$

VALUE DOCUMENT ID TECN COMMENT


0.293 ± 0.013 OUR FIT Error includes scale factor of 1.8.

0.295 ± 0.012 OUR AVERAGE Signs on measurements were ignored. Error includes scale factor of 1.4. See the ideogram below.

| | | | | |
|---|----------|-----|------|-----------------------------------|
| -0.31 ± 0.01 | ZHANG | 13A | DPWA | Multichannel |
| -0.28 ± 0.03 | GOPAL | 77 | DPWA | $\bar{K}N$ multichannel |
| -0.25 ± 0.02 | BAILLON | 75 | IPWA | $\bar{K}N \rightarrow \Lambda\pi$ |
| -0.28 ^{+0.04} _{-0.05} | VANHORN | 75 | DPWA | $K^- p \rightarrow \Lambda\pi^0$ |
| -0.259 ± 0.048 | DEVENISH | 74B | | Fixed- t dispersion rel. |

• • • We do not use the following data for averages, fits, limits, etc. • • •

WEIGHTED AVERAGE
 0.090 ± 0.011 (Error scaled by 1.6)



$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(1775) \rightarrow \Sigma\pi$

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(1775) \rightarrow \Sigma(1385)\pi$, D-wave **$(\Gamma_1 \Gamma_5)^{1/2} / \Gamma$**

VALUE DOCUMENT ID TECN COMMENT

0.155 ± 0.024 OUR AVERAGE Signs on measurements were ignored. Error includes scale factor of 3.5. See the ideogram below.


| | | | | |
|--------------------|----------------------|-----|------|-------------------------------------|
| -0.12 ± 0.01 | ZHANG | 13A | DPWA | Multichannel |
| -0.184 ± 0.011 | ¹ CAMERON | 78 | DPWA | $K^- p \rightarrow \Sigma(1385)\pi$ |
| $+0.20 \pm 0.02$ | PREVOST | 74 | DPWA | $K^- N \rightarrow \Sigma(1385)\pi$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-----------------|---------------|-----|-----------------------------------|-----------------------------------|
| 0.32 ± 0.06 | SIMS | 68 | DBC | $K^- N \rightarrow \Lambda\pi\pi$ |
| 0.24 ± 0.03 | ARMENTEROS67C | HBC | $K^- p \rightarrow \Lambda\pi\pi$ | |

¹ The CAMERON 78 upper limit on G-wave decay is 0.03.

WEIGHTED AVERAGE
 0.155 ± 0.024 (Error scaled by 3.5)



$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(1775) \rightarrow \Sigma(1385)\pi$, *D*-wave

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(1775) \rightarrow \Lambda(1520)\pi$, *P*-wave **$(\Gamma_1 \Gamma_8)^{1/2} / \Gamma$**

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|


0.10 ± 0.06 OUR FIT Error includes scale factor of 11.5.

0.20 ± 0.07 OUR AVERAGE Signs on measurements were ignored. Error includes scale factor of 10.7. See the ideogram below.

| | | | | |
|----------------|----------------------|-----|------|--|
| -0.06 ± 0.01 | ZHANG | 13A | DPWA | Multichannel |
| -0.305 ± 0.010 | ¹ CAMERON | 77 | DPWA | $K^- p \rightarrow \Lambda(1520)\pi^0$ |
| 0.31 ± 0.02 | BARLETTA | 72 | DPWA | $K^- p \rightarrow \Lambda(1520)\pi^0$ |
| 0.27 ± 0.03 | ARMENTEROS65C | HBC | HBC | $K^- p \rightarrow \Lambda(1520)\pi^0$ |

¹This rate combines *P*-wave- and *F*-wave decays. The CAMERON 77 results for the separate *P*-wave- and *F*-wave decays are -0.303 ± 0.010 and -0.037 ± 0.014 . The published signs have been changed here to be in accord with the baryon-first convention.

WEIGHTED AVERAGE
 0.20 ± 0.07 (Error scaled by 11.)



Values above of weighted average, error, and scale factor are based upon the data in this ideogram only. They are not necessarily the same as our 'best' values, obtained from a least-squares constrained fit utilizing measurements of other (related) quantities as additional information.

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(1775) \rightarrow \Lambda(1520)\pi$, *P*-wave

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(1775) \rightarrow \Delta(1232)\bar{K}$, *D*-wave $(\Gamma_1 \Gamma_{10})^{1/2} / \Gamma$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|------------------|-------------|------|-------------------|
| $+0.06 \pm 0.03$ | ZHANG | 13A | DPWA Multichannel |

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(1775) \rightarrow N\bar{K}^*(892)$, *S*=1/2 $(\Gamma_1 \Gamma_{11})^{1/2} / \Gamma$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|------------------|-------------|------|-------------------|
| $+0.04 \pm 0.01$ | ZHANG | 13A | DPWA Multichannel |

$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(1775) \rightarrow N\bar{K}^*(892)$, *S*=3/2, *D*-wave

$(\Gamma_1 \Gamma_{13})^{1/2} / \Gamma$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|------------------|-------------|------|-------------------|
| $+0.04 \pm 0.01$ | ZHANG | 13A | DPWA Multichannel |

$\Sigma(1775)$ REFERENCES

| | | | | |
|------------|-----|-------------------|---|-------------------|
| KAMANO | 15 | PR C92 025205 | H. Kamano <i>et al.</i> | (ANL, OSAK) |
| ZHANG | 13A | PR C88 035205 | H. Zhang <i>et al.</i> | (KSU) |
| PDG | 82 | PL 111B 1 | M. Roos <i>et al.</i> | (HELS, CIT, CERN) |
| GOPAL | 80 | Toronto Conf. 159 | G.P. Gopal | (RHEL) IJP |
| ALSTON-... | 78 | PR D18 182 | M. Alston-Garnjost <i>et al.</i> | (LBL, MTHO+) IJP |
| Also | | PRL 38 1007 | M. Alston-Garnjost <i>et al.</i> | (LBL, MTHO+) IJP |
| CAMERON | 78 | NP B143 189 | W. Cameron <i>et al.</i> | (RHEL, LOIC) IJP |
| CAMERON | 77 | NP B131 399 | W. Cameron <i>et al.</i> | (RHEL, LOIC) IJP |
| GOPAL | 77 | NP B119 362 | G.P. Gopal <i>et al.</i> | (LOIC, RHEL) IJP |
| MARTIN | 77 | NP B127 349 | B.R. Martin, M.K. Pidcock, R.G. Moorhouse | (LOUC+) IJP |
| Also | | NP B126 266 | B.R. Martin, M.K. Pidcock | (LOUC) |
| Also | | NP B126 285 | B.R. Martin, M.K. Pidcock | (LOUC) IJP |
| DEBELLEFON | 76 | NP B109 129 | A. de Bellefon, A. Berthon | (CDEF) IJP |

| | | | | |
|------------|-----|--------------|---|------------------------|
| BAILLON | 75 | NP B94 39 | P.H. Baillon, P.J. Litchfield | (CERN, RHEL) IJP |
| VANHORN | 75 | NP B87 145 | A.J. van Horn | (LBL) IJP |
| Also | | NP B87 157 | A.J. van Horn | (LBL) IJP |
| DEVENISH | 74B | NP B81 330 | R.C.E. Devenish, C.D. Froggatt, B.R. Martin | (DESY+) |
| KANE | 74 | LBL-2452 | D.F. Kane | (LBL) IJP |
| PREVOST | 74 | NP B69 246 | J. Prevost <i>et al.</i> | (SACL, CERN, HEID) |
| BARLETTA | 72 | NP B40 45 | W.A. Barletta | (EFI) IJP |
| Also | | PRL 17 841 | S. Fenster <i>et al.</i> | (CHIC, ANL, CERN) IJP |
| ARMENTEROS | 68C | NP B8 216 | R. Armenteros <i>et al.</i> | (CERN, HEID, SACL) I |
| SIMS | 68 | PRL 21 1413 | W.H. Sims <i>et al.</i> | (FSU, TUFTS, BRAN) |
| ARMENTEROS | 67C | ZPHY 202 486 | R. Armenteros <i>et al.</i> | (CERN, HEID, SACL) |
| UHLIG | 67 | PR 155 1448 | R.P. Uhlig <i>et al.</i> | (UMD, NRL) |
| ARMENTEROS | 65C | PL 19 338 | R. Armenteros <i>et al.</i> | (CERN, HEID, SACL) IJP |
| GALTIERI | 63 | PL 6 296 | A. Galtieri, A. Hussain, R. Tripp | (LRL) IJ |
