

$$I(J^P) = \frac{1}{2}(0^-)$$

## $D^\pm$ MASS

The fit includes  $D^\pm$ ,  $D^0$ ,  $D_s^\pm$ ,  $D^{*\pm}$ ,  $D^{*0}$ ,  $D_s^{*\pm}$ ,  $D_1(2420)^0$ ,  $D_2^*(2460)^0$ , and  $D_{s1}(2536)^\pm$  mass and mass difference measurements.

| VALUE (MeV)   | EVTS     | DOCUMENT ID            | TECN     | COMMENT                      |
|---|----------|------------------------|----------|------------------------------|
| <b>1869.65 ± 0.05 OUR FIT</b>   |          |                        |          |                              |
| <b>1869.5 ± 0.4 OUR AVERAGE</b>   |          |                        |          |                              |
| 1869.53 ± 0.49 ± 0.20   | 110 ± 15 | ANASHIN                | 10A KEDR | $e^+e^-$ at $\psi(3770)$     |
| 1870.0 ± 0.5 ± 1.0  | 317      | BARLAG                 | 90C ACCM | $\pi^-$ Cu 230 GeV           |
| 1869.4 ± 0.6  |          | <sup>1</sup> TRILLING  | 81 RVUE  | $e^+e^-$ 3.77 GeV            |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● |          |                        |          |                              |
| 1875 ± 10   | 9        | ADAMOVICH              | 87 EMUL  | Photoproduction              |
| 1860 ± 16   | 6        | ADAMOVICH              | 84 EMUL  | Photoproduction              |
| 1863 ± 4  |          | DERRICK                | 84 HRS   | $e^+e^-$ 29 GeV              |
| 1868.4 ± 0.5  |          | <sup>1</sup> SCHINDLER | 81 MRK2  | $e^+e^-$ 3.77 GeV            |
| 1874 ± 5  |          | GOLDHABER              | 77 MRK1  | $D^0$ , $D^+$ recoil spectra |
| 1868.3 ± 0.9  |          | <sup>1</sup> PERUZZI   | 77 LGW   | $e^+e^-$ 3.77 GeV            |
| 1874 ± 11   |          | PICCOLO                | 77 MRK1  | $e^+e^-$ 4.03, 4.41 GeV      |
| 1876 ± 15   | 50       | PERUZZI                | 76 MRK1  | $K^\mp \pi^\pm \pi^\pm$      |

<sup>1</sup>PERUZZI 77 and SCHINDLER 81 errors do not include the 0.13% uncertainty in the absolute SPEAR energy calibration. TRILLING 81 uses the high precision  $J/\psi(1S)$  and  $\psi(2S)$  measurements of ZHOLENTZ 80 to determine this uncertainty and combines the PERUZZI 77 and SCHINDLER 81 results to obtain the value quoted.

## $D^\pm$ MEAN LIFE

Measurements with an error  $> 100 \times 10^{-15}$  s have been omitted from the Listings.

| VALUE ( $10^{-15}$ s)   | EVTS | DOCUMENT ID         | TECN     | COMMENT  |
|---|------|---------------------|----------|--|
| <b>1040 ± 7 OUR AVERAGE</b>   |      |                     |          |  |
| 1039.4 ± 4.3 ± 7.0  | 110k | LINK                | 02F FOCS | $\gamma$ nucleus, $\approx 180$ GeV            |
| 1033.6 ± 22.1 <sup>+9.9</sup> <sub>-12.7</sub>                                | 3.7k | BONVICINI           | 99 CLEO  | $e^+e^- \approx \Upsilon(4S)$                  |
| 1048 ± 15 ± 11  | 9k   | FRABETTI            | 94D E687 | $D^+ \rightarrow K^- \pi^+ \pi^+$              |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● |      |                     |          |  |
| 1075 ± 40 ± 18  | 2.4k | FRABETTI            | 91 E687  | $\gamma$ Be, $D^+ \rightarrow K^- \pi^+ \pi^+$ |
| 1030 ± 80 ± 60  | 200  | ALVAREZ             | 90 NA14  | $\gamma$ , $D^+ \rightarrow K^- \pi^+ \pi^+$   |
| 1050 <sup>+77</sup> <sub>-72</sub>  | 317  | <sup>1</sup> BARLAG | 90C ACCM | $\pi^-$ Cu 230 GeV                             |
| 1050 ± 80 ± 70  | 363  | ALBRECHT            | 88i ARG  | $e^+e^-$ 10 GeV                                |
| 1090 ± 30 ± 25  | 2.9k | RAAB                | 88 E691  | Photoproduction                                |

<sup>1</sup>BARLAG 90C estimates the systematic error to be negligible.

## $D^+$ DECAY MODES

Most decay modes (other than the semileptonic modes) that involve a neutral  $K$  meson are now given as  $K_S^0$  modes, not as  $\bar{K}^0$  modes. Nearly always it is a  $K_S^0$  that is measured, and interference between Cabibbo-allowed and doubly Cabibbo-suppressed modes can invalidate the assumption that  $2\Gamma(K_S^0) = \Gamma(\bar{K}^0)$ .

| Mode   | Fraction ( $\Gamma_i/\Gamma$ )   | Scale factor/<br>Confidence level |
|--|----------------------------------|-----------------------------------|
| <b>Inclusive modes</b>   |                                  |                                   |
| $\Gamma_1$ $e^+$ semileptonic  | $(16.07 \pm 0.30)$ %             |                                   |
| $\Gamma_2$ $\mu^+$ anything  | $(17.6 \pm 3.2)$ %               |                                   |
| $\Gamma_3$ $K^-$ anything  | $(25.7 \pm 1.4)$ %               |                                   |
| $\Gamma_4$ $\bar{K}^0$ anything + $K^0$ anything   | $(61 \pm 5)$ %                   |                                   |
| $\Gamma_5$ $K^+$ anything  | $(5.9 \pm 0.8)$ %                |                                   |
| $\Gamma_6$ $K^*(892)^-$ anything   | $(6 \pm 5)$ %                    |                                   |
| $\Gamma_7$ $\bar{K}^*(892)^0$ anything   | $(23 \pm 5)$ %                   |                                   |
| $\Gamma_8$ $K^*(892)^0$ anything   | $< 6.6$ %                        | CL=90%                            |
| $\Gamma_9$ $\eta$ anything   | $(6.3 \pm 0.7)$ %                |                                   |
| $\Gamma_{10}$ $\eta'$ anything   | $(1.04 \pm 0.18)$ %              |                                   |
| $\Gamma_{11}$ $\phi$ anything  | $(1.03 \pm 0.12)$ %              |                                   |
| <b>Leptonic and semileptonic modes</b>   |                                  |                                   |
| $\Gamma_{12}$ $e^+ \nu_e$  | $< 8.8$ $\times 10^{-6}$         | CL=90%                            |
| $\Gamma_{13}$ $\gamma e^+ \nu_e$   | $< 3.0$ $\times 10^{-5}$         | CL=90%                            |
| $\Gamma_{14}$ $\mu^+ \nu_\mu$  | $(3.74 \pm 0.17) \times 10^{-4}$ |                                   |
| $\Gamma_{15}$ $\tau^+ \nu_\tau$  | $< 1.2$ $\times 10^{-3}$         | CL=90%                            |
| $\Gamma_{16}$ $\bar{K}^0 e^+ \nu_e$  | $(8.73 \pm 0.10)$ %              |                                   |
| $\Gamma_{17}$ $\bar{K}^0 \mu^+ \nu_\mu$  | $(8.76 \pm 0.19)$ %              |                                   |
| $\Gamma_{18}$ $K^- \pi^+ e^+ \nu_e$  | $(4.02 \pm 0.18)$ %              | S=3.2                             |
| $\Gamma_{19}$ $\bar{K}^*(892)^0 e^+ \nu_e, \bar{K}^*(892)^0 \rightarrow K^- \pi^+$       | $(3.77 \pm 0.17)$ %              |                                   |
| $\Gamma_{20}$ $(K^- \pi^+)_{[0.8-1.0]\text{GeV}} e^+ \nu_e$                              | $(3.39 \pm 0.09)$ %              |                                   |
| $\Gamma_{21}$ $(K^- \pi^+)_{S\text{-wave}} e^+ \nu_e$                                    | $(2.28 \pm 0.11) \times 10^{-3}$ |                                   |
| $\Gamma_{22}$ $\bar{K}^*(1410)^0 e^+ \nu_e, \bar{K}^*(1410)^0 \rightarrow K^- \pi^+$     | $< 6$ $\times 10^{-3}$           | CL=90%                            |
| $\Gamma_{23}$ $\bar{K}_2^*(1430)^0 e^+ \nu_e, \bar{K}_2^*(1430)^0 \rightarrow K^- \pi^+$ | $< 5$ $\times 10^{-4}$           | CL=90%                            |
| $\Gamma_{24}$ $K^- \pi^+ e^+ \nu_e$ nonresonant  | $< 7$ $\times 10^{-3}$           | CL=90%                            |
| $\Gamma_{25}$ $\bar{K}^*(892)^0 e^+ \nu_e$   | $(5.40 \pm 0.10)$ %              | S=1.1                             |
| $\Gamma_{26}$ $K^- \pi^+ \mu^+ \nu_\mu$  | $(3.65 \pm 0.34)$ %              |                                   |
| $\Gamma_{27}$ $\bar{K}^*(892)^0 \mu^+ \nu_\mu, \bar{K}^*(892)^0 \rightarrow K^- \pi^+$   | $(3.52 \pm 0.10)$ %              |                                   |

|               |   |   |        |
|---------------|---|---|--------|
| $\Gamma_{28}$ | $K^- \pi^+ \mu^+ \nu_\mu$ nonresonant                 | $(1.9 \pm 0.5) \times 10^{-3}$              |        |
| $\Gamma_{29}$ | $\bar{K}^*(892)^0 \mu^+ \nu_\mu$                      | $(5.27 \pm 0.15) \%$                        |        |
| $\Gamma_{30}$ | $K^- \pi^+ \pi^0 \mu^+ \nu_\mu$                       | $< 1.5 \times 10^{-3}$                      | CL=90% |
| $\Gamma_{31}$ | $\bar{K}_0^*(1430)^0 \mu^+ \nu_\mu$                   | $< 2.3 \times 10^{-4}$                      | CL=90% |
| $\Gamma_{32}$ | $\bar{K}^*(1680)^0 \mu^+ \nu_\mu$                     | $< 1.5 \times 10^{-3}$                      | CL=90% |
| $\Gamma_{33}$ | $\pi^0 e^+ \nu_e$                                     | $(3.72 \pm 0.17) \times 10^{-3}$            | S=2.0  |
| $\Gamma_{34}$ | $\pi^0 \mu^+ \nu_\mu$                                 | $(3.50 \pm 0.15) \times 10^{-3}$            |        |
| $\Gamma_{35}$ | $\eta e^+ \nu_e$                                      | $(1.11 \pm 0.07) \times 10^{-3}$            |        |
| $\Gamma_{36}$ | $\rho^0 e^+ \nu_e$                                    | $(2.18 \pm_{-0.25}^{+0.17}) \times 10^{-3}$ |        |
| $\Gamma_{37}$ | $\rho^0 \mu^+ \nu_\mu$                                | $(2.4 \pm 0.4) \times 10^{-3}$              |        |
| $\Gamma_{38}$ | $\omega e^+ \nu_e$                                    | $(1.69 \pm 0.11) \times 10^{-3}$            |        |
| $\Gamma_{39}$ | $\eta'(958) e^+ \nu_e$                                | $(2.0 \pm 0.4) \times 10^{-4}$              |        |
| $\Gamma_{40}$ | $a(980)^0 e^+ \nu_e, a(980)^0 \rightarrow \eta \pi^0$ | $(1.7 \pm_{-0.7}^{+0.8}) \times 10^{-4}$    |        |
| $\Gamma_{41}$ | $\phi e^+ \nu_e$                                      | $< 1.3 \times 10^{-5}$                      | CL=90% |
| $\Gamma_{42}$ | $D^0 e^+ \nu_e$                                       | $< 1.0 \times 10^{-4}$                      | CL=90% |

### Hadronic modes with a $\bar{K}$ or $\bar{K}K\bar{K}$

|               |   |  |       |
|---------------|---|--|-------|
| $\Gamma_{43}$ | $K_S^0 \pi^+$   | $(1.562 \pm 0.031) \%$                   | S=1.7 |
| $\Gamma_{44}$ | $K_L^0 \pi^+$   | $(1.46 \pm 0.05) \%$                     |       |
| $\Gamma_{45}$ | $K^- 2\pi^+$  | [a] $(9.38 \pm 0.16) \%$                 | S=1.6 |
| $\Gamma_{46}$ | $(K^- \pi^+)_{S\text{-wave}} \pi^+$   | $(7.52 \pm 0.17) \%$                     |       |
| $\Gamma_{47}$ | $\bar{K}_0^*(700)^0 \pi^+, \bar{K}_0^*(700) \rightarrow$                    |  |       |
| $\Gamma_{48}$ | $\bar{K}_0^*(1430)^0 \pi^+,$<br>$\bar{K}_0^*(1430)^0 \rightarrow K^- \pi^+$ | [b] $(1.25 \pm 0.06) \%$                 |       |
| $\Gamma_{49}$ | $\bar{K}^*(892)^0 \pi^+,$<br>$\bar{K}^*(892)^0 \rightarrow K^- \pi^+$       | $(1.04 \pm 0.12) \%$                     |       |
| $\Gamma_{50}$ | $\bar{K}^*(1410)^0 \pi^+, \bar{K}^{*0} \rightarrow$                         | not seen                                 |       |
| $\Gamma_{51}$ | $\bar{K}_2^*(1430)^0 \pi^+,$<br>$\bar{K}_2^*(1430)^0 \rightarrow K^- \pi^+$ | [b] $(2.3 \pm 0.7) \times 10^{-4}$       |       |
| $\Gamma_{52}$ | $\bar{K}^*(1680)^0 \pi^+,$<br>$\bar{K}^*(1680)^0 \rightarrow K^- \pi^+$     | [b] $(2.2 \pm 1.1) \times 10^{-4}$       |       |
| $\Gamma_{53}$ | $K^- (2\pi^+)_{I=2}$  | $(1.45 \pm 0.26) \%$                     |       |
| $\Gamma_{54}$ | $K^- 2\pi^+$ nonresonant  |  |       |
| $\Gamma_{55}$ | $K_S^0 \pi^+ \pi^0$   | [a] $(7.36 \pm 0.21) \%$                 |       |
| $\Gamma_{56}$ | $K_S^0 \rho^+$  | $(6.14 \pm_{-0.35}^{+0.60}) \%$          |       |
| $\Gamma_{57}$ | $K_S^0 \rho(1450)^+, \rho^+ \rightarrow \pi^+ \pi^0$                        | $(1.5 \pm_{-1.4}^{+1.2}) \times 10^{-3}$ |       |
| $\Gamma_{58}$ | $\bar{K}^*(892)^0 \pi^+,$<br>$\bar{K}^*(892)^0 \rightarrow K_S^0 \pi^0$     | $(2.64 \pm 0.32) \times 10^{-3}$         |       |

|                 |  |   |       |
|-----------------|--|---|-------|
| Γ <sub>59</sub> | $\bar{K}_0^*(1430)^0 \pi^+, \bar{K}_0^{*0} \rightarrow K_S^0 \pi^0$            | ( 2.7 ± 0.9 ) × 10 <sup>-3</sup>                          |       |
| Γ <sub>60</sub> | $\bar{K}_0^*(1680)^0 \pi^+, \bar{K}_0^{*0} \rightarrow K_S^0 \pi^0$            | ( 10 + <sup>7</sup> / <sub>-10</sub> ) × 10 <sup>-4</sup> |       |
| Γ <sub>61</sub> | $\bar{\kappa}^0 \pi^+, \bar{\kappa}^0 \rightarrow K_S^0 \pi^0$                 | ( 6 + <sup>5</sup> / <sub>-4</sub> ) × 10 <sup>-3</sup>   |       |
| Γ <sub>62</sub> | $K_S^0 \pi^+ \pi^0$ nonresonant  | ( 3 ± 4 ) × 10 <sup>-3</sup>                              |       |
| Γ <sub>63</sub> | $K_S^0 \pi^+ \pi^0$ nonresonant and $\bar{\kappa}^0 \pi^+$                     | ( 1.37 + <sup>0.21</sup> / <sub>-0.40</sub> ) %           |       |
| Γ <sub>64</sub> | $(K_S^0 \pi^0)_{S\text{-wave}} \pi^+$  | ( 1.27 + <sup>0.27</sup> / <sub>-0.33</sub> ) %           |       |
| Γ <sub>65</sub> | $K_S^0 \pi^+ \eta'(958)$   | ( 1.90 ± 0.21 ) × 10 <sup>-3</sup>                        |       |
| Γ <sub>66</sub> | $K^- 2\pi^+ \pi^0$   | [c] ( 6.25 ± 0.18 ) %                                     |       |
| Γ <sub>67</sub> | $K_S^0 2\pi^+ \pi^-$   | [c] ( 3.10 ± 0.09 ) %                                     |       |
| Γ <sub>68</sub> | $K^- 3\pi^+ \pi^-$   | [a] ( 5.7 ± 0.5 ) × 10 <sup>-3</sup>                      | S=1.1 |
| Γ <sub>69</sub> | $\bar{K}^*(892)^0 2\pi^+ \pi^-, \bar{K}^*(892)^0 \rightarrow K^- \pi^+$        | ( 1.2 ± 0.4 ) × 10 <sup>-3</sup>                          |       |
| Γ <sub>70</sub> | $\bar{K}^*(892)^0 \rho^0 \pi^+, \bar{K}^*(892)^0 \rightarrow K^- \pi^+$        | ( 2.3 ± 0.4 ) × 10 <sup>-3</sup>                          |       |
| Γ <sub>71</sub> | $\bar{K}^*(892)^0 a_1(1260)^+$   | [d] ( 9.3 ± 1.9 ) × 10 <sup>-3</sup>                      |       |
| Γ <sub>72</sub> | $\bar{K}^*(892)^0 2\pi^+ \pi^-$ no-ρ, $\bar{K}^*(892)^0 \rightarrow K^- \pi^+$ |   |       |
| Γ <sub>73</sub> | $K^- \rho^0 2\pi^+$  | ( 1.72 ± 0.28 ) × 10 <sup>-3</sup>                        |       |
| Γ <sub>74</sub> | $K^- 3\pi^+ \pi^-$ nonresonant   | ( 4.0 ± 2.9 ) × 10 <sup>-4</sup>                          |       |
| Γ <sub>75</sub> | $K^+ 2K_S^0$   | ( 2.54 ± 0.13 ) × 10 <sup>-3</sup>                        |       |
| Γ <sub>76</sub> | $K^+ K^- K_S^0 \pi^+$  | ( 2.4 ± 0.5 ) × 10 <sup>-4</sup>                          |       |

### Pionic modes

|                 |  |                                      |        |
|-----------------|--|--------------------------------------|--------|
| Γ <sub>77</sub> | $\pi^+ \pi^0$  | ( 1.247 ± 0.033 ) × 10 <sup>-3</sup> |        |
| Γ <sub>78</sub> | $2\pi^+ \pi^-$   | ( 3.27 ± 0.18 ) × 10 <sup>-3</sup>   |        |
| Γ <sub>79</sub> | $\rho^0 \pi^+$   | ( 8.3 ± 1.5 ) × 10 <sup>-4</sup>     |        |
| Γ <sub>80</sub> | $\pi^+ (\pi^+ \pi^-)_{S\text{-wave}}$                      | ( 1.83 ± 0.16 ) × 10 <sup>-3</sup>   |        |
| Γ <sub>81</sub> | $\sigma \pi^+, \sigma \rightarrow \pi^+ \pi^-$             | ( 1.38 ± 0.12 ) × 10 <sup>-3</sup>   |        |
| Γ <sub>82</sub> | $f_0(980) \pi^+, f_0(980) \rightarrow \pi^+ \pi^-$         | ( 1.56 ± 0.33 ) × 10 <sup>-4</sup>   |        |
| Γ <sub>83</sub> | $f_0(1370) \pi^+, f_0(1370) \rightarrow \pi^+ \pi^-$       | ( 8 ± 4 ) × 10 <sup>-5</sup>         |        |
| Γ <sub>84</sub> | $f_2(1270) \pi^+, f_2(1270) \rightarrow \pi^+ \pi^-$       | ( 5.0 ± 0.9 ) × 10 <sup>-4</sup>     |        |
| Γ <sub>85</sub> | $\rho(1450)^0 \pi^+, \rho(1450)^0 \rightarrow \pi^+ \pi^-$ | < 8 × 10 <sup>-5</sup>               | CL=95% |
| Γ <sub>86</sub> | $f_0(1500) \pi^+, f_0(1500) \rightarrow \pi^+ \pi^-$       | ( 1.1 ± 0.4 ) × 10 <sup>-4</sup>     |        |

|               |  |                   |                  |        |
|---------------|--|-------------------|------------------|--------|
| $\Gamma_{87}$ | $f_0(1710)\pi^+$ ,<br>$f_0(1710) \rightarrow \pi^+\pi^-$ | $< 5$             | $\times 10^{-5}$ | CL=95% |
| $\Gamma_{88}$ | $f_0(1790)\pi^+$ ,<br>$f_0(1790) \rightarrow \pi^+\pi^-$ | $< 7$             | $\times 10^{-5}$ | CL=95% |
| $\Gamma_{89}$ | $(\pi^+\pi^+)_{S\text{-wave}}\pi^-$                      | $< 1.2$           | $\times 10^{-4}$ | CL=95% |
| $\Gamma_{90}$ | $2\pi^+\pi^-$ nonresonant                                | $< 1.1$           | $\times 10^{-4}$ | CL=95% |
| $\Gamma_{91}$ | $\pi^+2\pi^0$  | $(4.7 \pm 0.4)$   | $\times 10^{-3}$ |        |
| $\Gamma_{92}$ | $2\pi^+\pi^-\pi^0$                                       | $(1.16 \pm 0.08)$ | %                |        |
| $\Gamma_{93}$ | $3\pi^+2\pi^-$   | $(1.66 \pm 0.16)$ | $\times 10^{-3}$ | S=1.1  |
| $\Gamma_{94}$ | $\eta\pi^+$  | $(3.77 \pm 0.09)$ | $\times 10^{-3}$ |        |
| $\Gamma_{95}$ | $\eta\pi^+\pi^0$   | $(1.38 \pm 0.35)$ | $\times 10^{-3}$ |        |
| $\Gamma_{96}$ | $\omega\pi^+$  | $(2.8 \pm 0.6)$   | $\times 10^{-4}$ |        |
| $\Gamma_{97}$ | $\eta'(958)\pi^+$  | $(4.97 \pm 0.19)$ | $\times 10^{-3}$ |        |
| $\Gamma_{98}$ | $\eta'(958)\pi^+\pi^0$                                   | $(1.6 \pm 0.5)$   | $\times 10^{-3}$ |        |

**Hadronic modes with a  $K\bar{K}$  pair**

|                |  |                              |                  |       |
|----------------|--|------------------------------|------------------|-------|
| $\Gamma_{99}$  | $K^+K_S^0$   | $(3.04 \pm 0.09)$            | $\times 10^{-3}$ | S=2.2 |
| $\Gamma_{100}$ | $K^+K^-\pi^+$  | [a] $(9.93 \pm 0.24)$        | $\times 10^{-3}$ |       |
| $\Gamma_{101}$ | $\phi\pi^+$ , $\phi \rightarrow K^+K^-$                                    | $(2.76 \pm_{-0.09}^{+0.08})$ | $\times 10^{-3}$ |       |
| $\Gamma_{102}$ | $K^+\bar{K}^*(892)^0$ ,<br>$\bar{K}^*(892)^0 \rightarrow K^-\pi^+$         | $(2.55 \pm_{-0.14}^{+0.09})$ | $\times 10^{-3}$ |       |
| $\Gamma_{103}$ | $K^+\bar{K}_0^*(1430)^0$ , $\bar{K}_0^*(1430)^0 \rightarrow$<br>$K^-\pi^+$ | $(1.9 \pm 0.4)$              | $\times 10^{-3}$ |       |
| $\Gamma_{104}$ | $K^+\bar{K}_2^*(1430)^0$ , $\bar{K}_2^* \rightarrow$<br>$K^-\pi^+$         | $(1.7 \pm_{-0.8}^{+1.3})$    | $\times 10^{-4}$ |       |
| $\Gamma_{105}$ | $K^+\bar{K}_0^*(700)$ , $\bar{K}_0^* \rightarrow K^-\pi^+$                 | $(6.9 \pm_{-2.1}^{+4.0})$    | $\times 10^{-4}$ |       |
| $\Gamma_{106}$ | $a_0(1450)^0\pi^+$ , $a_0^0 \rightarrow K^+K^-$                            | $(4.6 \pm_{-1.9}^{+7.0})$    | $\times 10^{-4}$ |       |
| $\Gamma_{107}$ | $\phi(1680)\pi^+$ , $\phi \rightarrow K^+K^-$                              | $(5.1 \pm_{-1.9}^{+4.0})$    | $\times 10^{-5}$ |       |
| $\Gamma_{108}$ | $K_S^0K_S^0\pi^+$  | $(2.70 \pm 0.13)$            | $\times 10^{-3}$ |       |
| $\Gamma_{109}$ | $K^+K_S^0\pi^+\pi^-$   | $(1.74 \pm 0.18)$            | $\times 10^{-3}$ |       |
| $\Gamma_{110}$ | $K_S^0K^-2\pi^+$   | $(2.38 \pm 0.17)$            | $\times 10^{-3}$ |       |
| $\Gamma_{111}$ | $K^+K^-2\pi^+\pi^-$  | $(2.3 \pm 1.2)$              | $\times 10^{-4}$ |       |

A few poorly measured branching fractions:

|                |                                |                           |   |        |
|----------------|--------------------------------|---------------------------|---|--------|
| $\Gamma_{112}$ | $\phi\pi^+\pi^0$               | $(2.3 \pm 1.0)$           | % |        |
| $\Gamma_{113}$ | $\phi\rho^+$                   | $< 1.5$                   | % | CL=90% |
| $\Gamma_{114}$ | $K^+K^-\pi^+\pi^0$ non- $\phi$ | $(1.5 \pm_{-0.6}^{+0.7})$ | % |        |
| $\Gamma_{115}$ | $K^*(892)^+K_S^0$              | $(1.7 \pm 0.8)$           | % |        |

### Doubly Cabibbo-suppressed modes

|                |  |                                    |       |
|----------------|--|------------------------------------|-------|
| $\Gamma_{116}$ | $K^+ \pi^0$  | $( 2.08 \pm 0.21 ) \times 10^{-4}$ | S=1.4 |
| $\Gamma_{117}$ | $K^+ \eta$   | $( 1.25 \pm 0.16 ) \times 10^{-4}$ | S=1.1 |
| $\Gamma_{118}$ | $K^+ \eta'(958)$   | $( 1.85 \pm 0.20 ) \times 10^{-4}$ |       |
| $\Gamma_{119}$ | $K^+ \pi^+ \pi^-$  | $( 5.42 \pm 0.22 ) \times 10^{-4}$ |       |
| $\Gamma_{120}$ | $K^+ \rho^0$   | $( 2.1 \pm 0.5 ) \times 10^{-4}$   |       |
| $\Gamma_{121}$ | $K^*(892)^0 \pi^+, K^*(892)^0 \rightarrow K^+ \pi^-$       | $( 2.5 \pm 0.4 ) \times 10^{-4}$   |       |
| $\Gamma_{122}$ | $K^+ f_0(980), f_0(980) \rightarrow \pi^+ \pi^-$           | $( 4.8 \pm 2.9 ) \times 10^{-5}$   |       |
| $\Gamma_{123}$ | $K_2^*(1430)^0 \pi^+, K_2^*(1430)^0 \rightarrow K^+ \pi^-$ | $( 4.3 \pm 2.9 ) \times 10^{-5}$   |       |
| $\Gamma_{124}$ | $K^+ \pi^+ \pi^-$ nonresonant                              | not seen                           |       |
| $\Gamma_{125}$ | $2K^+ K^-$   | $( 8.9 \pm 2.1 ) \times 10^{-5}$   |       |

### $\Delta C = 1$ weak neutral current (*C1*) modes, or Lepton Family number (*LF*) or Lepton number (*L*) violating modes

|                |  |           |                       |                  |        |
|----------------|--|-----------|-----------------------|------------------|--------|
| $\Gamma_{126}$ | $\pi^+ e^+ e^-$                            | <i>C1</i> | $< 1.1$               | $\times 10^{-6}$ | CL=90% |
| $\Gamma_{127}$ | $\pi^+ \pi^0 e^+ e^-$                      |           | $< 1.4$               | $\times 10^{-5}$ | CL=90% |
| $\Gamma_{128}$ | $\pi^+ \phi, \phi \rightarrow e^+ e^-$     | [e]       | $( 1.7 \pm 0.9 )$     | $\times 10^{-6}$ |        |
| $\Gamma_{129}$ | $\pi^+ \mu^+ \mu^-$                        | <i>C1</i> | $< 7.3$               | $\times 10^{-8}$ | CL=90% |
| $\Gamma_{130}$ | $\pi^+ \phi, \phi \rightarrow \mu^+ \mu^-$ | [e]       | $( 1.8 \pm 0.8 )$     | $\times 10^{-6}$ |        |
| $\Gamma_{131}$ | $\rho^+ \mu^+ \mu^-$                       | <i>C1</i> | $< 5.6$               | $\times 10^{-4}$ | CL=90% |
| $\Gamma_{132}$ | $K^+ e^+ e^-$                              | [f]       | $< 1.0$               | $\times 10^{-6}$ | CL=90% |
| $\Gamma_{133}$ | $K^+ \pi^0 e^+ e^-$                        |           | $< 1.5$               | $\times 10^{-5}$ | CL=90% |
| $\Gamma_{134}$ | $K_S^0 \pi^+ e^+ e^-$                      |           | $< 2.6$               | $\times 10^{-5}$ | CL=90% |
| $\Gamma_{135}$ | $K_S^0 K^+ e^+ e^-$                        |           | $< 1.1$               | $\times 10^{-5}$ | CL=90% |
| $\Gamma_{136}$ | $K^+ \mu^+ \mu^-$                          | [f]       | $< 4.3$               | $\times 10^{-6}$ | CL=90% |
| $\Gamma_{137}$ | $\pi^+ e^+ \mu^-$                          | <i>LF</i> | $< 2.9$               | $\times 10^{-6}$ | CL=90% |
| $\Gamma_{138}$ | $\pi^+ e^- \mu^+$                          | <i>LF</i> | $< 3.6$               | $\times 10^{-6}$ | CL=90% |
| $\Gamma_{139}$ | $K^+ e^+ \mu^-$                            | <i>LF</i> | $< 1.2$               | $\times 10^{-6}$ | CL=90% |
| $\Gamma_{140}$ | $K^+ e^- \mu^+$                            | <i>LF</i> | $< 2.8$               | $\times 10^{-6}$ | CL=90% |
| $\Gamma_{141}$ | $\pi^- 2e^+$                               | <i>L</i>  | $< 1.1$               | $\times 10^{-6}$ | CL=90% |
| $\Gamma_{142}$ | $\pi^- 2\mu^+$                             | <i>L</i>  | $< 2.2$               | $\times 10^{-8}$ | CL=90% |
| $\Gamma_{143}$ | $\pi^- e^+ \mu^+$                          | <i>L</i>  | $< 2.0$               | $\times 10^{-6}$ | CL=90% |
| $\Gamma_{144}$ | $\rho^- 2\mu^+$                            | <i>L</i>  | $< 5.6$               | $\times 10^{-4}$ | CL=90% |
| $\Gamma_{145}$ | $K^- 2e^+$                                 | <i>L</i>  | $< 9$                 | $\times 10^{-7}$ | CL=90% |
| $\Gamma_{146}$ | $K^- 2\mu^+$                               | <i>L</i>  | $< 1.0$               | $\times 10^{-5}$ | CL=90% |
| $\Gamma_{147}$ | $K^- e^+ \mu^+$                            | <i>L</i>  | $< 1.9$               | $\times 10^{-6}$ | CL=90% |
| $\Gamma_{148}$ | $K^*(892)^- 2\mu^+$                        | <i>L</i>  | $< 8.5$               | $\times 10^{-4}$ | CL=90% |
| $\Gamma_{149}$ | Unaccounted decay modes                    |           | $( 63.3 \pm 0.4 ) \%$ |                  | S=1.3  |

- [a] The branching fraction for this mode may differ from the sum of the submodes that contribute to it, due to interference effects. See the relevant papers.
- [b] These subfractions of the  $K^- 2\pi^+$  mode are uncertain: see the Particle Listings.
- [c] Submodes of the  $D^+ \rightarrow K^- 2\pi^+ \pi^0$  and  $K_S^0 2\pi^+ \pi^-$  modes were studied by ANJOS 92C and COFFMAN 92B, but with at most 142 events for the first mode and 229 for the second – not enough for precise results. With nothing new for 18 years, we refer to our 2008 edition, Physics Letters **B667** 1 (2008), for those results.
- [d] The unseen decay modes of the resonances are included.
- [e] This is *not* a test for the  $\Delta C=1$  weak neutral current, but leads to the  $\pi^+ \ell^+ \ell^-$  final state.
- [f] This mode is not a useful test for a  $\Delta C=1$  weak neutral current because both quarks must change flavor in this decay.

---

### CONSTRAINED FIT INFORMATION

An overall fit to 31 branching ratios uses 41 measurements and one constraint to determine 17 parameters. The overall fit has a  $\chi^2 = 62.8$  for 25 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$ , in percent, from the fit to the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ . The fit constrains the  $x_i$  whose labels appear in this array to sum to one.

|           |          |          |          |          |          |          |          |          |          |          |
|-----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|
| $x_{18}$  | 0        |          |          |          |          |          |          |          |          |          |
| $x_{25}$  | 0        | 0        |          |          |          |          |          |          |          |          |
| $x_{29}$  | 8        | 0        | 0        |          |          |          |          |          |          |          |
| $x_{36}$  | 0        | 0        | 0        | 0        |          |          |          |          |          |          |
| $x_{43}$  | 0        | 5        | 0        | 0        | 0        |          |          |          |          |          |
| $x_{45}$  | 0        | 28       | 0        | 0        | 0        | 19       |          |          |          |          |
| $x_{68}$  | 0        | 5        | 0        | 0        | 0        | 4        | 19       |          |          |          |
| $x_{77}$  | 0        | 6        | 0        | 0        | 0        | 4        | 22       | 4        |          |          |
| $x_{93}$  | 0        | 5        | 0        | 0        | 0        | 3        | 17       | 75       | 4        |          |
| $x_{94}$  | 0        | 4        | 0        | 0        | 0        | 3        | 14       | 3        | 3        | 2        |
| $x_{97}$  | 0        | 5        | 0        | 0        | 0        | 4        | 19       | 4        | 4        | 3        |
| $x_{99}$  | 0        | 9        | 0        | 0        | 0        | 29       | 31       | 6        | 7        | 5        |
| $x_{116}$ | 0        | 1        | 0        | 0        | 0        | 1        | 5        | 1        | 1        | 1        |
| $x_{117}$ | 0        | 1        | 0        | 0        | 0        | 0        | 2        | 0        | 0        | 0        |
| $x_{118}$ | 0        | 2        | 0        | 0        | 0        | 1        | 6        | 1        | 1        | 1        |
| $x_{149}$ | -49      | -57      | -26      | -41      | -5       | -19      | -58      | -26      | -14      | -23      |
|           | $x_{17}$ | $x_{18}$ | $x_{25}$ | $x_{29}$ | $x_{36}$ | $x_{43}$ | $x_{45}$ | $x_{68}$ | $x_{77}$ | $x_{93}$ |

|           |          |          |          |           |           |           |  |  |  |  |
|-----------|----------|----------|----------|-----------|-----------|-----------|--|--|--|--|
| $x_{97}$  | 3        |          |          |           |           |           |  |  |  |  |
| $x_{99}$  | 4        | 6        |          |           |           |           |  |  |  |  |
| $x_{116}$ | 1        | 1        | 1        |           |           |           |  |  |  |  |
| $x_{117}$ | 14       | 0        | 1        | 0         |           |           |  |  |  |  |
| $x_{118}$ | 1        | 32       | 2        | 0         | 0         |           |  |  |  |  |
| $x_{149}$ | -10      | -16      | -22      | -3        | -2        | -5        |  |  |  |  |
|           | $x_{94}$ | $x_{97}$ | $x_{99}$ | $x_{116}$ | $x_{117}$ | $x_{118}$ |  |  |  |  |

---

### $D^+$ BRANCHING RATIOS

Some now-obsolete measurements have been omitted from these Listings.

#### ———— c-quark decays ————

#### $\Gamma(c \rightarrow e^+ \text{ anything}) / \Gamma(c \rightarrow \text{ anything})$

For the Summary Table, we only use the average of  $e^+$  and  $\mu^+$  measurements from  $Z^0 \rightarrow c \bar{c}$  decays; see the second data block below.



| <u>VALUE</u>  | <u>EVTS</u> | <u>DOCUMENT ID</u>    | <u>TECN</u> | <u>COMMENT</u>             |
|---|-------------|-----------------------|-------------|----------------------------|
| <b><math>0.103 \pm 0.009^{+0.009}_{-0.008}</math></b> | 378         | <sup>1</sup> ABBIENDI | 99K OPAL    | $Z^0 \rightarrow c\bar{c}$ |

<sup>1</sup> ABBIENDI 99K uses the excess of right-sign over wrong-sign leptons opposite reconstructed  $D^*(2010)^+ \rightarrow D^0\pi^+$  decays in  $Z^0 \rightarrow c\bar{c}$ .

### $\Gamma(c \rightarrow \mu^+ \text{ anything})/\Gamma(c \rightarrow \text{ anything})$

For the Summary Table, we only use the average of  $e^+$  and  $\mu^+$  measurements from  $Z^0 \rightarrow c\bar{c}$  decays; see the next data block.

| <u>VALUE</u>                                    | <u>EVTS</u> | <u>DOCUMENT ID</u>    | <u>TECN</u> | <u>COMMENT</u>                                 |
|---|-------------|-----------------------|-------------|--|
| <b><math>0.082 \pm 0.005</math> OUR AVERAGE</b> |             |                       |             |  |
| $0.073 \pm 0.008 \pm 0.002$                     | 73          | KAYIS-TOPAK.05        | CHRS        | $\nu_\mu$ emulsion                             |
| $0.095 \pm 0.007^{+0.014}_{-0.013}$             | 2829        | ASTIER                | 00D NOMD    | $\nu_\mu \text{ Fe} \rightarrow \mu^- \mu^+ X$ |
| $0.090 \pm 0.007^{+0.007}_{-0.006}$             | 476         | <sup>1</sup> ABBIENDI | 99K OPAL    | $Z^0 \rightarrow c\bar{c}$                     |
| $0.086 \pm 0.017^{+0.008}_{-0.007}$             | 69          | <sup>2</sup> ALBRECHT | 92F ARG     | $e^+e^- \approx 10 \text{ GeV}$                |
| $0.078 \pm 0.009 \pm 0.012$                     |             | ONG                   | 88 MRK2     | $e^+e^- 29 \text{ GeV}$                        |
| $0.078 \pm 0.015 \pm 0.02$                      |             | BARTEL                | 87 JADE     | $e^+e^- 34.6 \text{ GeV}$                      |
| $0.082 \pm 0.012^{+0.02}_{-0.01}$               |             | ALTHOFF               | 84G TASS    | $e^+e^- 34.5 \text{ GeV}$                      |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                             |    |                |          |                      |
|-----------------------------|----|----------------|----------|----------------------|
| $0.093 \pm 0.009 \pm 0.009$ | 88 | KAYIS-TOPAK.02 | CHRS     | See KAYIS-TOPAKSU 05 |
| $0.089 \pm 0.018 \pm 0.025$ |    | BARTEL         | 85J JADE | See BARTEL 87        |

<sup>1</sup> ABBIENDI 99K uses the excess of right-sign over wrong-sign leptons opposite reconstructed  $D^*(2010)^+ \rightarrow D^0\pi^+$  decays in  $Z^0 \rightarrow c\bar{c}$ .

<sup>2</sup> ALBRECHT 92F uses the excess of right-sign over wrong-sign leptons in a sample of events tagged by fully reconstructed  $D^*(2010)^+ \rightarrow D^0\pi^+$  decays.

### $\Gamma(c \rightarrow \ell^+ \text{ anything})/\Gamma(c \rightarrow \text{ anything})$

This is an average (not a sum) of  $e^+$  and  $\mu^+$  measurements.

| <u>VALUE</u>                                    | <u>EVTS</u> | <u>DOCUMENT ID</u>    | <u>TECN</u> | <u>COMMENT</u>             |
|---|-------------|-----------------------|-------------|----------------------------|
| <b><math>0.096 \pm 0.004</math> OUR AVERAGE</b> |             |                       |             |                            |
| $0.0958 \pm 0.0042 \pm 0.0028$                  | 1828        | <sup>1</sup> ABREU    | 000 DLPH    | $Z^0 \rightarrow c\bar{c}$ |
| $0.095 \pm 0.006^{+0.007}_{-0.006}$             | 854         | <sup>2</sup> ABBIENDI | 99K OPAL    | $Z^0 \rightarrow c\bar{c}$ |

<sup>1</sup> ABREU 000 uses leptons opposite fully reconstructed  $D^*(2010)^+$ ,  $D^+$ , or  $D^0$  mesons.

<sup>2</sup> ABBIENDI 99K uses the excess of right-sign over wrong-sign leptons opposite reconstructed  $D^*(2010)^+ \rightarrow D^0\pi^+$  decays in  $Z^0 \rightarrow c\bar{c}$ .

### $\Gamma(c \rightarrow D^*(2010)^+ \text{ anything})/\Gamma(c \rightarrow \text{ anything})$

| <u>VALUE</u>                                  | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>             |
|---|-------------|--------------------|-------------|----------------------------|
| <b><math>0.255 \pm 0.015 \pm 0.008</math></b> | 2371        | <sup>1</sup> ABREU | 000 DLPH    | $Z^0 \rightarrow c\bar{c}$ |

<sup>1</sup> ABREU 000 uses slow pions opposite fully reconstructed  $D^*(2010)^+$ ,  $D^+$ , or  $D^0$  mesons as a signal of  $D^*(2010)^-$  production.

————— Inclusive modes —————

### $\Gamma(e^+ \text{ semileptonic})/\Gamma_{\text{total}}$ $\Gamma_1/\Gamma$

The sum of our  $\bar{K}^0 e^+ \nu_e$ ,  $\bar{K}^*(892)^0 e^+ \nu_e$ ,  $\pi^0 e^+ \nu_e$ ,  $\eta e^+ \nu_e$ ,  $\rho^0 e^+ \nu_e$ , and  $\omega e^+ \nu_e$  branching fractions is  $15.3 \pm 0.3\%$ .

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------|------|-------------|------|---------|
|-----------|------|-------------|------|---------|

#### 16.07±0.30 OUR AVERAGE

|                 |           |                    |          |                              |
|-----------------|-----------|--------------------|----------|------------------------------|
| 16.13±0.10±0.29 | 26.2±0.2k | <sup>1</sup> ASNER | 10 CLEO  | $e^+ e^-$ at 3774 MeV        |
| 15.2 ±0.9 ±0.8  | 521 ± 32  | ABLIKIM            | 07G BES2 | $e^+ e^- \approx \psi(3770)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                 |            |                   |          |                    |
|-----------------|------------|-------------------|----------|--------------------|
| 16.13±0.20±0.33 | 8798 ± 105 | <sup>2</sup> ADAM | 06A CLEO | See ASNER 10       |
| 17.0 ±1.9 ±0.7  | 158        | BALTRUSAIT..85B   | MRK3     | $e^+ e^-$ 3.77 GeV |

<sup>1</sup> Using the  $D^+$  and  $D^0$  lifetimes, ASNER 10 finds that the ratio of the  $D^+$  and  $D^0$  semileptonic widths is  $0.985 \pm 0.015 \pm 0.024$ .

<sup>2</sup> Using the  $D^+$  and  $D^0$  lifetimes, ADAM 06A finds that the ratio of the  $D^+$  and  $D^0$  inclusive  $e^+$  widths is  $0.985 \pm 0.028 \pm 0.015$ , consistent with the isospin-invariance prediction of 1.

### $\Gamma(\mu^+ \text{ anything})/\Gamma_{\text{total}}$ $\Gamma_2/\Gamma$

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------|------|-------------|------|---------|
|-----------|------|-------------|------|---------|

|                     |          |                      |          |                              |
|---------------------|----------|----------------------|----------|------------------------------|
| <b>17.6±2.7±1.8</b> | 100 ± 12 | <sup>1</sup> ABLIKIM | 08L BES2 | $e^+ e^- \approx \psi(3772)$ |
|---------------------|----------|----------------------|----------|------------------------------|

<sup>1</sup> ABLIKIM 08L finds the ratio of  $D^+ \rightarrow \mu^+ X$  and  $D^0 \rightarrow \mu^+ X$  branching fractions to be  $2.59 \pm 0.70 \pm 0.25$ , in accord with the ratio of  $D^+$  and  $D^0$  lifetimes,  $2.54 \pm 0.02$ .

### $\Gamma(K^- \text{ anything})/\Gamma_{\text{total}}$ $\Gamma_3/\Gamma$

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------|------|-------------|------|---------|
|-----------|------|-------------|------|---------|

#### 25.7±1.4 OUR AVERAGE

|                                      |          |         |          |                              |
|--------------------------------------|----------|---------|----------|------------------------------|
| 24.7±1.3±1.2                         | 631 ± 33 | ABLIKIM | 07G BES2 | $e^+ e^- \approx \psi(3770)$ |
| 27.8 <sup>+3.6</sup> <sub>-3.1</sub> |          | BARLAG  | 92C ACCM | $\pi^-$ Cu 230 GeV           |
| 27.1±2.3±2.4                         |          | COFFMAN | 91 MRK3  | $e^+ e^-$ 3.77 GeV           |

### $[\Gamma(\bar{K}^0 \text{ anything}) + \Gamma(K^0 \text{ anything})]/\Gamma_{\text{total}}$ $\Gamma_4/\Gamma$

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------|------|-------------|------|---------|
|-----------|------|-------------|------|---------|

#### 61 ±5 OUR AVERAGE

|              |          |         |          |                       |
|--------------|----------|---------|----------|-----------------------|
| 60.5±5.5±3.3 | 244 ± 22 | ABLIKIM | 06U BES2 | $e^+ e^-$ at 3773 MeV |
| 61.2±6.5±4.3 |          | COFFMAN | 91 MRK3  | $e^+ e^-$ 3.77 GeV    |

### $\Gamma(K^+ \text{ anything})/\Gamma_{\text{total}}$ $\Gamma_5/\Gamma$

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------|------|-------------|------|---------|
|-----------|------|-------------|------|---------|

#### 5.9±0.8 OUR AVERAGE

|             |          |         |          |                              |
|-------------|----------|---------|----------|------------------------------|
| 6.1±0.9±0.4 | 189 ± 27 | ABLIKIM | 07G BES2 | $e^+ e^- \approx \psi(3770)$ |
| 5.5±1.3±0.9 |          | COFFMAN | 91 MRK3  | $e^+ e^-$ 3.77 GeV           |

### $\Gamma(K^*(892)^- \text{ anything})/\Gamma_{\text{total}}$ $\Gamma_6/\Gamma$

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------|------|-------------|------|---------|
|-----------|------|-------------|------|---------|

|                    |           |         |          |                       |
|--------------------|-----------|---------|----------|-----------------------|
| <b>5.7±5.2±0.7</b> | 7.2 ± 6.5 | ABLIKIM | 06U BES2 | $e^+ e^-$ at 3773 MeV |
|--------------------|-----------|---------|----------|-----------------------|

$\Gamma(\bar{K}^*(892)^0 \text{ anything})/\Gamma_{\text{total}}$   $\Gamma_7/\Gamma$

| VALUE (%)               | EVTS     | DOCUMENT ID | TECN | COMMENT                            |
|-------------------------|----------|-------------|------|------------------------------------|
| <b>23.2 ± 4.5 ± 3.0</b> | 189 ± 36 | ABLIKIM 05P | BES  | $e^+ e^- \approx 3773 \text{ MeV}$ |

$\Gamma(K^*(892)^0 \text{ anything})/\Gamma_{\text{total}}$   $\Gamma_8/\Gamma$

| VALUE (%)      | CL% | DOCUMENT ID | TECN | COMMENT                            |
|----------------|-----|-------------|------|------------------------------------|
| <b>&lt;6.6</b> | 90  | ABLIKIM 05P | BES  | $e^+ e^- \approx 3773 \text{ MeV}$ |

$\Gamma(\eta \text{ anything})/\Gamma_{\text{total}}$   $\Gamma_9/\Gamma$

This ratio includes  $\eta$  particles from  $\eta'$  decays.

| VALUE (%)              | EVTS       | DOCUMENT ID | TECN | COMMENT                   |
|------------------------|------------|-------------|------|---------------------------|
| <b>6.3 ± 0.5 ± 0.5</b> | 1972 ± 142 | HUANG 06B   | CLEO | $e^+ e^-$ at $\psi(3770)$ |

$\Gamma(\eta' \text{ anything})/\Gamma_{\text{total}}$   $\Gamma_{10}/\Gamma$

| VALUE (%)                 | EVTS    | DOCUMENT ID | TECN | COMMENT                   |
|---------------------------|---------|-------------|------|---------------------------|
| <b>1.04 ± 0.16 ± 0.09</b> | 82 ± 13 | HUANG 06B   | CLEO | $e^+ e^-$ at $\psi(3770)$ |

$\Gamma(\phi \text{ anything})/\Gamma_{\text{total}}$   $\Gamma_{11}/\Gamma$

| VALUE (%)                 | EVTS     | DOCUMENT ID | TECN | COMMENT                   |
|---------------------------|----------|-------------|------|---------------------------|
| <b>1.03 ± 0.10 ± 0.07</b> | 248 ± 21 | HUANG 06B   | CLEO | $e^+ e^-$ at $\psi(3770)$ |

———— Leptonic and semileptonic modes ————

$\Gamma(e^+ \nu_e)/\Gamma_{\text{total}}$   $\Gamma_{12}/\Gamma$

| VALUE                            | CL% | DOCUMENT ID   | TECN | COMMENT                   |
|----------------------------------|-----|---------------|------|---------------------------|
| <b>&lt;8.8 × 10<sup>-6</sup></b> | 90  | EISENSTEIN 08 | CLEO | $e^+ e^-$ at $\psi(3770)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                         |    |            |      |                   |
|-------------------------|----|------------|------|-------------------|
| <2.4 × 10 <sup>-5</sup> | 90 | ARTUSO 05A | CLEO | See EISENSTEIN 08 |
|-------------------------|----|------------|------|-------------------|

$\Gamma(\gamma e^+ \nu_e)/\Gamma_{\text{total}}$   $\Gamma_{13}/\Gamma$

| VALUE                            | CL% | DOCUMENT ID              | TECN | COMMENT                |
|----------------------------------|-----|--------------------------|------|------------------------|
| <b>&lt;3.0 × 10<sup>-5</sup></b> | 90  | <sup>1</sup> ABLIKIM 17M | BES3 | $e^+ e^-$ at 3.773 GeV |

<sup>1</sup>This ABLIKIM 17M limit is for photons with energies greater than 10 MeV.

$\Gamma(\mu^+ \nu_\mu)/\Gamma_{\text{total}}$   $\Gamma_{14}/\Gamma$

See the note on "Decay Constants of Charged Pseudoscalar Mesons" in the  $D_s^+$  Listings.

| VALUE (units 10 <sup>-4</sup> ) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------------------|------|-------------|------|---------|
| <b>3.74 ± 0.17 OUR AVERAGE</b>  |      |             |      |         |

|                    |          |                            |      |                           |
|--------------------|----------|----------------------------|------|---------------------------|
| 3.71 ± 0.19 ± 0.06 | 409 ± 21 | <sup>1</sup> ABLIKIM 14F   | BES3 | $e^+ e^-$ at $\psi(3770)$ |
| 3.82 ± 0.32 ± 0.09 | 150 ± 12 | <sup>2</sup> EISENSTEIN 08 | CLEO | $e^+ e^-$ at $\psi(3770)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|   |   |                          |     |                                     |
|---|---|--------------------------|-----|-------------------------------------|
| 12.2 <sup>+11.1</sup> / <sub>-5.3</sub> ± 1.0 | 3 | <sup>3</sup> ABLIKIM 05D | BES | $e^+ e^- \approx 3.773 \text{ GeV}$ |
|---|---|--------------------------|-----|-------------------------------------|

|   |        |                         |      |                   |
|---|--------|-------------------------|------|-------------------|
| 4.40 ± 0.66 <sup>+0.09</sup> / <sub>-0.12</sub> | 47 ± 7 | <sup>4</sup> ARTUSO 05A | CLEO | See EISENSTEIN 08 |
|---|--------|-------------------------|------|-------------------|

|                 |   |                            |      |                     |
|-----------------|---|----------------------------|------|---------------------|
| 3.5 ± 1.4 ± 0.6 | 7 | <sup>5</sup> BONVICINI 04A | CLEO | Incl. in ARTUSO 05A |
|-----------------|---|----------------------------|------|---------------------|

|  |   |                      |     |                                  |
|--|---|----------------------|-----|----------------------------------|
| 8 <sup>+16</sup> / <sub>-5</sub> <sup>+5</sup> / <sub>-2</sub> | 1 | <sup>6</sup> BAI 98B | BES | $e^+ e^- \rightarrow D^{*+} D^-$ |
|--|---|----------------------|-----|----------------------------------|

- <sup>1</sup> ABLIKIM 14F obtain  $|V_{cd}| \cdot f_{D^+} = (45.75 \pm 1.20 \pm 0.39)$  MeV, and using  $|V_{cd}| = 0.22520 \pm 0.00065$  gets  $f_{D^+} = (203.2 \pm 5.3 \pm 1.8)$  MeV.
- <sup>2</sup> EISENSTEIN 08, using the  $D^+$  lifetime and assuming  $|V_{cd}| = |V_{us}|$ , gets  $f_{D^+} = (205.8 \pm 8.5 \pm 2.5)$  MeV from this measurement.
- <sup>3</sup> ABLIKIM 05D finds a background-subtracted  $2.67 \pm 1.74$   $D^+ \rightarrow \mu^+ \nu_\mu$  events, and from this obtains  $f_{D^+} = 371_{-119}^{+129} \pm 25$  MeV.
- <sup>4</sup> ARTUSO 05A obtains  $f_{D^+} = 222.6 \pm 16.7_{-3.4}^{+2.8}$  MeV from this measurement.
- <sup>5</sup> BONVICINI 04A finds eight events with an estimated background of one, and from the branching fraction obtains  $f_{D^+} = 202 \pm 41 \pm 17$  MeV.
- <sup>6</sup> BAI 98B obtains  $f_{D^+} = (300_{-150}^{+180} +_{-40}^{+80})$  MeV from this measurement.

| $\Gamma(\tau^+ \nu_\tau)/\Gamma_{\text{total}}$                               |     |               |      |                           | $\Gamma_{15}/\Gamma$ |
|---|-----|---------------|------|---------------------------|----------------------|
| VALUE   | CL% | DOCUMENT ID   | TECN | COMMENT                   |                      |
| $<1.2 \times 10^{-3}$   | 90  | EISENSTEIN 08 | CLEO | $e^+ e^-$ at $\psi(3770)$ |                      |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |     |               |      |                           |                      |
| $<2.1 \times 10^{-3}$   | 90  | RUBIN 06A     | CLEO | See EISENSTEIN 08         |                      |

| $\Gamma(\bar{K}^0 e^+ \nu_e)/\Gamma_{\text{total}}$                           |      |                           |      |   | $\Gamma_{16}/\Gamma$ |
|---|------|---------------------------|------|---|----------------------|
| VALUE (%)   | EVTS | DOCUMENT ID               | TECN | COMMENT                                   |                      |
| <b>8.73 ± 0.10 OUR AVERAGE</b>  |      |                           |      |   |                      |
| $8.60 \pm 0.06 \pm 0.15$  | 26k  | ABLIKIM 17S               | BES3 | Using $\bar{K}^0 \rightarrow \pi^+ \pi^-$ |                      |
| $8.59 \pm 0.14 \pm 0.21$  | 5013 | ABLIKIM 16V               | BES3 | Using $\bar{K}^0 \rightarrow 2\pi^0$      |                      |
| $8.962 \pm 0.054 \pm 0.206$   | 40k  | <sup>1</sup> ABLIKIM 15AF | BES3 | from $D^+ \rightarrow K_L e^+ \nu_e$      |                      |
| $8.83 \pm 0.10 \pm 0.20$  | 8.5k | <sup>2</sup> BESSON 09    | CLEO | from $D^+ \rightarrow K_S e^+ \nu_e$      |                      |
| $8.95 \pm 1.59 \pm 0.67$  | 34   | <sup>3</sup> ABLIKIM 05A  | BES  | from $D^+ \rightarrow K_S e^+ \nu_e$      |                      |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |      |                           |      |   |                      |
| $8.53 \pm 0.13 \pm 0.23$  |      | <sup>4</sup> DOBBS 08     | CLEO | See BESSON 09                             |                      |
| $8.71 \pm 0.38 \pm 0.37$  | 545  | HUANG 05B                 | CLEO | See DOBBS 08                              |                      |

- <sup>1</sup> ABLIKIM 15AF report  $\Gamma(D^+ \rightarrow K_L e^+ \nu_e)/\Gamma_{\text{total}} = (4.481 \pm 0.027 \pm 0.103)\%$ . See also the form-factor parameters near the end of this  $D^+$  Listing.
- <sup>2</sup> See the form-factor parameters near the end of this  $D^+$  Listing.
- <sup>3</sup> The ABLIKIM 05A result together with the  $D^0 \rightarrow K^- e^+ \nu_e$  branching fraction of ABLIKIM 04C and Particle Data Group lifetimes gives  $\Gamma(D^0 \rightarrow K^- e^+ \nu_e) / \Gamma(D^+ \rightarrow \bar{K}^0 e^+ \nu_e) = 1.08 \pm 0.22 \pm 0.07$ ; isospin invariance predicts the ratio is 1.0.
- <sup>4</sup> DOBBS 08 establishes  $|\frac{V_{cd}}{V_{cs}} \cdot \frac{f_+^\pi(0)}{f_+^K(0)}| = 0.188 \pm 0.008 \pm 0.002$  from the  $D^+$  and  $D^0$  decays to  $\bar{K} e^+ \nu_e$  and  $\pi e^+ \nu_e$ . It also finds  $\Gamma(D^0 \rightarrow K^- e^+ \nu_e) / \Gamma(D^+ \rightarrow \bar{K}^0 e^+ \nu_e) = 1.06 \pm 0.02 \pm 0.03$ ; isospin invariance predicts the ratio is 1.0.

| $\Gamma(\bar{K}^0 \mu^+ \nu_\mu)/\Gamma_{\text{total}}$                       |            |             |      |                       | $\Gamma_{17}/\Gamma$ |
|---|------------|-------------|------|-----------------------|----------------------|
| VALUE (units $10^{-2}$ )  | EVTS       | DOCUMENT ID | TECN | COMMENT               |                      |
| <b>8.76 ± 0.19 OUR FIT</b>  |            |             |      |                       |                      |
| <b>8.72 ± 0.07 ± 0.18</b>   | 21k        | ABLIKIM 16G | BES3 | $e^+ e^-$ at 3773 MeV |                      |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |            |             |      |                       |                      |
| $10.3 \pm 2.3 \pm 0.8$  | $29 \pm 6$ | ABLIKIM 07  | BES2 | $e^+ e^-$ at 3773 MeV |                      |

$\Gamma(\bar{K}^0 \mu^+ \nu_\mu) / \Gamma(K^- 2\pi^+) \quad \Gamma_{17} / \Gamma_{45}$ 

| VALUE                        | EVTS     | DOCUMENT ID | TECN     | COMMENT  |
|------------------------------|----------|-------------|----------|--|
| <b>0.934 ± 0.025 OUR FIT</b> |          |             |          | Error includes scale factor of 1.2.                |
| <b>1.019 ± 0.076 ± 0.065</b> | 555 ± 39 | LINK        | 04E FOCS | $\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV |

 $\Gamma(K^- \pi^+ e^+ \nu_e) / \Gamma_{\text{total}} \quad \Gamma_{18} / \Gamma$ 

| VALUE (units $10^{-2}$ )  | EVTS  | DOCUMENT ID | TECN     | COMMENT                             |
|---|-------|-------------|----------|-------------------------------------|
| <b>4.02 ± 0.18 OUR FIT</b>  |       |             |          | Error includes scale factor of 3.2. |
| <b>3.77 ± 0.03 ± 0.08</b>   | 18.3k | ABLIKIM     | 16F BES3 | $e^+ e^-$ at $\psi(3770)$           |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |       |             |          |                                     |
| 3.50 ± 0.75 ± 0.27  | 29    | ABLIKIM     | 060 BES2 | $e^+ e^-$ at 3773 MeV               |
| 3.5 $^{+1.2}_{-0.7}$ ± 0.4  | 14    | BAI         | 91 MRK3  | $e^+ e^- \approx 3.77$ GeV          |

 $\Gamma(K^- \pi^+ e^+ \nu_e) / \Gamma(K^- 2\pi^+) \quad \Gamma_{18} / \Gamma_{45}$ 

| VALUE                           | EVTS      | DOCUMENT ID     | TECN | COMMENT                             |
|---------------------------------|-----------|-----------------|------|-------------------------------------|
| <b>0.428 ± 0.018 OUR FIT</b>    |           |                 |      | Error includes scale factor of 3.7. |
| <b>0.4380 ± 0.0036 ± 0.0042</b> | 70k ± 363 | DEL-AMO-SA..11I | BABR | $e^+ e^- \approx 10.6$ GeV          |

 $\Gamma(\bar{K}^*(892)^0 e^+ \nu_e, \bar{K}^*(892)^0 \rightarrow K^- \pi^+) / \Gamma(K^- \pi^+ e^+ \nu_e) \quad \Gamma_{19} / \Gamma_{18}$ 

| VALUE (%)                       | DOCUMENT ID     | TECN     | COMMENT                    |
|---------------------------------|-----------------|----------|----------------------------|
| <b>93.94 ± 0.27 OUR AVERAGE</b> |                 |          |                            |
| 93.93 ± 0.22 ± 0.18             | ABLIKIM         | 16F BES3 | $e^+ e^-$ at $\psi(3770)$  |
| 94.11 ± 0.74 ± 0.75             | DEL-AMO-SA..11I | BABR     | $e^+ e^- \approx 10.6$ GeV |

 $\Gamma((K^- \pi^+)_{[0.8-1.0]\text{GeV}} e^+ \nu_e) / \Gamma_{\text{total}} \quad \Gamma_{20} / \Gamma$ 

| VALUE (units $10^{-2}$ )  | EVTS  | DOCUMENT ID | TECN     | COMMENT                   |
|---------------------------|-------|-------------|----------|---------------------------|
| <b>3.39 ± 0.03 ± 0.08</b> | 16.2k | ABLIKIM     | 16F BES3 | $e^+ e^-$ at $\psi(3770)$ |

 $\Gamma((K^- \pi^+)_{S\text{-wave}} e^+ \nu_e) / \Gamma_{\text{total}} \quad \Gamma_{21} / \Gamma$ 

| VALUE (units $10^{-3}$ )  | DOCUMENT ID | TECN     | COMMENT                   |
|---------------------------|-------------|----------|---------------------------|
| <b>2.28 ± 0.08 ± 0.08</b> | ABLIKIM     | 16F BES3 | $e^+ e^-$ at $\psi(3770)$ |

 $\Gamma((K^- \pi^+)_{S\text{-wave}} e^+ \nu_e) / \Gamma(K^- \pi^+ e^+ \nu_e) \quad \Gamma_{21} / \Gamma_{18}$ 

| VALUE (%)                      | DOCUMENT ID     | TECN     | COMMENT                    |
|--------------------------------|-----------------|----------|----------------------------|
| <b>5.89 ± 0.17 OUR AVERAGE</b> |                 |          |                            |
| 6.05 ± 0.22 ± 0.18             | ABLIKIM         | 16F BES3 | $e^+ e^-$ at $\psi(3770)$  |
| 5.79 ± 0.16 ± 0.15             | DEL-AMO-SA..11I | BABR     | $e^+ e^- \approx 10.6$ GeV |

 $\Gamma(\bar{K}^*(1410)^0 e^+ \nu_e, \bar{K}^*(1410)^0 \rightarrow K^- \pi^+) / \Gamma_{\text{total}} \quad \Gamma_{22} / \Gamma$ 

| VALUE                           | CL% | DOCUMENT ID     | TECN | COMMENT                    |
|---------------------------------|-----|-----------------|------|----------------------------|
| <b>&lt; 6 × 10<sup>-3</sup></b> | 90  | DEL-AMO-SA..11I | BABR | $e^+ e^- \approx 10.6$ GeV |

 $\Gamma(\bar{K}_2^*(1430)^0 e^+ \nu_e, \bar{K}_2^*(1430)^0 \rightarrow K^- \pi^+) / \Gamma_{\text{total}} \quad \Gamma_{23} / \Gamma$ 

| VALUE                           | CL% | DOCUMENT ID     | TECN | COMMENT                    |
|---------------------------------|-----|-----------------|------|----------------------------|
| <b>&lt; 5 × 10<sup>-4</sup></b> | 90  | DEL-AMO-SA..11I | BABR | $e^+ e^- \approx 10.6$ GeV |

$\Gamma(K^- \pi^+ e^+ \nu_e \text{ nonresonant})/\Gamma_{\text{total}}$   $\Gamma_{24}/\Gamma$ 

| VALUE  | CL% | DOCUMENT ID | TECN     | COMMENT         |
|--------|-----|-------------|----------|-----------------|
| <0.007 | 90  | ANJOS       | 89B E691 | Photoproduction |

 $\Gamma(\bar{K}^*(892)^0 e^+ \nu_e)/\Gamma_{\text{total}}$   $\Gamma_{25}/\Gamma$ 

Unseen decay modes of  $\bar{K}^*(892)^0$  are included. See the end of the  $D^+$  Listings for measurements of  $D^+ \rightarrow \bar{K}^*(892)^0 \ell^+ \nu_\ell$  form-factor ratios.

| VALUE (units $10^{-2}$ ) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|

**5.40±0.10 OUR FIT** Error includes scale factor of 1.1.

**5.40±0.10 OUR AVERAGE** Error includes scale factor of 1.1.

5.31±0.05±0.12 16.2k ABLIKIM 16F BES3  $e^+ e^-$  at  $\psi(3770)$

5.52±0.07±0.13  $\approx 5k$  BRIERE 10 CLEO  $e^+ e^-$  at  $\psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.06±1.21±0.40  $28 \pm 7$  ABLIKIM 06O BES2  $e^+ e^-$  at 3773 MeV

5.56±0.27±0.23  $422 \pm 21$  <sup>1</sup>HUANG 05B CLEO  $e^+ e^-$  at  $\psi(3770)$

<sup>1</sup>HUANG 05B finds  $\Gamma(D^0 \rightarrow K^{*-} e^+ \nu_e) / \Gamma(D^+ \rightarrow \bar{K}^{*0} e^+ \nu_e) = 0.98 \pm 0.08 \pm 0.04$ ; isospin invariance predicts the ratio is 1.0.

 $\Gamma(\bar{K}^*(892)^0 e^+ \nu_e)/\Gamma(K^- 2\pi^+)$   $\Gamma_{25}/\Gamma_{45}$ 

Unseen decay modes of the  $\bar{K}^*(892)^0$  are included. See the end of the  $D^+$  Listings for measurements of  $D^+ \rightarrow \bar{K}^*(892)^0 \ell^+ \nu_\ell$  form-factor ratios.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------|------|-------------|------|---------|
|-------|------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.74±0.04±0.05 BRANDENB... 02 CLEO  $e^+ e^- \approx \Upsilon(4S)$

0.62±0.15±0.09 35 ADAMOVICH 91 OMEG  $\pi^-$  340 GeV

0.55±0.08±0.10 880 ALBRECHT 91 ARG  $e^+ e^- \approx 10.4$  GeV

0.49±0.04±0.05 ANJOS 89B E691 Photoproduction

 $\Gamma(K^- \pi^+ \mu^+ \nu_\mu)/\Gamma(\bar{K}^0 \mu^+ \nu_\mu)$   $\Gamma_{26}/\Gamma_{17}$ 

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------|------|-------------|------|---------|
|-------|------|-------------|------|---------|

**0.417±0.030±0.023**  $555 \pm 39$  LINK 04E FOCS  $\gamma$  nucleus,  $\bar{E}_\gamma \approx 180$  GeV

 $\Gamma(K^- \pi^+ \mu^+ \nu_\mu \text{ nonresonant})/\Gamma(K^- \pi^+ \mu^+ \nu_\mu)$   $\Gamma_{28}/\Gamma_{26}$ 

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------|------|-------------|------|---------|
|-------|------|-------------|------|---------|

**0.0530±0.0074<sup>+0.0099</sup><sub>-0.0096</sub>** 14k LINK 05I FOCS  $\gamma$  nucleus,  $\bar{E}_\gamma \approx 180$  GeV

 $\Gamma(\bar{K}^*(892)^0 \mu^+ \nu_\mu)/\Gamma_{\text{total}}$   $\Gamma_{29}/\Gamma$ 

Unseen decay modes of the  $\bar{K}^*(892)^0$  are included. See the end of the  $D^+$  Listings for measurements of  $D^+ \rightarrow \bar{K}^*(892)^0 \ell^+ \nu_\ell$  form-factor ratios.

| VALUE (units $10^{-2}$ ) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|

**5.27±0.15 OUR FIT**

**5.27±0.07±0.14**  $\approx 5k$  BRIERE 10 CLEO  $e^+ e^-$  at  $\psi(3770)$

$\Gamma(\bar{K}^*(892)^0 \mu^+ \nu_\mu) / \Gamma(\bar{K}^0 \mu^+ \nu_\mu)$   $\Gamma_{29} / \Gamma_{17}$

Unseen decay modes of the  $\bar{K}^*(892)^0$  are included. See the end of the  $D^+$  Listings for measurements of  $D^+ \rightarrow \bar{K}^*(892)^0 \ell^+ \nu_\ell$  form-factor ratios.

| VALUE                        | EVTS     | DOCUMENT ID | TECN     | COMMENT  |
|------------------------------|----------|-------------|----------|--|
| <b>0.602 ± 0.021 OUR FIT</b> |          |             |          |  |
| <b>0.594 ± 0.043 ± 0.033</b> | 555 ± 39 | LINK        | 04E FOCS | $\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV |

$\Gamma(\bar{K}^*(892)^0 \mu^+ \nu_\mu) / \Gamma(K^- 2\pi^+)$   $\Gamma_{29} / \Gamma_{45}$

Unseen decay modes of the  $\bar{K}^*(892)^0$  are included. See the end of the  $D^+$  Listings for measurements of  $D^+ \rightarrow \bar{K}^*(892)^0 \ell^+ \nu_\ell$  form-factor ratios.

| VALUE   | EVTS | DOCUMENT ID       | TECN     | COMMENT                                      |
|---|------|-------------------|----------|--|
| <b>0.562 ± 0.018 OUR FIT</b>  |      |                   |          | Error includes scale factor of 1.1.          |
| <b>0.57 ± 0.06 OUR AVERAGE</b>  |      |                   |          | Error includes scale factor of 1.2.          |
| 0.72 ± 0.10 ± 0.05  |      | BRANDENB...       | 02 CLEO  | $e^+ e^- \approx \mathcal{R}(4S)$            |
| 0.56 ± 0.04 ± 0.06  | 875  | FRABETTI          | 93E E687 | $\gamma$ Be $\bar{E}_\gamma \approx 200$ GeV |
| 0.46 ± 0.07 ± 0.08  | 224  | KODAMA            | 92C E653 | $\pi^-$ emulsion 600 GeV                     |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |      |                   |          |  |
| 0.602 ± 0.010 ± 0.021   | 12k  | <sup>1</sup> LINK | 02J FOCS | $\gamma$ nucleus, $\approx 180$ GeV          |

<sup>1</sup>This LINK 02J result includes the effects of an interference of a small  $S$ -wave  $K^- \pi^+$  amplitude with the dominant  $\bar{K}^{*0}$  amplitude. (The interference effect is reported in LINK 02E.) This result is redundant with results of LINK 04E elsewhere in these Listings.

$\Gamma(K^- \pi^+ \pi^0 \mu^+ \nu_\mu) / \Gamma(K^- \pi^+ \mu^+ \nu_\mu)$   $\Gamma_{30} / \Gamma_{26}$

| VALUE            | CL% | DOCUMENT ID | TECN     | COMMENT                                      |
|------------------|-----|-------------|----------|--|
| <b>&lt;0.042</b> | 90  | FRABETTI    | 93E E687 | $\gamma$ Be $\bar{E}_\gamma \approx 200$ GeV |

$\Gamma(\bar{K}_0^*(1430)^0 \mu^+ \nu_\mu) / \Gamma(K^- \pi^+ \mu^+ \nu_\mu)$   $\Gamma_{31} / \Gamma_{26}$

Unseen decay modes of the  $\bar{K}_0^*(1430)^0$  are included.

| VALUE             | CL% | DOCUMENT ID | TECN     | COMMENT                                      |
|-------------------|-----|-------------|----------|--|
| <b>&lt;0.0064</b> | 90  | LINK        | 05I FOCS | $\gamma$ A, $\bar{E}_\gamma \approx 180$ GeV |

$\Gamma(\bar{K}^*(1680)^0 \mu^+ \nu_\mu) / \Gamma(K^- \pi^+ \mu^+ \nu_\mu)$   $\Gamma_{32} / \Gamma_{26}$

Unseen decay modes of the  $\bar{K}^*(1680)^0$  are included.

| VALUE           | CL% | DOCUMENT ID | TECN     | COMMENT                                      |
|-----------------|-----|-------------|----------|--|
| <b>&lt;0.04</b> | 90  | LINK        | 05I FOCS | $\gamma$ A, $\bar{E}_\gamma \approx 180$ GeV |

$\Gamma(\pi^0 e^+ \nu_e) / \Gamma_{\text{total}}$   $\Gamma_{33} / \Gamma$

| VALUE (%)   | EVTS   | DOCUMENT ID         | TECN     | COMMENT                             |
|---|--------|---------------------|----------|-------------------------------------|
| <b>0.372 ± 0.017 OUR AVERAGE</b>  |        |                     |          | Error includes scale factor of 2.0. |
| 0.363 ± 0.008 ± 0.005   | 3.4k   | ABLIKIM             | 17S BES3 | Using $\pi^0 \rightarrow 2\gamma$   |
| 0.405 ± 0.016 ± 0.009   | 838    | <sup>1</sup> BESSON | 09 CLEO  | $e^+ e^-$ at $\psi(3770)$           |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |        |                     |          |                                     |
| 0.373 ± 0.022 ± 0.013   |        | <sup>2</sup> DOBBS  | 08 CLEO  | See BESSON 09                       |
| 0.44 ± 0.06 ± 0.03  | 63 ± 9 | HUANG               | 05B CLEO | See DOBBS 08                        |

<sup>1</sup>See the form-factor parameters near the end of this  $D^+$  Listing.

<sup>2</sup>DOBBS 08 establishes  $|\frac{V_{cd}}{V_{cs}} \cdot \frac{f_\pi^+(0)}{f_K^+(0)}| = 0.188 \pm 0.008 \pm 0.002$  from the  $D^+$  and  $D^0$  decays to  $\bar{K} e^+ \nu_e$  and  $\pi e^+ \nu_e$ . It finds  $\Gamma(D^0 \rightarrow \pi^- e^+ \nu_e) / \Gamma(D^+ \rightarrow \pi^0 e^+ \nu_e) = 2.03 \pm 0.14 \pm 0.08$ ; isospin invariance predicts the ratio is 2.0.

$\Gamma(\pi^0 \mu^+ \nu_\mu) / \Gamma_{\text{total}}$   $\Gamma_{34} / \Gamma$

| VALUE (%)                    | EVTS | DOCUMENT ID | TECN      | COMMENT              |
|------------------------------|------|-------------|-----------|----------------------|
| <b>0.350 ± 0.011 ± 0.010</b> | 1.3k | ABLIKIM     | 18AE BES3 | $e^+ e^-$ , 3773 MeV |

$\Gamma(\eta e^+ \nu_e) / \Gamma_{\text{total}}$   $\Gamma_{35} / \Gamma$

| VALUE (units $10^{-4}$ )  | EVTS | DOCUMENT ID | TECN     | COMMENT                   |
|---|------|-------------|----------|---------------------------|
| <b>11.1 ± 0.7 OUR AVERAGE</b>   |      |             |          |                           |
| 10.74 ± 0.81 ± 0.51   | 373  | ABLIKIM     | 18R BES3 | $e^+ e^-$ , 3773 MeV      |
| 11.4 ± 0.9 ± 0.4  |      | YELTON      | 11 CLEO  | $e^+ e^-$ at $\psi(3770)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |      |             |          |                           |
| 13.3 ± 2.0 ± 0.6  | 46   | MITCHELL    | 09B CLEO | See YELTON 11             |

$\Gamma(\rho^0 e^+ \nu_e) / \Gamma_{\text{total}}$   $\Gamma_{36} / \Gamma$

| VALUE (units $10^{-3}$ )                            | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|------|---------|
| <b>2.18<sup>+0.17</sup><sub>-0.25</sub> OUR FIT</b> |      |             |      |         |

|   |          |                    |          |                           |
|---|----------|--------------------|----------|---------------------------|
| <b>2.17 ± 0.12<sup>+0.12</sup><sub>-0.22</sub></b>                            | 447 ± 25 | <sup>1</sup> DOBBS | 13 CLEO  | $e^+ e^-$ at $\psi(3770)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |          |                    |          |                           |
| 2.1 ± 0.4 ± 0.1   | 27 ± 6   | <sup>2</sup> HUANG | 05B CLEO | See DOBBS 13              |

<sup>1</sup> DOBBS 13 finds  $\Gamma(D^0 \rightarrow \rho^- e^+ \nu_e) / 2 \Gamma(D^+ \rightarrow \rho^0 e^+ \nu_e) = 1.03 \pm 0.09^{+0.08}_{-0.02}$ ; isospin invariance predicts the ratio is 1.0.

<sup>2</sup> HUANG 05B finds  $\Gamma(D^0 \rightarrow \rho^- e^+ \nu_e) / 2 \Gamma(D^+ \rightarrow \rho^0 e^+ \nu_e) = 1.2^{+0.4}_{-0.3} \pm 0.1$ ; isospin invariance predicts the ratio is 1.0.

$\Gamma(\rho^0 e^+ \nu_e) / \Gamma(\bar{K}^*(892)^0 e^+ \nu_e)$   $\Gamma_{36} / \Gamma_{25}$

| VALUE   | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|------|---------|
| <b>0.0404<sup>+0.0033</sup><sub>-0.0050</sub> OUR FIT</b> |      |             |      |         |

|   |    |                     |         |                          |
|---|----|---------------------|---------|--------------------------|
| <b>0.045 ± 0.014 ± 0.009</b>  | 49 | <sup>1</sup> AITALA | 97 E791 | $\pi^-$ nucleus, 500 GeV |
| <sup>1</sup> AITALA 97 explicitly subtracts $D^+ \rightarrow \eta' e^+ \nu_e$ and other backgrounds to get this result. |    |                     |         |                          |

$\Gamma(\rho^0 \mu^+ \nu_\mu) / \Gamma(\bar{K}^*(892)^0 \mu^+ \nu_\mu)$   $\Gamma_{37} / \Gamma_{29}$

| VALUE  | EVTS     | DOCUMENT ID           | TECN     | COMMENT                                       |
|--|----------|-----------------------|----------|---|
| <b>0.045 ± 0.007 OUR AVERAGE</b> Error includes scale factor of 1.1. |          |                       |          |   |
| 0.041 ± 0.006 ± 0.004  | 320 ± 44 | LINK                  | 06B FOCS | $\gamma$ A, $\bar{E}_\gamma \approx 180$ GeV  |
| 0.051 ± 0.015 ± 0.009  | 54       | <sup>1</sup> AITALA   | 97 E791  | $\pi^-$ nucleus, 500 GeV                      |
| 0.079 ± 0.019 ± 0.013  | 39       | <sup>2</sup> FRABETTI | 97 E687  | $\gamma$ Be, $\bar{E}_\gamma \approx 220$ GeV |

<sup>1</sup> AITALA 97 explicitly subtracts  $D^+ \rightarrow \eta' \mu^+ \nu_\mu$  and other backgrounds to get this result.

<sup>2</sup> Because the reconstruction efficiency for photons is low, this FRABETTI 97 result also includes any  $D^+ \rightarrow \eta' \mu^+ \nu_\mu \rightarrow \gamma \rho^0 \mu^+ \nu_\mu$  events in the numerator.



$\Gamma(\omega e^+ \nu_e)/\Gamma_{\text{total}}$   $\Gamma_{38}/\Gamma$ 

| VALUE (units $10^{-3}$ )  | EVTS                                | DOCUMENT ID | TECN     | COMMENT                                       |
|---|-------------------------------------|-------------|----------|---|
| <b>1.69 ± 0.11 OUR AVERAGE</b>  |                                     |             |          |   |
| 1.63 ± 0.11 ± 0.08  | 491 ± 32                            | ABLIKIM     | 15W BES3 | 292 fb <sup>-1</sup> , 3773 MeV               |
| 1.82 ± 0.18 ± 0.07  | 129 ± 13                            | DOBBS       | 13 CLEO  | e <sup>+</sup> e <sup>-</sup> at $\psi(3770)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                                     |             |          |   |
| 1.6 <sup>+0.7</sup> <sub>-0.6</sub> ± 0.1                                     | 7.6 <sup>+3.3</sup> <sub>-2.7</sub> | HUANG       | 05B CLEO | See DOBBS 13                                  |

 $\Gamma(\eta'(958) e^+ \nu_e)/\Gamma_{\text{total}}$   $\Gamma_{39}/\Gamma$ 

| VALUE (units $10^{-4}$ )  | CL% | EVTS | DOCUMENT ID | TECN     | COMMENT                                       |
|---|-----|------|-------------|----------|---|
| <b>2.0 ± 0.4 OUR AVERAGE</b>  |     |      |             |          |   |
| 1.91 ± 0.51 ± 0.13  |     | 32   | ABLIKIM     | 18R BES3 | e <sup>+</sup> e <sup>-</sup> , 3773 MeV      |
| 2.16 ± 0.53 ± 0.07  |     |      | YELTON      | 11 CLEO  | e <sup>+</sup> e <sup>-</sup> at $\psi(3770)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |     |      |             |          |   |
| <3.5  |     | 90   | MITCHELL    | 09B CLEO | See YELTON 11                                 |

 $\Gamma(a(980)^0 e^+ \nu_e, a(980)^0 \rightarrow \eta \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{40}/\Gamma$ 

| VALUE (units $10^{-4}$ )                            | EVTS                           | DOCUMENT ID          | TECN     | COMMENT                                   |
|---|--------------------------------|----------------------|----------|---|
| <b>1.66 <sup>+0.81</sup><sub>-0.66</sub> ± 0.11</b> | 10 <sup>+5</sup> <sub>-4</sub> | <sup>1</sup> ABLIKIM | 18F BES3 | e <sup>+</sup> e <sup>-</sup> at 3773 MeV |

<sup>1</sup>Signal observed at 2.9  $\sigma$  C.L. $\Gamma(\phi e^+ \nu_e)/\Gamma_{\text{total}}$   $\Gamma_{41}/\Gamma$ Unseen decay modes of the  $\phi$  are included.

| VALUE   | CL% | DOCUMENT ID | TECN     | COMMENT                                       |
|---|-----|-------------|----------|---|
| <b>&lt;1.3 × 10<sup>-5</sup></b>  | 90  | ABLIKIM     | 15W BES3 | 292 fb <sup>-1</sup> , 3773 MeV               |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |     |             |          |   |
| <0.9 × 10 <sup>-4</sup>   | 90  | YELTON      | 11 CLEO  | e <sup>+</sup> e <sup>-</sup> at $\psi(3770)$ |
| <1.6 × 10 <sup>-4</sup>   | 90  | MITCHELL    | 09B CLEO | See YELTON 11                                 |
| <0.0201   | 90  | ABLIKIM     | 06P BES2 | e <sup>+</sup> e <sup>-</sup> at 3773 MeV     |
| <0.0209   | 90  | BAI         | 91 MRK3  | e <sup>+</sup> e <sup>-</sup> ≈ 3.77 GeV      |

 $\Gamma(D^0 e^+ \nu_e)/\Gamma_{\text{total}}$   $\Gamma_{42}/\Gamma$ 

| VALUE                            | CL% | DOCUMENT ID | TECN      | COMMENT                                    |
|----------------------------------|-----|-------------|-----------|--|
| <b>&lt;1.0 × 10<sup>-4</sup></b> | 90  | ABLIKIM     | 17AD BES3 | e <sup>+</sup> e <sup>-</sup> at 3.773 GeV |

————— Hadronic modes with a  $\bar{K}$  or  $\bar{K}K\bar{K}$  ————— $\Gamma(K_S^0 \pi^+)/\Gamma_{\text{total}}$   $\Gamma_{43}/\Gamma$ 

| VALUE (units $10^{-2}$ )  | EVTS                                | DOCUMENT ID        | TECN     | COMMENT                                  |
|---|-------------------------------------|--------------------|----------|--|
| <b>1.562 ± 0.031 OUR FIT</b>  | Error includes scale factor of 1.7. |                    |          |  |
| <b>1.591 ± 0.006 ± 0.030</b>  | 94k                                 | ABLIKIM            | 18W BES3 | e <sup>+</sup> e <sup>-</sup> , 3773 MeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                                     |                    |          |  |
| 1.526 ± 0.022 ± 0.038   |                                     | <sup>1</sup> DOBBS | 07 CLEO  | See MENDEZ 10                            |
| 1.55 ± 0.05 ± 0.06  | 2.2k                                | <sup>1</sup> HE    | 05 CLEO  | See DOBBS 07                             |
| 1.6 ± 0.3 ± 0.1   | 161                                 | ADLER              | 88C MRK3 | e <sup>+</sup> e <sup>-</sup> 3.77 GeV   |

<sup>1</sup>DOBBS 07 and HE 05 use single- and double-tagged events in an overall fit. DOBBS 07 supersedes HE 05.

$\Gamma(K_S^0 \pi^+)/\Gamma(K^- 2\pi^+)$  $\Gamma_{43}/\Gamma_{45}$ 

| VALUE   | EVTS  | DOCUMENT ID         | TECN     | COMMENT  |
|---|-------|---------------------|----------|--|
| <b>0.167 ± 0.004 OUR FIT</b>  |       |                     |          | Error includes scale factor of 2.4.                |
| <b>0.162 ± 0.009 OUR AVERAGE</b>  |       |                     |          | Error includes scale factor of 4.5.                |
| 0.171 ± 0.002 ± 0.002   |       | BONVICINI 14        | CLEO     | All CLEO-c runs                                    |
| 0.1530 ± 0.0023 ± 0.0016  | 10.6k | LINK                | 02B FOCS | $\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |       |                     |          |  |
| 0.1682 ± 0.0012 ± 0.0037  | 30k   | MENDEZ              | 10 CLEO  | See BONVICINI 14                                   |
| 0.174 ± 0.012 ± 0.011   | 473   | <sup>1</sup> BISHAI | 97 CLEO  | $e^+ e^- \approx \Upsilon(4S)$                     |
| 0.137 ± 0.015 ± 0.016   | 264   | ANJOS               | 90C E691 | Photoproduction                                    |

<sup>1</sup> See BISHAI 97 for an isospin analysis of  $D^+ \rightarrow \bar{K} \pi$  amplitudes. $\Gamma(K_L^0 \pi^+)/\Gamma_{\text{total}}$  $\Gamma_{44}/\Gamma$ 

| VALUE (units $10^{-2}$ )     | EVTS      | DOCUMENT ID     | TECN    | COMMENT                   |
|------------------------------|-----------|-----------------|---------|---------------------------|
| <b>1.460 ± 0.040 ± 0.035</b> | 2023 ± 54 | <sup>1</sup> HE | 08 CLEO | $e^+ e^-$ at $\psi(3770)$ |

<sup>1</sup> The difference of CLEO  $D^+ \rightarrow K_S^0 \pi^+$  and  $K_L^0 \pi^+$  branching fractions over the sum (DOBBS 07 and HE 08) is  $+0.022 \pm 0.016 \pm 0.018$ . $\Gamma(K^- 2\pi^+)/\Gamma_{\text{total}}$  $\Gamma_{45}/\Gamma$ 

| VALUE (units $10^{-2}$ )  | EVTS  | DOCUMENT ID            | TECN     | COMMENT                             |
|---|-------|------------------------|----------|-------------------------------------|
| <b>9.38 ± 0.16 OUR FIT</b>  |       |                        |          | Error includes scale factor of 1.6. |
| <b>9.224 ± 0.059 ± 0.157</b>  |       | BONVICINI 14           | CLEO     | All CLEO-c runs                     |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |       |                        |          |                                     |
| 9.14 ± 0.10 ± 0.17  |       | <sup>1</sup> DOBBS     | 07 CLEO  | See BONVICINI 14                    |
| 9.5 ± 0.2 ± 0.3   | 15.1k | <sup>1</sup> HE        | 05 CLEO  | See DOBBS 07                        |
| 9.3 ± 0.6 ± 0.8   | 1502  | <sup>2</sup> BALEST    | 94 CLEO  | $e^+ e^- \approx \Upsilon(4S)$      |
| 6.4 <sup>+1.5</sup><br><sub>-1.4</sub>  |       | <sup>3</sup> BARLAG    | 92C ACCM | $\pi^-$ Cu 230 GeV                  |
| 9.1 ± 1.3 ± 0.4   | 1164  | ADLER                  | 88C MRK3 | $e^+ e^-$ 3.77 GeV                  |
| 9.1 ± 1.9   | 239   | <sup>4</sup> SCHINDLER | 81 MRK2  | $e^+ e^-$ 3.771 GeV                 |

<sup>1</sup> DOBBS 07 and HE 05 use single- and double-tagged events in an overall fit. DOBBS 07 supersedes HE 05.<sup>2</sup> BALEST 94 measures the ratio of  $D^+ \rightarrow K^- \pi^+ \pi^+$  and  $D^0 \rightarrow K^- \pi^+$  branching fractions to be  $2.35 \pm 0.16 \pm 0.16$  and uses their absolute measurement of the  $D^0 \rightarrow K^- \pi^+$  fraction (AKERIB 93).<sup>3</sup> BARLAG 92C computes the branching fraction by topological normalization.<sup>4</sup> SCHINDLER 81 (MARK-2) measures  $\sigma(e^+ e^- \rightarrow \psi(3770)) \times$  branching fraction to be  $0.38 \pm 0.05$  nb. We use the MARK-3 (ADLER 88C) value of  $\sigma = 4.2 \pm 0.6 \pm 0.3$  nb.

See the related review(s):

[Review of Multibody Charm Analyses](#) $\Gamma((K^- \pi^+)_{S\text{-wave}} \pi^+)/\Gamma(K^- 2\pi^+)$  $\Gamma_{46}/\Gamma_{45}$ This is the "fit fraction" from the Dalitz-plot analysis. The  $K^- \pi^+$  S-wave includes a broad scalar  $\kappa$  ( $\bar{K}_0^*(700)$ ), the  $\bar{K}_0^*(1430)^0$ , and non-resonant background.

| VALUE   | DOCUMENT ID            | TECN     | COMMENT                  |
|---|------------------------|----------|--------------------------|
| <b>0.801 ± 0.012 OUR AVERAGE</b>  |                        |          |                          |
| 0.8024 ± 0.0138 ± 0.0043  | <sup>1</sup> LINK      | 09 FOCS  | MIPWA fit, 53k evts      |
| 0.838 ± 0.038   | <sup>2</sup> BONVICINI | 08A CLEO | QMIPWA fit, 141k evts    |
| 0.786 ± 0.014 ± 0.018   | AITALA                 | 06 E791  | Dalitz fit, 15.1k events |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                        |          |                          |

0.8323±0.0150±0.0008      <sup>3</sup> LINK      07B FOCS    See LINK 09

<sup>1</sup> This LINK 09 model-independent partial-wave analysis of the  $K^- \pi^+$   $S$ -wave slices the  $K^- \pi^+$  mass range into 39 bins.

<sup>2</sup> The BONVICINI 08A QMIPWA (quasi-model-independent partial-wave analysis) of the  $K^- \pi^+$   $S$ -wave amplitude slices the  $K^- \pi^+$  mass range into 26 bins but keeps the Breit-Wigner  $\bar{K}_0^*(1430)^0$ .

<sup>3</sup> This LINK 07B fit uses a K matrix. The  $K^- \pi^+$   $S$ -wave fit fraction given above breaks down into (207.3 ± 25.5 ± 12.4)% isospin-1/2 and (40.5 ± 9.6 ± 3.2)% isospin-3/2 — with large interference between the two. The isospin-1/2 component includes the  $\kappa$  (or  $\bar{K}_0^*(700)^0$ ) and  $\bar{K}_0^*(1430)^0$ .

$\Gamma(\bar{K}_0^*(700)^0 \pi^+, \bar{K}_0^*(700) \rightarrow K^- \pi^+)/\Gamma(K^- 2\pi^+)$        $\Gamma_{47}/\Gamma_{45}$

This is the “fit fraction” from the Dalitz-plot analysis.

| VALUE             | DOCUMENT ID   | TECN    | COMMENT       |
|-------------------|---|---------|---------------|
| • • •             | We do not use the following data for averages, fits, limits, etc. | • • •   |               |
| 0.478±0.121±0.053 | AITALA  | 02 E791 | See AITALA 06 |

$\Gamma(\bar{K}_0^*(1430)^0 \pi^+, \bar{K}_0^*(1430)^0 \rightarrow K^- \pi^+)/\Gamma(K^- 2\pi^+)$        $\Gamma_{48}/\Gamma_{45}$

This is the “fit fraction” from the Dalitz-plot analysis.

| VALUE                | DOCUMENT ID   | TECN     | COMMENT                |
|----------------------|---|----------|------------------------|
| <b>0.1330±0.0062</b> | BONVICINI   | 08A CLEO | QMIPWA fit, 141k evts  |
| • • •                | We do not use the following data for averages, fits, limits, etc. | • • •    |                        |
| 0.125 ±0.014 ±0.005  | AITALA  | 02 E791  | See AITALA 06          |
| 0.284 ±0.022 ±0.059  | FRABETTI  | 94G E687 | Dalitz fit, 8800 evts  |
| 0.248 ±0.019 ±0.017  | ANJOS   | 93 E691  | $\gamma$ Be 90–260 GeV |

$\Gamma(\bar{K}^*(892)^0 \pi^+, \bar{K}^*(892)^0 \rightarrow K^- \pi^+)/\Gamma(K^- 2\pi^+)$        $\Gamma_{49}/\Gamma_{45}$

This is the “fit fraction” from the Dalitz-plot analysis.

| VALUE                           | DOCUMENT ID   | TECN     | COMMENT                  |
|---------------------------------|---|----------|--------------------------|
| <b>0.111 ±0.012 OUR AVERAGE</b> | Error includes scale factor of 3.7.                               |          |                          |
| 0.1236±0.0034±0.0034            | LINK  | 09 FOCS  | MIPWA fit, 53k evts      |
| 0.0988±0.0046                   | BONVICINI   | 08A CLEO | QMIPWA fit, 141k evts    |
| 0.119 ±0.002 ±0.020             | AITALA  | 06 E791  | Dalitz fit, 15.1k events |
| • • •                           | We do not use the following data for averages, fits, limits, etc. | • • •    |                          |
| 0.1361±0.0041±0.0030            | <sup>1</sup> LINK   | 07B FOCS | See LINK 09              |
| 0.123 ±0.010 ±0.009             | AITALA  | 02 E791  | See AITALA 06            |
| 0.137 ±0.006 ±0.009             | FRABETTI  | 94G E687 | Dalitz fit, 8800 evts    |
| 0.170 ±0.009 ±0.034             | ANJOS   | 93 E691  | $\gamma$ Be 90–260 GeV   |
| 0.14 ±0.04 ±0.04                | ALVAREZ   | 91B NA14 | Photoproduction          |
| 0.13 ±0.01 ±0.07                | ADLER   | 87 MRK3  | $e^+ e^-$ 3.77 GeV       |

<sup>1</sup> The statistical error on this LINK 07B value is corrected in LINK 09.

$\Gamma(\bar{K}^*(1410)^0 \pi^+, \bar{K}^{*0} \rightarrow K^- \pi^+)/\Gamma(K^- 2\pi^+)$        $\Gamma_{50}/\Gamma_{45}$


| VALUE (units 10 <sup>-3</sup> ) | DOCUMENT ID   | TECN     | COMMENT               |
|---------------------------------|---|----------|-----------------------|
| <b>not seen</b>                 | LINK  | 09 FOCS  | MIPWA fit, 53k evts   |
| <b>not seen</b>                 | BONVICINI   | 08A CLEO | QMIPWA fit, 141k evts |
| • • •                           | We do not use the following data for averages, fits, limits, etc. | • • •    |                       |
| 4.8±2.1±1.7                     | LINK  | 07B FOCS | See LINK 09           |

$\Gamma(\bar{K}_2^*(1430)^0 \pi^+, \bar{K}_2^*(1430)^0 \rightarrow K^- \pi^+)/\Gamma(K^- 2\pi^+)$   $\Gamma_{51}/\Gamma_{45}$

This is the "fit fraction" from the Dalitz-plot analysis.

| VALUE (units $10^{-2}$ )  | DOCUMENT ID   | TECN | COMMENT                       |
|---|---|------|-------------------------------|
| <b>0.24 ± 0.08 OUR AVERAGE</b>  | Error includes scale factor of 2.2. See the ideogram below. |      |                               |
| 0.58 ± 0.10 ± 0.06  | LINK  | 09   | FOCS MIPWA fit, 53k evts      |
| 0.204 ± 0.040   | BONVICINI   | 08A  | CLEO QMIPWA fit, 141k evts    |
| 0.2 ± 0.1 ± 0.1   | AITALA  | 06   | E791 Dalitz fit, 15.1k events |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● |   |      |                               |
| 0.39 ± 0.09 ± 0.05  | LINK  | 07B  | FOCS See LINK 09              |
| 0.5 ± 0.1 ± 0.2   | AITALA  | 02   | E791 See AITALA 06            |

WEIGHTED AVERAGE  
0.24±0.08 (Error scaled by 2.2)



$\Gamma(\bar{K}_2^*(1430)^0 \pi^+, \bar{K}_2^*(1430)^0 \rightarrow K^- \pi^+)/\Gamma(K^- 2\pi^+)$   $\Gamma_{51}/\Gamma_{45}$   
(units  $10^{-2}$ )

$\Gamma(\bar{K}^*(1680)^0 \pi^+, \bar{K}^*(1680)^0 \rightarrow K^- \pi^+)/\Gamma(K^- 2\pi^+)$   $\Gamma_{52}/\Gamma_{45}$

This is the "fit fraction" from the Dalitz-plot analysis.

| VALUE (units $10^{-2}$ )  | DOCUMENT ID | TECN | COMMENT                       |
|---|-------------|------|-------------------------------|
| <b>0.23 ± 0.12 OUR AVERAGE</b>  |             |      |                               |
| 1.75 ± 0.62 ± 0.54  | LINK        | 09   | FOCS MIPWA fit, 53k evts      |
| 0.196 ± 0.118   | BONVICINI   | 08A  | CLEO QMIPWA fit, 141k evts    |
| 1.2 ± 0.6 ± 1.2   | AITALA      | 06   | E791 Dalitz fit, 15.1k events |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● |             |      |                               |
| 1.90 ± 0.63 ± 0.43  | LINK        | 07B  | FOCS See LINK 09              |
| 2.5 ± 0.7 ± 0.3   | AITALA      | 02   | E791 See AITALA 06            |
| 4.7 ± 0.6 ± 0.7   | FRABETTI    | 94G  | E687 Dalitz fit, 8800 evts    |
| 3.0 ± 0.4 ± 1.3   | ANJOS       | 93   | E691 $\gamma$ Be 90–260 GeV   |

$\Gamma(K^-(2\pi^+)_{I=2})/\Gamma(K^-2\pi^+)$   $\Gamma_{53}/\Gamma_{45}$

| VALUE              | DOCUMENT ID   | TECN | COMMENT               |
|--------------------|---------------|------|-----------------------|
| <b>0.155±0.028</b> | BONVICINI 08A | CLEO | QMIPWA fit, 141k evts |

$\Gamma(K^-2\pi^+ \text{ nonresonant})/\Gamma(K^-2\pi^+)$   $\Gamma_{54}/\Gamma_{45}$

This is the “fit fraction” from the Dalitz-plot analysis. Later analyses find little need for this decay mode.

| VALUE   | DOCUMENT ID  | TECN | COMMENT                |
|---|--------------|------|------------------------|
| • • • We do not use the following data for averages, fits, limits, etc. • • • |              |      |                        |
| 0.130±0.058±0.044   | AITALA 02    | E791 | See AITALA 06          |
| 0.998±0.037±0.072   | FRABETTI 94G | E687 | Dalitz fit, 8800 evts  |
| 0.838±0.088±0.275   | ANJOS 93     | E691 | $\gamma$ Be 90–260 GeV |
| 0.79 ±0.07 ±0.15  | ADLER 87     | MRK3 | $e^+e^-$ 3.77 GeV      |

$\Gamma(K_S^0\pi^+\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{55}/\Gamma$

| VALUE (units $10^{-2}$ )  | EVTS | DOCUMENT ID           | TECN | COMMENT           |
|---|------|-----------------------|------|-------------------|
| • • • We do not use the following data for averages, fits, limits, etc. • • • |      |                       |      |                   |
| 6.99±0.09±0.25  |      | <sup>1</sup> DOBBS 07 | CLEO | See BONVICINI 14  |
| 7.2 ±0.2 ±0.4   | 5.1k | <sup>1</sup> HE 05    | CLEO | See DOBBS 07      |
| 5.1 ±1.3 ±0.8   | 159  | ADLER 88C             | MRK3 | $e^+e^-$ 3.77 GeV |

<sup>1</sup> DOBBS 07 and HE 05 use single- and double-tagged events in an overall fit. DOBBS 07 supersedes HE 05.

$\Gamma(K_S^0\rho^+\pi^0)/\Gamma(K^-2\pi^+)$   $\Gamma_{55}/\Gamma_{45}$

| VALUE                    | DOCUMENT ID  | TECN | COMMENT         |
|--------------------------|--------------|------|-----------------|
| <b>0.785±0.007±0.016</b> | BONVICINI 14 | CLEO | All CLEO-c runs |

$\Gamma(K_S^0\rho^+)/\Gamma(K_S^0\pi^+\pi^0)$   $\Gamma_{56}/\Gamma_{55}$

This is the “fit fraction” from the Dalitz-plot analysis.

| VALUE (units $10^{-2}$ )                      | DOCUMENT ID              | TECN | COMMENT                  |
|---|--------------------------|------|--------------------------|
| <b>83.4±2.2<sup>+7.1</sup><sub>-3.6</sub></b> | <sup>1</sup> ABLIKIM 14E | BES3 | $e^+e^-$ at $\psi(3770)$ |

<sup>1</sup> Fit fraction from Dalitz plot analysis of 142k  $D^+ \rightarrow K_S^0\pi^+\pi^0$  events.

$\Gamma(K_S^0\rho(1450)^+, \rho^+ \rightarrow \pi^+\pi^0)/\Gamma(K_S^0\pi^+\pi^0)$   $\Gamma_{57}/\Gamma_{55}$

This is the “fit fraction” from the Dalitz-plot analysis.

| VALUE (%)                                    | DOCUMENT ID | TECN | COMMENT                  |
|--|-------------|------|--------------------------|
| <b>2.1±0.3<sup>+1.6</sup><sub>-1.9</sub></b> | ABLIKIM 14E | BES3 | $e^+e^-$ at $\psi(3770)$ |

$\Gamma(\bar{K}^*(892)^0\pi^+, \bar{K}^*(892)^0 \rightarrow K_S^0\pi^0)/\Gamma(K_S^0\pi^+\pi^0)$   $\Gamma_{58}/\Gamma_{55}$

This is the “fit fraction” from the Dalitz-plot analysis.

| VALUE (units $10^{-2}$ )                         | DOCUMENT ID              | TECN | COMMENT                  |
|--|--------------------------|------|--------------------------|
| <b>3.58±0.17<sup>+0.39</sup><sub>-0.38</sub></b> | <sup>1</sup> ABLIKIM 14E | BES3 | $e^+e^-$ at $\psi(3770)$ |

<sup>1</sup> Fit fraction from Dalitz plot analysis of 142k  $D^+ \rightarrow K_S^0\pi^+\pi^0$  events.

$\Gamma(\bar{K}_0^*(1430)^0 \pi^+, \bar{K}_0^{*0} \rightarrow K_S^0 \pi^0) / \Gamma(K_S^0 \pi^+ \pi^0)$   $\Gamma_{59} / \Gamma_{55}$

This is the "fit fraction" from the Dalitz-plot analysis.

| VALUE (%)             | DOCUMENT ID | TECN | COMMENT                        |
|-----------------------|-------------|------|--------------------------------|
| $3.7 \pm 0.6 \pm 1.1$ | ABLIKIM     | 14E  | BES3 $e^+ e^-$ at $\psi(3770)$ |

$\Gamma(\bar{K}_0^*(1680)^0 \pi^+, \bar{K}_0^{*0} \rightarrow K_S^0 \pi^0) / \Gamma(K_S^0 \pi^+ \pi^0)$   $\Gamma_{60} / \Gamma_{55}$

This is the "fit fraction" from the Dalitz-plot analysis.

| VALUE (%)                   | DOCUMENT ID | TECN | COMMENT                        |
|-----------------------------|-------------|------|--------------------------------|
| $1.3 \pm 0.2^{+0.9}_{-1.3}$ | ABLIKIM     | 14E  | BES3 $e^+ e^-$ at $\psi(3770)$ |

$\Gamma(\bar{\kappa}^0 \pi^+, \bar{\kappa}^0 \rightarrow K_S^0 \pi^0) / \Gamma(K_S^0 \pi^+ \pi^0)$   $\Gamma_{61} / \Gamma_{55}$

This is the "fit fraction" from the Dalitz-plot analysis.

| VALUE (%)                   | DOCUMENT ID | TECN | COMMENT                        |
|-----------------------------|-------------|------|--------------------------------|
| $7.7 \pm 1.2^{+6.5}_{-4.8}$ | ABLIKIM     | 14E  | BES3 $e^+ e^-$ at $\psi(3770)$ |

$\Gamma(K_S^0 \pi^+ \pi^0 \text{ nonresonant}) / \Gamma(K_S^0 \pi^+ \pi^0)$   $\Gamma_{62} / \Gamma_{55}$

This is the "fit fraction" from the Dalitz-plot analysis.

| VALUE (units $10^{-2}$ )    | DOCUMENT ID          | TECN | COMMENT                        |
|-----------------------------|----------------------|------|--------------------------------|
| $4.6 \pm 0.7^{+5.4}_{-5.1}$ | <sup>1</sup> ABLIKIM | 14E  | BES3 $e^+ e^-$ at $\psi(3770)$ |

<sup>1</sup> Fit fraction from Dalitz plot analysis of 142k  $D^+ \rightarrow K_S^0 \pi^+ \pi^0$  events.

$\Gamma(K_S^0 \pi^+ \pi^0 \text{ nonresonant and } \bar{\kappa}^0 \pi^+) / \Gamma(K_S^0 \pi^+ \pi^0)$   $\Gamma_{63} / \Gamma_{55}$

This is the "fit fraction" from the Dalitz-plot analysis.

| VALUE (%)                    | DOCUMENT ID | TECN | COMMENT                        |
|------------------------------|-------------|------|--------------------------------|
| $18.6 \pm 1.7^{+2.3}_{-4.6}$ | ABLIKIM     | 14E  | BES3 $e^+ e^-$ at $\psi(3770)$ |

$\Gamma((K_S^0 \pi^0)_{S\text{-wave}} \pi^+) / \Gamma(K_S^0 \pi^+ \pi^0)$   $\Gamma_{64} / \Gamma_{55}$

The numerator here is the coherent sum of the  $\bar{K}_0^*(1430)^0 \pi^+$ ,  $\bar{\kappa}^0 \pi^+$ , and nonresonant contributions.

| VALUE (%)                    | DOCUMENT ID | TECN | COMMENT                        |
|------------------------------|-------------|------|--------------------------------|
| $17.3 \pm 1.4^{+3.4}_{-4.3}$ | ABLIKIM     | 14E  | BES3 $e^+ e^-$ at $\psi(3770)$ |

$\Gamma(K_S^0 \pi^+ \eta'(958)) / \Gamma_{\text{total}}$   $\Gamma_{65} / \Gamma$

| VALUE (units $10^{-3}$ ) | EVTS | DOCUMENT ID | TECN      | COMMENT              |
|--------------------------|------|-------------|-----------|----------------------|
| $1.90 \pm 0.17 \pm 0.13$ | 267  | ABLIKIM     | 18AC BES3 | $e^+ e^-$ , 3773 MeV |

$\Gamma(K^- 2\pi^+ \pi^0) / \Gamma_{\text{total}}$   $\Gamma_{66} / \Gamma$

See our 2008 Review (Physics Letters **B667** 1 (2008)) for measurements of submodes of this mode. There is nothing new since 1992, and the two papers, ANJOS 92C, with  $91 \pm 12$  events above background, and COFFMAN 92B, with  $142 \pm 20$  such events, could not determine submode fractions with much accuracy.

| VALUE (units $10^{-2}$ )    | EVTS | DOCUMENT ID        | TECN | COMMENT                 |
|-----------------------------|------|--------------------|------|-------------------------|
| $5.98 \pm 0.08 \pm 0.16$    |      | <sup>1</sup> DOBBS | 07   | CLEO See BONVICINI 14   |
| $6.0 \pm 0.2 \pm 0.2$       | 4.8k | <sup>1</sup> HE    | 05   | CLEO See DOBBS 07       |
| $5.8 \pm 1.2 \pm 1.2$       | 142  | COFFMAN            | 92B  | MRK3 $e^+ e^-$ 3.77 GeV |
| $6.3^{+1.4}_{-1.3} \pm 1.2$ | 175  | BALTRUSAIT..       | 86E  | MRK3 See COFFMAN 92B    |

<sup>1</sup> DOBBS 07 and HE 05 use single- and double-tagged events in an overall fit. DOBBS 07 supersedes HE 05.

$\Gamma(K^- 2\pi^+ \pi^0)/\Gamma(K^- 2\pi^+)$   $\Gamma_{66}/\Gamma_{45}$

| VALUE                        | DOCUMENT ID  | TECN | COMMENT         |
|------------------------------|--------------|------|-----------------|
| <b>0.666 ± 0.006 ± 0.014</b> | BONVICINI 14 | CLEO | All CLEO-c runs |

$\Gamma(K_S^0 2\pi^+ \pi^-)/\Gamma_{\text{total}}$   $\Gamma_{67}/\Gamma$

See our 2008 Review (Physics Letters **B667** 1 (2008)) for measurements of submodes of this mode. There is nothing new since 1992, and the two papers, ANJOS 92C, with  $229 \pm 17$  events above background, and COFFMAN 92B, with  $209 \pm 20$  such events, could not determine submode fractions with much accuracy.

| VALUE (units $10^{-2}$ ) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

|  |      |                     |     |      |                    |
|--|------|---------------------|-----|------|--------------------|
| $3.122 \pm 0.046 \pm 0.096$                              |      | <sup>1</sup> DOBBS  | 07  | CLEO | See BONVICINI 14   |
| $3.2 \pm 0.1 \pm 0.2$                                    | 3.2k | <sup>1</sup> HE     | 05  | CLEO | See DOBBS 07       |
| $2.1 \begin{smallmatrix} +1.0 \\ -0.9 \end{smallmatrix}$ |      | <sup>2</sup> BARLAG | 92C | ACCM | $\pi^-$ Cu 230 GeV |
| $3.3 \pm 0.8 \pm 0.2$                                    | 168  | ADLER               | 88C | MRK3 | $e^+ e^-$ 3.77 GeV |

<sup>1</sup> DOBBS 07 and HE 05 use single- and double-tagged events in an overall fit. DOBBS 07 supersedes HE 05.

<sup>2</sup> BARLAG 92C computes the branching fraction by topological normalization.

$\Gamma(K_S^0 2\pi^+ \pi^-)/\Gamma(K^- 2\pi^+)$   $\Gamma_{67}/\Gamma_{45}$

| VALUE                        | DOCUMENT ID  | TECN | COMMENT         |
|------------------------------|--------------|------|-----------------|
| <b>0.331 ± 0.004 ± 0.006</b> | BONVICINI 14 | CLEO | All CLEO-c runs |

$\Gamma(K^- 3\pi^+ \pi^-)/\Gamma(K^- 2\pi^+)$   $\Gamma_{68}/\Gamma_{45}$

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------|------|-------------|------|---------|
|-------|------|-------------|------|---------|

**0.061 ± 0.005 OUR FIT** Error includes scale factor of 1.1.

**0.062 ± 0.008 OUR AVERAGE** Error includes scale factor of 1.3.

|                             |      |          |     |      |   |
|-----------------------------|------|----------|-----|------|---|
| $0.058 \pm 0.002 \pm 0.006$ | 2923 | LINK     | 03D | FOCS | $\gamma$ A, $\bar{E}_\gamma \approx 180$ GeV  |
| $0.077 \pm 0.008 \pm 0.010$ | 239  | FRABETTI | 97C | E687 | $\gamma$ Be, $\bar{E}_\gamma \approx 200$ GeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                          |     |       |     |      |                 |
|--------------------------|-----|-------|-----|------|-----------------|
| $0.09 \pm 0.01 \pm 0.01$ | 113 | ANJOS | 90D | E691 | Photoproduction |
|--------------------------|-----|-------|-----|------|-----------------|

$\Gamma(\bar{K}^*(892)^0 2\pi^+ \pi^-, \bar{K}^*(892)^0 \rightarrow K^- \pi^+)/\Gamma(K^- 3\pi^+ \pi^-)$   $\Gamma_{69}/\Gamma_{68}$

| VALUE                     | DOCUMENT ID | TECN | COMMENT   |
|---------------------------|-------------|------|---|
| <b>0.21 ± 0.04 ± 0.06</b> | LINK        | 03D  | FOCS $\gamma$ A, $\bar{E}_\gamma \approx 180$ GeV |

$\Gamma(\bar{K}^*(892)^0 \rho^0 \pi^+, \bar{K}^*(892)^0 \rightarrow K^- \pi^+)/\Gamma(K^- 2\pi^+)$   $\Gamma_{70}/\Gamma_{45}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                             |          |     |  |
|-----------------------------|----------|-----|--|
| $0.016 \pm 0.007 \pm 0.004$ | FRABETTI | 97C | E687 $\gamma$ Be, $\bar{E}_\gamma \approx 200$ GeV |
|-----------------------------|----------|-----|--|

$\Gamma(\bar{K}^*(892)^0 \rho^0 \pi^+, \bar{K}^*(892)^0 \rightarrow K^- \pi^+)/\Gamma(K^- 3\pi^+ \pi^-)$   $\Gamma_{70}/\Gamma_{68}$

| VALUE                     | DOCUMENT ID | TECN | COMMENT   |
|---------------------------|-------------|------|---|
| <b>0.40 ± 0.03 ± 0.06</b> | LINK        | 03D  | FOCS $\gamma$ A, $\bar{E}_\gamma \approx 180$ GeV |

$\Gamma(\bar{K}^*(892)^0 a_1(1260)^+)/\Gamma(K^- 2\pi^+)$   $\Gamma_{71}/\Gamma_{45}$

Unseen decay modes of the  $\bar{K}^*(892)^0$  and  $a_1(1260)^+$  are included.

| VALUE                    | DOCUMENT ID | TECN     | COMMENT                                      |
|--------------------------|-------------|----------|--|
| <b>0.099±0.008±0.018</b> | LINK        | 03D FOCS | $\gamma$ A, $\bar{E}_\gamma \approx 180$ GeV |

$\Gamma(\bar{K}^*(892)^0 2\pi^+ \pi^- \text{no-}\rho, \bar{K}^*(892)^0 \rightarrow K^- \pi^+)/\Gamma(K^- 2\pi^+)$   $\Gamma_{72}/\Gamma_{45}$

| VALUE                    | DOCUMENT ID | TECN     | COMMENT                                       |
|--------------------------|-------------|----------|---|
| <b>0.032±0.010±0.008</b> | FRABETTI    | 97C E687 | $\gamma$ Be, $\bar{E}_\gamma \approx 200$ GeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\Gamma(K^- \rho^0 2\pi^+)/\Gamma(K^- 2\pi^+)$   $\Gamma_{73}/\Gamma_{45}$

| VALUE                    | DOCUMENT ID | TECN     | COMMENT                                       |
|--------------------------|-------------|----------|---|
| <b>0.034±0.009±0.005</b> | FRABETTI    | 97C E687 | $\gamma$ Be, $\bar{E}_\gamma \approx 200$ GeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\Gamma(K^- \rho^0 2\pi^+)/\Gamma(K^- 3\pi^+ \pi^-)$   $\Gamma_{73}/\Gamma_{68}$

| VALUE                 | DOCUMENT ID | TECN     | COMMENT                                      |
|-----------------------|-------------|----------|--|
| <b>0.30±0.04±0.01</b> | LINK        | 03D FOCS | $\gamma$ A, $\bar{E}_\gamma \approx 180$ GeV |

$\Gamma(K^- 3\pi^+ \pi^- \text{nonresonant})/\Gamma(K^- 3\pi^+ \pi^-)$   $\Gamma_{74}/\Gamma_{68}$

| VALUE                  | CL% | DOCUMENT ID | TECN     | COMMENT                                       |
|------------------------|-----|-------------|----------|---|
| <b>0.07 ±0.05±0.01</b> |     | LINK        | 03D FOCS | $\gamma$ A, $\bar{E}_\gamma \approx 180$ GeV  |
| <0.026                 | 90  | FRABETTI    | 97C E687 | $\gamma$ Be, $\bar{E}_\gamma \approx 200$ GeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\Gamma(K^+ 2K_S^0)/\Gamma_{\text{total}}$   $\Gamma_{75}/\Gamma$

| VALUE (units $10^{-4}$ ) | EVTS | DOCUMENT ID | TECN     | COMMENT                          |
|--------------------------|------|-------------|----------|----------------------------------|
| <b>25.4±0.5±1.2</b>      | 3551 | ABLIKIM     | 17A BES3 | $e^+ e^- \rightarrow \psi(3770)$ |

$\Gamma(K^+ 2K_S^0)/\Gamma(K^- 2\pi^+)$   $\Gamma_{75}/\Gamma_{45}$

| VALUE                    | EVTS    | DOCUMENT ID | TECN    | COMMENT                    |
|--------------------------|---------|-------------|---------|----------------------------|
| <b>0.035±0.010±0.005</b> | 39 ± 9  | ALBRECHT    | 94i ARG | $e^+ e^- \approx 10$ GeV   |
| <b>0.085±0.018</b>       | 70 ± 12 | AMMAR       | 91 CLEO | $e^+ e^- \approx 10.5$ GeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\Gamma(K^+ K^- K_S^0 \pi^+)/\Gamma(K_S^0 2\pi^+ \pi^-)$   $\Gamma_{76}/\Gamma_{67}$

| VALUE (units $10^{-3}$ ) | EVTS   | DOCUMENT ID | TECN     | COMMENT  |
|--------------------------|--------|-------------|----------|--|
| <b>7.7±1.5±0.9</b>       | 35 ± 7 | LINK        | 01C FOCS | $\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV |

————— Pionic modes —————

$\Gamma(\pi^+ \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{77}/\Gamma$

| VALUE (units $10^{-3}$ )   | EVTS | DOCUMENT ID | TECN     | COMMENT              |
|----------------------------|------|-------------|----------|----------------------|
| <b>1.247±0.033 OUR FIT</b> |      |             |          |                      |
| <b>1.259±0.033±0.023</b>   | 10k  | ABLIKIM     | 18w BES3 | $e^+ e^-$ , 3773 MeV |



$\Gamma(\pi^+\pi^0)/\Gamma(K^-2\pi^+)$

$\Gamma_{77}/\Gamma_{45}$

VALUE (units  $10^{-2}$ )      EVTS      DOCUMENT ID      TECN      COMMENT

**1.33±0.04 OUR FIT** Error includes scale factor of 1.1.

**1.31±0.06 OUR AVERAGE**

|                |           |          |     |      |                               |
|----------------|-----------|----------|-----|------|-------------------------------|
| 1.29±0.04±0.05 | 2649 ± 76 | MENDEZ   | 10  | CLEO | $e^+e^-$ at 3774 MeV          |
| 1.33±0.11±0.09 | 1229 ± 99 | AUBERT,B | 06F | BABR | $e^+e^- \approx \Upsilon(4S)$ |
| 1.44±0.19±0.10 | 171 ± 22  | ARMS     | 04  | CLEO | $e^+e^- \approx 10$ GeV       |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                |          |       |    |      |               |
|----------------|----------|-------|----|------|---------------|
| 1.33±0.07±0.06 | 914 ± 46 | RUBIN | 06 | CLEO | See MENDEZ 10 |
|----------------|----------|-------|----|------|---------------|

$\Gamma(2\pi^+\pi^-)/\Gamma(K^-2\pi^+)$


$\Gamma_{78}/\Gamma_{45}$

VALUE (units  $10^{-2}$ )      EVTS      DOCUMENT ID      TECN      COMMENT

**3.48±0.19 OUR AVERAGE** Error includes scale factor of 1.4. See the ideogram below.

|   |           |          |     |      |                               |
|---|-----------|----------|-----|------|-------------------------------|
| 3.52±0.11±0.12                              | 3303 ± 95 | RUBIN    | 06  | CLEO | $e^+e^-$ at $\psi(3770)$      |
| 4.1 ± 1.1 ± 0.3                             | 85 ± 22   | ABLIKIM  | 05F | BES  | $e^+e^- \approx \psi(3770)$   |
| 3.11±0.18 <sup>+0.16</sup> <sub>-0.26</sub> | 1172      | AITALA   | 01B | E791 | $\pi^-$ nucleus, 500 GeV      |
| 4.3 ± 0.3 ± 0.3                             | 236       | FRABETTI | 97D | E687 | $\gamma$ Be $\approx 200$ GeV |
| 3.5 ± 0.7 ± 0.3                             | 83        | ANJOS    | 89  | E691 | Photoproduction               |

WEIGHTED AVERAGE  
3.48±0.19 (Error scaled by 1.4)




### $\Gamma(\rho^0 \pi^+)/\Gamma(2\pi^+ \pi^-)$

$\Gamma_{79}/\Gamma_{78}$

This is the "fit fraction" from the Dalitz-plot analysis.

| VALUE                          | DOCUMENT ID  | TECN | COMMENT   |
|--------------------------------|--------------|------|---|
| <b>0.25 ± 0.04 OUR AVERAGE</b> |              |      | Error includes scale factor of 2.4. See the ideogram below. |
| 0.200 ± 0.023 ± 0.009          | BONVICINI 07 | CLEO | Dalitz fit, ≈ 2240 evts                                     |
| 0.3082 ± 0.0314 ± 0.0230       | LINK 04      | FOCS | Dalitz fit, 1527 ± 51 evts                                  |
| 0.336 ± 0.032 ± 0.022          | AITALA 01B   | E791 | Dalitz fit, 1172 evts                                       |



### $\Gamma(\pi^+(\pi^+ \pi^-)_{S\text{-wave}})/\Gamma(2\pi^+ \pi^-)$

$\Gamma_{80}/\Gamma_{78}$

This is the "fit fraction" from the Dalitz-plot analysis. See also the next three data blocks.

| VALUE                           | DOCUMENT ID          | TECN | COMMENT                    |
|---------------------------------|----------------------|------|----------------------------|
| <b>0.5600 ± 0.0324 ± 0.0214</b> | <sup>1</sup> LINK 04 | FOCS | Dalitz fit, 1527 ± 51 evts |

<sup>1</sup> LINK 04 borrows a K-matrix parametrization from ANISOVICH 03 of the full  $\pi\text{-}\pi$  S-wave isoscalar scattering amplitude to describe the  $\pi^+ \pi^-$  S-wave component of the  $\pi^+ \pi^+ \pi^-$  state. The fit fraction given above is a sum over five  $f_0$  mesons, the  $f_0(980)$ ,  $f_0(1300)$ ,  $f_0(1200\text{-}1600)$ ,  $f_0(1500)$ , and  $f_0(1750)$ . See LINK 04 for details and discussion.

### $\Gamma(\sigma \pi^+, \sigma \rightarrow \pi^+ \pi^-)/\Gamma(2\pi^+ \pi^-)$

$\Gamma_{81}/\Gamma_{78}$

This is the "fit fraction" from the Dalitz-plot analysis.

| VALUE                            | DOCUMENT ID  | TECN | COMMENT                 |
|----------------------------------|--------------|------|-------------------------|
| <b>0.422 ± 0.027 OUR AVERAGE</b> |              |      |                         |
| 0.418 ± 0.014 ± 0.025            | BONVICINI 07 | CLEO | Dalitz fit, ≈ 2240 evts |
| 0.463 ± 0.090 ± 0.021            | AITALA 01B   | E791 | Dalitz fit, 1172 evts   |

$\Gamma(f_0(980)\pi^+, f_0(980) \rightarrow \pi^+\pi^-)/\Gamma(2\pi^+\pi^-)$   $\Gamma_{82}/\Gamma_{78}$

This is the "fit fraction" from the Dalitz-plot analysis.

| VALUE                          | DOCUMENT ID                         | TECN | COMMENT                      |
|--------------------------------|-------------------------------------|------|------------------------------|
| <b>0.048±0.010 OUR AVERAGE</b> | Error includes scale factor of 1.3. |      |                              |
| 0.041±0.009±0.003              | BONVICINI                           | 07   | CLEO Dalitz fit, ≈ 2240 evts |
| 0.062±0.013±0.004              | AITALA                              | 01B  | E791 Dalitz fit, 1172 evts   |

$\Gamma(f_0(1370)\pi^+, f_0(1370) \rightarrow \pi^+\pi^-)/\Gamma(2\pi^+\pi^-)$   $\Gamma_{83}/\Gamma_{78}$


This is the "fit fraction" from the Dalitz-plot analysis.

| VALUE                          | DOCUMENT ID | TECN | COMMENT                      |
|--------------------------------|-------------|------|------------------------------|
| <b>0.024±0.013 OUR AVERAGE</b> |             |      |                              |
| 0.026±0.018±0.006              | BONVICINI   | 07   | CLEO Dalitz fit, ≈ 2240 evts |
| 0.023±0.015±0.008              | AITALA      | 01B  | E791 Dalitz fit, 1172 evts   |

$\Gamma(f_2(1270)\pi^+, f_2(1270) \rightarrow \pi^+\pi^-)/\Gamma(2\pi^+\pi^-)$   $\Gamma_{84}/\Gamma_{78}$

This is the "fit fraction" from the Dalitz-plot analysis.

| VALUE                           | DOCUMENT ID   | TECN | COMMENT                         |
|---------------------------------|---|------|---------------------------------|
| <b>0.154 ±0.025 OUR AVERAGE</b> | Error includes scale factor of 1.9. See the ideogram below. |      |                                 |
| 0.182 ±0.026 ±0.007             | BONVICINI   | 07   | CLEO Dalitz fit, ≈ 2240 evts    |
| 0.1174±0.0190±0.0029            | LINK  | 04   | FOCS Dalitz fit, 1527 ± 51 evts |
| 0.194 ±0.025 ±0.004             | AITALA  | 01B  | E791 Dalitz fit, 1172 evts      |



$\Gamma(\rho(1450)^0\pi^+, \rho(1450)^0 \rightarrow \pi^+\pi^-)/\Gamma(2\pi^+\pi^-)$   $\Gamma_{85}/\Gamma_{78}$

This is the "fit fraction" from the Dalitz-plot analysis.

| VALUE   | CL% | DOCUMENT ID  | TECN | COMMENT                         |
|---|-----|--------------|------|---------------------------------|
| <b>&lt;0.024</b>  | 95  | BONVICINI 07 | CLEO | Dalitz fit, $\approx 2240$ evts |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |     |              |      |                                 |
| $0.007 \pm 0.007 \pm 0.003$   |     | AITALA 01B   | E791 | Dalitz fit, 1172 evts           |

$\Gamma(f_0(1500)\pi^+, f_0(1500) \rightarrow \pi^+\pi^-)/\Gamma(2\pi^+\pi^-)$   $\Gamma_{86}/\Gamma_{78}$

This is the "fit fraction" from the Dalitz-plot analysis.

| VALUE  | DOCUMENT ID  | TECN | COMMENT                         |
|--|--------------|------|---------------------------------|
| <b>0.034 <math>\pm</math> 0.010 <math>\pm</math> 0.008</b> | BONVICINI 07 | CLEO | Dalitz fit, $\approx 2240$ evts |

$\Gamma(f_0(1710)\pi^+, f_0(1710) \rightarrow \pi^+\pi^-)/\Gamma(2\pi^+\pi^-)$   $\Gamma_{87}/\Gamma_{78}$

This is the "fit fraction" from the Dalitz-plot analysis.

| VALUE            | CL% | DOCUMENT ID  | TECN | COMMENT                         |
|------------------|-----|--------------|------|---------------------------------|
| <b>&lt;0.016</b> | 95  | BONVICINI 07 | CLEO | Dalitz fit, $\approx 2240$ evts |

$\Gamma(f_0(1790)\pi^+, f_0(1790) \rightarrow \pi^+\pi^-)/\Gamma(2\pi^+\pi^-)$   $\Gamma_{88}/\Gamma_{78}$

This is the "fit fraction" from the Dalitz-plot analysis.

| VALUE           | CL% | DOCUMENT ID  | TECN | COMMENT                         |
|-----------------|-----|--------------|------|---------------------------------|
| <b>&lt;0.02</b> | 95  | BONVICINI 07 | CLEO | Dalitz fit, $\approx 2240$ evts |

$\Gamma((\pi^+\pi^+)_{S\text{-wave}}\pi^-)/\Gamma(2\pi^+\pi^-)$   $\Gamma_{89}/\Gamma_{78}$

This is the "fit fraction" from the Dalitz-plot analysis.

| VALUE            | CL% | DOCUMENT ID  | TECN | COMMENT                         |
|------------------|-----|--------------|------|---------------------------------|
| <b>&lt;0.037</b> | 95  | BONVICINI 07 | CLEO | Dalitz fit, $\approx 2240$ evts |

$\Gamma(2\pi^+\pi^- \text{ nonresonant})/\Gamma(2\pi^+\pi^-)$   $\Gamma_{90}/\Gamma_{78}$

This is the "fit fraction" from the Dalitz-plot analysis.

| VALUE   | CL% | DOCUMENT ID  | TECN | COMMENT                         |
|---|-----|--------------|------|---------------------------------|
| <b>&lt;0.035</b>  | 95  | BONVICINI 07 | CLEO | Dalitz fit, $\approx 2240$ evts |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |     |              |      |                                 |
| $0.078 \pm 0.060 \pm 0.027$   |     | AITALA 01B   | E791 | Dalitz fit, 1172 evts           |

$\Gamma(\pi^+2\pi^0)/\Gamma(K^-2\pi^+)$   $\Gamma_{91}/\Gamma_{45}$

| VALUE (units $10^{-2}$ )                             | EVTS          | DOCUMENT ID | TECN | COMMENT                  |
|--|---------------|-------------|------|--------------------------|
| <b>5.0 <math>\pm</math> 0.3 <math>\pm</math> 0.3</b> | $1535 \pm 89$ | RUBIN 06    | CLEO | $e^+e^-$ at $\psi(3770)$ |

$\Gamma(2\pi^+\pi^-\pi^0)/\Gamma(K^-2\pi^+)$   $\Gamma_{92}/\Gamma_{45}$

| VALUE (units $10^{-2}$ )                              | EVTS           | DOCUMENT ID | TECN | COMMENT                  |
|---|----------------|-------------|------|--------------------------|
| <b>12.4 <math>\pm</math> 0.5 <math>\pm</math> 0.6</b> | $5701 \pm 205$ | RUBIN 06    | CLEO | $e^+e^-$ at $\psi(3770)$ |

$\Gamma(3\pi^+2\pi^-)/\Gamma(K^-2\pi^+)$   $\Gamma_{93}/\Gamma_{45}$

| VALUE (units $10^{-2}$ )  | EVTS         | DOCUMENT ID  | TECN | COMMENT   |
|---|--------------|--------------|------|---|
| <b>1.77 <math>\pm</math> 0.17 OUR FIT</b>                                     |              |              |      |   |
| <b>1.73 <math>\pm</math> 0.20 <math>\pm</math> 0.17</b>                       | $732 \pm 77$ | RUBIN 06     | CLEO | $e^+e^-$ at $\psi(3770)$                          |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |              |              |      |   |
| $2.3 \pm 0.4 \pm 0.2$   | 58           | FRABETTI 97C | E687 | $\gamma\text{Be}, \bar{E}_\gamma \approx 200$ GeV |

$\Gamma(3\pi^+2\pi^-)/\Gamma(K^-3\pi^+\pi^-)$   $\Gamma_{93}/\Gamma_{68}$

| VALUE                      | EVTS | DOCUMENT ID | TECN     | COMMENT                                      |
|----------------------------|------|-------------|----------|--|
| <b>0.289±0.019 OUR FIT</b> |      |             |          |  |
| <b>0.290±0.017±0.011</b>   | 835  | LINK        | 03D FOCS | $\gamma$ A, $\bar{E}_\gamma \approx 180$ GeV |

$\Gamma(\eta\pi^+)/\Gamma_{\text{total}}$   $\Gamma_{94}/\Gamma$

Unseen decay modes of the  $\eta$  are included.

| VALUE (units $10^{-4}$ )  | EVTS      | DOCUMENT ID | TECN     | COMMENT              |
|---|-----------|-------------|----------|----------------------|
| <b>37.7 ±0.9 OUR FIT</b>  |           |             |          |                      |
| <b>37.90±0.70±0.68</b>  | 12k       | ABLIKIM     | 18W BES3 | $e^+e^-$ , 3773 MeV  |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |           |             |          |                      |
| 30.7 ±2.2 ±1.3  | 258       | ABLIKIM     | 16D BES3 | $e^+e^-$ at 3773 MeV |
| 34.3 ±1.4 ±1.7  | 1033 ± 42 | ARTUSO      | 08 CLEO  | See MENDEZ 10        |

$\Gamma(\eta\pi^+)/\Gamma(K^-2\pi^+)$   $\Gamma_{94}/\Gamma_{45}$

Unseen decay modes of the  $\eta$  are included.

| VALUE (units $10^{-2}$ )  | EVTS      | DOCUMENT ID | TECN    | COMMENT                             |
|---|-----------|-------------|---------|-------------------------------------|
| <b>4.02±0.11 OUR FIT</b>  |           |             |         | Error includes scale factor of 1.1. |
| <b>3.87±0.09±0.19</b>   | 2940 ± 68 | MENDEZ      | 10 CLEO | $e^+e^-$ at 3774 MeV                |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |           |             |         |                                     |
| 3.81±0.26±0.21  | 377 ± 26  | RUBIN       | 06 CLEO | See ARTUSO 08                       |

$\Gamma(\eta\pi^+\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{95}/\Gamma$

| VALUE (units $10^{-4}$ ) | EVTS     | DOCUMENT ID | TECN    | COMMENT                  |
|--------------------------|----------|-------------|---------|--------------------------|
| <b>13.8±3.1±1.6</b>      | 149 ± 34 | ARTUSO      | 08 CLEO | $e^+e^-$ at $\psi(3770)$ |

$\Gamma(\omega\pi^+)/\Gamma_{\text{total}}$   $\Gamma_{96}/\Gamma$

Unseen decay modes of the  $\omega$  are included.

| VALUE (units $10^{-4}$ )  | CL% | EVTS | DOCUMENT ID | TECN     | COMMENT                  |
|---|-----|------|-------------|----------|--------------------------|
| <b>2.79±0.57±0.16</b>   |     | 79   | ABLIKIM     | 16D BES3 | $e^+e^-$ at 3773 MeV     |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |     |      |             |          |                          |
| <3.4  |     | 90   | RUBIN       | 06 CLEO  | $e^+e^-$ at $\psi(3770)$ |

$\Gamma(\eta'(958)\pi^+)/\Gamma_{\text{total}}$   $\Gamma_{97}/\Gamma$

Unseen decay modes of the  $\eta'(958)$  are included.

| VALUE (units $10^{-4}$ )  | EVTS     | DOCUMENT ID | TECN     | COMMENT             |
|---|----------|-------------|----------|---------------------|
| <b>49.7±1.9 OUR FIT</b>   |          |             |          |                     |
| <b>51.2±1.4±2.1</b>   | 3.1k     | ABLIKIM     | 18W BES3 | $e^+e^-$ , 3773 MeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |          |             |          |                     |
| 44.2±2.5±2.9  | 352 ± 20 | ARTUSO      | 08 CLEO  | See MENDEZ 10       |

$\Gamma(\eta'(958)\pi^+)/\Gamma(K^-2\pi^+)$   $\Gamma_{97}/\Gamma_{45}$

Unseen decay modes of the  $\eta'(958)$  are included.

| VALUE (units $10^{-2}$ ) | EVTS      | DOCUMENT ID | TECN    | COMMENT              |
|--------------------------|-----------|-------------|---------|----------------------|
| <b>5.30±0.21 OUR FIT</b> |           |             |         |                      |
| <b>5.12±0.17±0.25</b>    | 1037 ± 35 | MENDEZ      | 10 CLEO | $e^+e^-$ at 3774 MeV |

$\Gamma(\eta'(958)\pi^+\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{98}/\Gamma$ Unseen decay modes of the  $\eta'(958)$  are included.

| VALUE (units $10^{-4}$ ) | EVTS   | DOCUMENT ID | TECN | COMMENT                       |
|--------------------------|--------|-------------|------|-------------------------------|
| <b>15.7±4.3±2.5</b>      | 33 ± 9 | ARTUSO      | 08   | CLEO $e^+e^-$ at $\psi(3770)$ |

————— Hadronic modes with a  $K\bar{K}$  pair ————— $\Gamma(K^+K_S^0)/\Gamma_{\text{total}}$   $\Gamma_{99}/\Gamma$ 

| VALUE (units $10^{-3}$ )   | EVTS  | DOCUMENT ID | TECN | COMMENT                  |
|----------------------------|---|-------------|------|--------------------------|
| <b>3.04 ± 0.09 OUR FIT</b> | Error includes scale factor of 2.2.                                     |             |      |                          |
| <b>3.183±0.029±0.060</b>   | 18k   | ABLIKIM     | 18W  | BES3 $e^+e^-$ , 3773 MeV |
| • • •                      | We do not use the following data for averages, fits, limits, etc. • • • |             |      |                          |
| 3.14 ± 0.09 ± 0.08         | 1971 ± 51   | BONVICINI   | 08   | CLEO See MENDEZ 10       |

 $\Gamma(K^+K_S^0)/\Gamma(K_S^0\pi^+)$   $\Gamma_{99}/\Gamma_{43}$ 

| VALUE                            | EVTS  | DOCUMENT ID     | TECN | COMMENT   |
|----------------------------------|---|-----------------|------|---|
| <b>0.194 ± 0.006 OUR FIT</b>     | Error includes scale factor of 2.8.                                     |                 |      |   |
| <b>0.1901±0.0024 OUR AVERAGE</b> |   |                 |      |   |
| 0.1899±0.0011±0.0022             | 101k±561  | WON             | 09   | BELL $e^+e^-$ at $\Upsilon(4S)$                   |
| 0.1892±0.0155±0.0073             | 278 ± 21  | ARMS            | 04   | CLEO $e^+e^- \approx 10$ GeV                      |
| 0.1996±0.0119±0.0096             | 949   | LINK            | 02B  | FOCS $\gamma$ A, $\bar{E}_\gamma \approx 180$ GeV |
| • • •                            | We do not use the following data for averages, fits, limits, etc. • • • |                 |      |   |
| 0.222 ± 0.037 ± 0.013            | 63 ± 10   | ABLIKIM         | 05F  | BES $e^+e^- \approx \psi(3770)$                   |
| 0.222 ± 0.041 ± 0.019            | 70  | BISHAI          | 97   | CLEO See ARMS 04                                  |
| 0.25 ± 0.04 ± 0.02               | 129   | FRABETTI        | 95   | E687 $\gamma$ Be $\bar{E}_\gamma \approx 200$ GeV |
| 0.271 ± 0.065 ± 0.039            | 69  | ANJOS           | 90C  | E691 $\gamma$ Be                                  |
| 0.317 ± 0.086 ± 0.048            | 31  | BALTRUSAIT..85E | MRK3 | $e^+e^-$ 3.77 GeV                                 |
| 0.25 ± 0.15                      | 6   | SCHINDLER       | 81   | MRK2 $e^+e^-$ 3.771 GeV                           |

 $\Gamma(K^+K_S^0)/\Gamma(K^-2\pi^+)$   $\Gamma_{99}/\Gamma_{45}$ 

| VALUE (units $10^{-2}$ ) | EVTS  | DOCUMENT ID       | TECN | COMMENT   |
|--------------------------|---|-------------------|------|---|
| <b>3.24±0.09 OUR FIT</b> | Error includes scale factor of 2.3.                                     |                   |      |   |
| <b>3.35±0.06±0.07</b>    | 5161 ± 86   | MENDEZ            | 10   | CLEO $e^+e^-$ at 3774 MeV                               |
| • • •                    | We do not use the following data for averages, fits, limits, etc. • • • |                   |      |   |
| 3.02±0.18±0.15           | 949   | <sup>1</sup> LINK | 02B  | FOCS $\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV |

<sup>1</sup>This LINK 02B result is redundant with a result in the previous datablock. $\Gamma(K^+K^-\pi^+)/\Gamma_{\text{total}}$   $\Gamma_{100}/\Gamma$ 

| VALUE (units $10^{-2}$ ) | EVTS  | DOCUMENT ID        | TECN | COMMENT               |
|--------------------------|---|--------------------|------|-----------------------|
| • • •                    | We do not use the following data for averages, fits, limits, etc. • • • |                    |      |                       |
| 0.935±0.017±0.024        |   | <sup>1</sup> DOBBS | 07   | CLEO See BONVICINI 14 |
| 0.97 ± 0.04 ± 0.04       | 1250 ± 40   | <sup>1</sup> HE    | 05   | CLEO See DOBBS 07     |

<sup>1</sup>DOBBS 07 and HE 05 use single- and double-tagged events in an overall fit. DOBBS 07 supersedes HE 05.

$$\Gamma(K^+ K^- \pi^+)/\Gamma(K^- 2\pi^+) \quad \Gamma_{100}/\Gamma_{45}$$

| <u>VALUE</u>                     | <u>EVTS</u>          | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>                                |
|----------------------------------|----------------------|--------------------|-------------|---|
| <b>0.1059±0.0018 OUR AVERAGE</b> |                      |                    |             |   |
| 0.106 ±0.002 ±0.003              |                      | BONVICINI          | 14 CLEO     | All CLEO-c runs                               |
| 0.117 ±0.013 ±0.007              | 181 ± 20             | ABLIKIM            | 05F BES     | $e^+ e^- \approx \psi(3770)$                  |
| 0.107 ±0.001 ±0.002              | 43k                  | AUBERT             | 05S BABR    | $e^+ e^- \approx \Upsilon(4S)$                |
| 0.093 ±0.010                     | $^{+0.008}_{-0.006}$ | JUN                | 00 SELX     | $\Sigma^-$ nucleus, 600 GeV                   |
| 0.0976±0.0042±0.0046             |                      | FRABETTI           | 95B E687    | $\gamma$ Be, $\bar{E}_\gamma \approx 200$ GeV |

$$\Gamma(\phi\pi^+, \phi \rightarrow K^+ K^-)/\Gamma(K^+ K^- \pi^+) \quad \Gamma_{101}/\Gamma_{100}$$

This is the “fit fraction” from the Dalitz-plot analysis.

| <u>VALUE (%)</u>  | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>              |
|---|--------------------|-------------|-----------------------------|
| <b>27.8±0.4<math>^{+0.2}_{-0.5}</math></b>                                    | RUBIN              | 08 CLEO     | Dalitz fit, 19,458±163 evts |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                    |             |                             |
| 29.2±3.1±3.0  | FRABETTI           | 95B E687    | Dalitz fit, 915 evts        |

$$\Gamma(K^+ \bar{K}^*(892)^0, \bar{K}^*(892)^0 \rightarrow K^- \pi^+)/\Gamma(K^+ K^- \pi^+) \quad \Gamma_{102}/\Gamma_{100}$$

This is the “fit fraction” from the Dalitz-plot analysis.

| <u>VALUE (%)</u>  | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>              |
|---|--------------------|-------------|-----------------------------|
| <b>25.7±0.5<math>^{+0.4}_{-1.2}</math></b>                                    | RUBIN              | 08 CLEO     | Dalitz fit, 19,458±163 evts |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                    |             |                             |
| 30.1±2.0±2.5  | FRABETTI           | 95B E687    | Dalitz fit, 915 evts        |

$$\Gamma(K^+ \bar{K}_0^*(1430)^0, \bar{K}_0^*(1430)^0 \rightarrow K^- \pi^+)/\Gamma(K^+ K^- \pi^+) \quad \Gamma_{103}/\Gamma_{100}$$

This is the “fit fraction” from the Dalitz-plot analysis.

| <u>VALUE (%)</u>  | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>              |
|---|--------------------|-------------|-----------------------------|
| <b>18.8±1.2<math>^{+3.3}_{-3.4}</math></b>                                    | RUBIN              | 08 CLEO     | Dalitz fit, 19,458±163 evts |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                    |             |                             |
| 37.0±3.5±1.8  | FRABETTI           | 95B E687    | Dalitz fit, 915 evts        |

$$\Gamma(K^+ \bar{K}_2^*(1430)^0, \bar{K}_2^* \rightarrow K^- \pi^+)/\Gamma(K^+ K^- \pi^+) \quad \Gamma_{104}/\Gamma_{100}$$

This is the “fit fraction” from the Dalitz-plot analysis.

| <u>VALUE (%)</u>                          | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>              |
|---|--------------------|-------------|-----------------------------|
| <b>1.7±0.4<math>^{+1.2}_{-0.7}</math></b> | RUBIN              | 08 CLEO     | Dalitz fit, 19,458±163 evts |

$$\Gamma(K^+ \bar{K}_0^*(700), \bar{K}_0^* \rightarrow K^- \pi^+)/\Gamma(K^+ K^- \pi^+) \quad \Gamma_{105}/\Gamma_{100}$$

This is the “fit fraction” from the Dalitz-plot analysis.

| <u>VALUE (%)</u>                          | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>              |
|---|--------------------|-------------|-----------------------------|
| <b>7.0±0.8<math>^{+3.5}_{-2.0}</math></b> | RUBIN              | 08 CLEO     | Dalitz fit, 19,458±163 evts |

$$\Gamma(a_0(1450)^0 \pi^+, a_0^0 \rightarrow K^+ K^-)/\Gamma(K^+ K^- \pi^+) \quad \Gamma_{106}/\Gamma_{100}$$

This is the “fit fraction” from the Dalitz-plot analysis.

| <u>VALUE (%)</u>                          | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>              |
|---|--------------------|-------------|-----------------------------|
| <b>4.6±0.6<math>^{+7.2}_{-1.8}</math></b> | RUBIN              | 08 CLEO     | Dalitz fit, 19,458±163 evts |

$\Gamma(\phi(1680)\pi^+, \phi \rightarrow K^+K^-)/\Gamma(K^+K^-\pi^+)$   $\Gamma_{107}/\Gamma_{100}$

This is the "fit fraction" from the Dalitz-plot analysis.

| VALUE (%)                       | DOCUMENT ID | TECN    | COMMENT                           |
|---------------------------------|-------------|---------|-----------------------------------|
| $0.51 \pm 0.11^{+0.37}_{-0.16}$ | RUBIN       | 08 CLEO | Dalitz fit, $19,458 \pm 163$ evts |

$\Gamma(K_S^0 K_S^0 \pi^+)/\Gamma_{\text{total}}$   $\Gamma_{108}/\Gamma$

| VALUE (units $10^{-4}$ ) | EVTS | DOCUMENT ID | TECN     | COMMENT                         |
|--------------------------|------|-------------|----------|---------------------------------|
| $27.0 \pm 0.5 \pm 1.2$   | 4897 | ABLIKIM     | 17A BES3 | $e^+e^- \rightarrow \psi(3770)$ |

$\Gamma(K^+ K_S^0 \pi^+ \pi^-)/\Gamma(K_S^0 2\pi^+ \pi^-)$   $\Gamma_{109}/\Gamma_{67}$

| VALUE (units $10^{-2}$ ) | EVTS         | DOCUMENT ID | TECN     | COMMENT  |
|--------------------------|--------------|-------------|----------|--|
| $5.62 \pm 0.39 \pm 0.40$ | $469 \pm 32$ | LINK        | 01C FOCS | $\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV |

$\Gamma(K_S^0 K^- 2\pi^+)/\Gamma(K_S^0 2\pi^+ \pi^-)$   $\Gamma_{110}/\Gamma_{67}$

| VALUE (units $10^{-2}$ ) | EVTS         | DOCUMENT ID | TECN     | COMMENT  |
|--------------------------|--------------|-------------|----------|--|
| $7.68 \pm 0.41 \pm 0.32$ | $670 \pm 35$ | LINK        | 01C FOCS | $\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV |

$\Gamma(K^+ K^- 2\pi^+ \pi^-)/\Gamma(K^- 3\pi^+ \pi^-)$   $\Gamma_{111}/\Gamma_{68}$

| VALUE                       | EVTS | DOCUMENT ID | TECN     | COMMENT                                      |
|-----------------------------|------|-------------|----------|--|
| $0.040 \pm 0.009 \pm 0.019$ | 38   | LINK        | 03D FOCS | $\gamma$ A, $\bar{E}_\gamma \approx 180$ GeV |

$\Gamma(\phi\pi^+\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{112}/\Gamma$

Unseen decay modes of the  $\phi$  are included.

| VALUE             | DOCUMENT ID         | TECN     | COMMENT            |
|-------------------|---------------------|----------|--------------------|
| $0.023 \pm 0.010$ | <sup>1</sup> BARLAG | 92C ACCM | $\pi^-$ Cu 230 GeV |

<sup>1</sup> BARLAG 92C computes the branching fraction using topological normalization.

$\Gamma(\phi\rho^+)/\Gamma(K^- 2\pi^+)$   $\Gamma_{113}/\Gamma_{45}$

Unseen decay modes of the  $\phi$  are included.

| VALUE   | CL% | DOCUMENT ID | TECN    | COMMENT                   |
|---------|-----|-------------|---------|---------------------------|
| $<0.16$ | 90  | DAOUDI      | 92 CLEO | $e^+e^- \approx 10.5$ GeV |

$\Gamma(K^+ K^- \pi^+ \pi^0 \text{ non-}\phi)/\Gamma_{\text{total}}$   $\Gamma_{114}/\Gamma$

| VALUE                     | DOCUMENT ID         | TECN     | COMMENT            |
|---------------------------|---------------------|----------|--------------------|
| $0.015^{+0.007}_{-0.006}$ | <sup>1</sup> BARLAG | 92C ACCM | $\pi^-$ Cu 230 GeV |

<sup>1</sup> BARLAG 92C computes the branching fraction using topological normalization.

$\Gamma(K^+ K^- \pi^+ \pi^0 \text{ non-}\phi)/\Gamma(K^- 2\pi^+)$   $\Gamma_{114}/\Gamma_{45}$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

|         |    |       |          |                 |
|---------|----|-------|----------|-----------------|
| $<0.25$ | 90 | ANJOS | 89E E691 | Photoproduction |
|---------|----|-------|----------|-----------------|

$\Gamma(K^*(892)^+ K_S^0)/\Gamma(K_S^0 \pi^+)$   $\Gamma_{115}/\Gamma_{43}$

Unseen decay modes of the  $K^*(892)^+$  are included.

| VALUE                 | EVTS | DOCUMENT ID | TECN    | COMMENT                                      |
|-----------------------|------|-------------|---------|--|
| $1.1 \pm 0.3 \pm 0.4$ | 67   | FRABETTI    | 95 E687 | $\gamma$ Be $\bar{E}_\gamma \approx 200$ GeV |



————— Doubly Cabibbo-suppressed modes —————

$\Gamma(K^+\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{116}/\Gamma$

| VALUE (units $10^{-4}$ )  | EVTS                                | DOCUMENT ID | TECN     | COMMENT                       |
|---|-------------------------------------|-------------|----------|-------------------------------|
| <b>2.08±0.21 OUR FIT</b>  | Error includes scale factor of 1.4. |             |          |                               |
| <b>2.35±0.20 OUR AVERAGE</b>  |                                     |             |          |                               |
| 2.32±0.21±0.06  | 1.8k                                | ABLIKIM     | 18W BES3 | $e^+e^-$ , 3773 MeV           |
| 2.52±0.47±0.26  | 189 ± 37                            | AUBERT,B    | 06F BABR | $e^+e^- \approx \Upsilon(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                                     |             |          |                               |
| 2.28±0.36±0.17  | 148 ± 23                            | DYTMAN      | 06 CLEO  | See MENDEZ 10                 |

$\Gamma(K^+\pi^0)/\Gamma(K^-2\pi^+)$   $\Gamma_{116}/\Gamma_{45}$

| VALUE (units $10^{-3}$ ) | EVTS                                | DOCUMENT ID | TECN    | COMMENT              |
|--------------------------|-------------------------------------|-------------|---------|----------------------|
| <b>2.21±0.23 OUR FIT</b> | Error includes scale factor of 1.5. |             |         |                      |
| <b>1.9 ±0.2 ±0.1</b>     | 343 ± 37                            | MENDEZ      | 10 CLEO | $e^+e^-$ at 3774 MeV |

$\Gamma(K^+\eta)/\Gamma_{\text{total}}$   $\Gamma_{117}/\Gamma$

| VALUE (units $10^{-3}$ )   | EVTS                                | DOCUMENT ID | TECN     | COMMENT             |
|----------------------------|-------------------------------------|-------------|----------|---------------------|
| <b>0.125±0.016 OUR FIT</b> | Error includes scale factor of 1.1. |             |          |                     |
| <b>0.151±0.025±0.014</b>   | 439                                 | ABLIKIM     | 18W BES3 | $e^+e^-$ , 3773 MeV |

$\Gamma(K^+\eta)/\Gamma(\eta\pi^+)$   $\Gamma_{117}/\Gamma_{94}$

| VALUE (%)               | EVTS                                | DOCUMENT ID | TECN    | COMMENT                       |
|-------------------------|-------------------------------------|-------------|---------|-------------------------------|
| <b>3.3 ±0.4 OUR FIT</b> | Error includes scale factor of 1.1. |             |         |                               |
| <b>3.06±0.43±0.14</b>   | 166 ± 23                            | WON         | 11 BELL | $e^+e^- \approx \Upsilon(4S)$ |

$\Gamma(K^+\eta'(958))/\Gamma_{\text{total}}$   $\Gamma_{118}/\Gamma$

| VALUE (units $10^{-3}$ )   | EVTS | DOCUMENT ID | TECN     | COMMENT             |
|----------------------------|------|-------------|----------|---------------------|
| <b>0.185±0.020 OUR FIT</b> |      |             |          |                     |
| <b>0.164±0.051±0.024</b>   | 87   | ABLIKIM     | 18W BES3 | $e^+e^-$ , 3773 MeV |

$\Gamma(K^+\eta'(958))/\Gamma(\eta'(958)\pi^+)$   $\Gamma_{118}/\Gamma_{97}$

| VALUE (%)               | EVTS     | DOCUMENT ID | TECN    | COMMENT                       |
|-------------------------|----------|-------------|---------|-------------------------------|
| <b>3.7 ±0.4 OUR FIT</b> |          |             |         |                               |
| <b>3.77±0.39±0.10</b>   | 180 ± 19 | WON         | 11 BELL | $e^+e^- \approx \Upsilon(4S)$ |

$\Gamma(K^+\pi^+\pi^-)/\Gamma(K^-2\pi^+)$   $\Gamma_{119}/\Gamma_{45}$

| VALUE (units $10^{-3}$ )     | EVTS      | DOCUMENT ID | TECN     | COMMENT                                       |
|------------------------------|-----------|-------------|----------|---|
| <b>5.77±0.22 OUR AVERAGE</b> |           |             |          |   |
| 5.69±0.18±0.14               | 2638 ± 84 | KO          | 09 BELL  | $e^+e^-$ at $\Upsilon(4S)$                    |
| 6.5 ±0.8 ±0.4                | 189 ± 24  | LINK        | 04F FOCS | $\gamma A$ , $\bar{E}_\gamma \approx 180$ GeV |
| 7.7 ±1.7 ±0.8                | 59 ± 13   | AITALA      | 97C E791 | $\pi^- A$ , 500 GeV                           |
| 7.2 ±2.3 ±1.7                | 21        | FRABETTI    | 95E E687 | $\gamma Be$ , $\bar{E}_\gamma = 220$ GeV      |

$\Gamma(K^+\rho^0)/\Gamma(K^+\pi^+\pi^-)$   $\Gamma_{120}/\Gamma_{119}$

This is the "fit fraction" from the Dalitz-plot analysis.

| VALUE                         | DOCUMENT ID | TECN     | COMMENT              |
|-------------------------------|-------------|----------|----------------------|
| <b>0.39 ±0.09 OUR AVERAGE</b> |             |          |                      |
| 0.3943±0.0787±0.0815          | LINK        | 04F FOCS | Dalitz fit, 189 evts |
| 0.37 ±0.14 ±0.07              | AITALA      | 97C E791 | Dalitz fit, 59 evts  |

$\Gamma(K^*(892)^0 \pi^+, K^*(892)^0 \rightarrow K^+ \pi^-) / \Gamma(K^+ \pi^+ \pi^-)$   $\Gamma_{121} / \Gamma_{119}$

This is the "fit fraction" from the Dalitz-plot analysis.

| VALUE                    | DOCUMENT ID        | TECN | COMMENT                   |
|--------------------------|--------------------|------|---------------------------|
| <b>0.47 ± 0.08</b>       | <b>OUR AVERAGE</b> |      |                           |
| 0.5220 ± 0.0684 ± 0.0638 | LINK               | 04F  | FOCS Dalitz fit, 189 evts |
| 0.35 ± 0.14 ± 0.01       | AITALA             | 97C  | E791 Dalitz fit, 59 evts  |

$\Gamma(K^+ f_0(980), f_0(980) \rightarrow \pi^+ \pi^-) / \Gamma(K^+ \pi^+ \pi^-)$   $\Gamma_{122} / \Gamma_{119}$

This is the "fit fraction" from the Dalitz-plot analysis.

| VALUE                           | DOCUMENT ID | TECN | COMMENT                   |
|---------------------------------|-------------|------|---------------------------|
| <b>0.0892 ± 0.0333 ± 0.0412</b> | LINK        | 04F  | FOCS Dalitz fit, 189 evts |

$\Gamma(K_2^*(1430)^0 \pi^+, K_2^*(1430)^0 \rightarrow K^+ \pi^-) / \Gamma(K^+ \pi^+ \pi^-)$   $\Gamma_{123} / \Gamma_{119}$

This is the "fit fraction" from the Dalitz-plot analysis.

| VALUE                           | DOCUMENT ID | TECN | COMMENT                   |
|---------------------------------|-------------|------|---------------------------|
| <b>0.0803 ± 0.0372 ± 0.0391</b> | LINK        | 04F  | FOCS Dalitz fit, 189 evts |

$\Gamma(K^+ \pi^+ \pi^- \text{ nonresonant}) / \Gamma(K^+ \pi^+ \pi^-)$   $\Gamma_{124} / \Gamma_{119}$

This is the "fit fraction" from the Dalitz-plot analysis.

| VALUE              | DOCUMENT ID   | TECN | COMMENT   |
|--------------------|---|------|---|
| • • •              | We do not use the following data for averages, fits, limits, etc. • • • |      |   |
| 0.36 ± 0.14 ± 0.07 | <sup>1</sup> AITALA   | 97C  | E791 Dalitz fit, 59 evts  |
|                    | <sup>1</sup> LINK   | 04F  | with three times as many events, finds no need for a nonresonant amplitude. |

$\Gamma(2K^+ K^-) / \Gamma(K^- 2\pi^+)$   $\Gamma_{125} / \Gamma_{45}$

| VALUE (units 10 <sup>-4</sup> ) | EVTS | DOCUMENT ID       | TECN | COMMENT  |
|---------------------------------|------|-------------------|------|--|
| <b>9.49 ± 2.17 ± 0.22</b>       | 65   | <sup>1</sup> LINK | 02I  | FOCS $\gamma$ nucleus, $\approx 180$ GeV                         |
|                                 |      | <sup>1</sup> LINK | 02I  | finds little evidence for $\phi K^+$ or $f_0(980) K^+$ submodes. |

————— Rare or forbidden modes —————

$\Gamma(\pi^+ e^+ e^-) / \Gamma_{\text{total}}$   $\Gamma_{126} / \Gamma$

A test for the  $\Delta C = 1$  weak neutral current. Allowed by higher-order electroweak interactions.

| VALUE                             | CL%   | DOCUMENT ID        | TECN | COMMENT  |
|-----------------------------------|---|--------------------|------|--|
| <b>&lt; 1.1 × 10<sup>-6</sup></b> | 90  | LEES               | 11G  | BABR $e^+ e^- \approx \Upsilon(4S)$                |
| • • •                             | We do not use the following data for averages, fits, limits, etc. • • • |                    |      |  |
| < 5.9 × 10 <sup>-6</sup>          | 90  | <sup>1</sup> RUBIN | 10   | CLEO $e^+ e^-$ at $\psi(3770)$                     |
| < 7.4 × 10 <sup>-6</sup>          | 90  | HE                 | 05A  | CLEO See RUBIN 10                                  |
| < 5.2 × 10 <sup>-5</sup>          | 90  | AITALA             | 99G  | E791 $\pi^- N$ 500 GeV                             |
| < 1.1 × 10 <sup>-4</sup>          | 90  | FRABETTI           | 97B  | E687 $\gamma$ Be, $\bar{E}_\gamma \approx 220$ GeV |
| < 6.6 × 10 <sup>-5</sup>          | 90  | AITALA             | 96   | E791 $\pi^- N$ 500 GeV                             |
| < 2.5 × 10 <sup>-3</sup>          | 90  | WEIR               | 90B  | MRK2 $e^+ e^-$ 29 GeV                              |
| < 2.6 × 10 <sup>-3</sup>          | 90  | HAAS               | 88   | CLEO $e^+ e^-$ 10 GeV                              |

<sup>1</sup>This RUBIN 10 limit is for the  $e^+ e^-$  mass in the continuum away from the  $\phi(1020)$ . See the next data block.

$\Gamma(\pi^+ \pi^0 e^+ e^-) / \Gamma_{\text{total}}$   $\Gamma_{127} / \Gamma$

| VALUE                             | CL% | DOCUMENT ID | TECN | COMMENT                   |
|-----------------------------------|-----|-------------|------|---------------------------|
| <b>&lt; 1.4 × 10<sup>-5</sup></b> | 90  | ABLIKIM     | 18P  | BES3 $e^+ e^-$ , 3773 MeV |

$\Gamma(\pi^+ \phi, \phi \rightarrow e^+ e^-) / \Gamma_{\text{total}}$   $\Gamma_{128} / \Gamma$

This is *not* a test for the  $\Delta C = 1$  weak neutral current, but leads to the  $\pi^+ e^+ e^-$  final state.

| VALUE  | EVTs | DOCUMENT ID        | TECN | COMMENT                        |
|--|------|--------------------|------|--------------------------------|
| $(1.7^{+1.4}_{-0.9} \pm 0.1) \times 10^{-6}$ | 4    | <sup>1</sup> RUBIN | 10   | CLEO $e^+ e^-$ at $\psi(3770)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|  |   |    |     |                   |
|--|---|----|-----|-------------------|
| $(2.7^{+3.6}_{-1.8} \pm 0.2) \times 10^{-6}$ | 2 | HE | 05A | CLEO See RUBIN 10 |
|--|---|----|-----|-------------------|

<sup>1</sup>This RUBIN 10 result is consistent with the known  $D^+ \rightarrow \phi \pi^+$  and  $\phi \rightarrow e^+ e^-$  fractions.

$\Gamma(\pi^+ \mu^+ \mu^-) / \Gamma_{\text{total}}$   $\Gamma_{129} / \Gamma$

A test for the  $\Delta C = 1$  weak neutral current. Allowed by higher-order electroweak interactions.

| VALUE                  | CL% | DOCUMENT ID | TECN | COMMENT                  |
|------------------------|-----|-------------|------|--------------------------|
| $< 7.3 \times 10^{-8}$ | 90  | AAIJ        | 13AF | LHCb $p\bar{p}$ at 7 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                        |    |                     |     |  |
|------------------------|----|---------------------|-----|--|
| $< 6.5 \times 10^{-6}$ | 90 | LEES                | 11G | BABR $e^+ e^- \approx \Upsilon(4S)$                |
| $< 3.9 \times 10^{-6}$ | 90 | <sup>1</sup> ABAZOV | 08D | D0 $p\bar{p}$ , $E_{\text{cm}} = 1.96$ TeV         |
| $< 8.8 \times 10^{-6}$ | 90 | LINK                | 03F | FOCS $\gamma$ A, $\bar{E}_\gamma \approx 180$ GeV  |
| $< 1.5 \times 10^{-5}$ | 90 | AITALA              | 99G | E791 $\pi^- N$ 500 GeV                             |
| $< 8.9 \times 10^{-5}$ | 90 | FRABETTI            | 97B | E687 $\gamma$ Be, $\bar{E}_\gamma \approx 220$ GeV |
| $< 1.8 \times 10^{-5}$ | 90 | AITALA              | 96  | E791 $\pi^- N$ 500 GeV                             |
| $< 2.2 \times 10^{-4}$ | 90 | KODAMA              | 95  | E653 $\pi^-$ emulsion 600 GeV                      |
| $< 5.9 \times 10^{-3}$ | 90 | WEIR                | 90B | MRK2 $e^+ e^-$ 29 GeV                              |
| $< 2.9 \times 10^{-3}$ | 90 | HAAS                | 88  | CLEO $e^+ e^-$ 10 GeV                              |

<sup>1</sup>This ABAZOV 08D limit is for the  $\mu^+ \mu^-$  mass in the continuum away from the  $\phi(1020)$ . See the next data block.

$\Gamma(\pi^+ \phi, \phi \rightarrow \mu^+ \mu^-) / \Gamma_{\text{total}}$   $\Gamma_{130} / \Gamma$

This is *not* a test for the  $\Delta C = 1$  weak neutral current, but leads to the  $\pi^+ \mu^+ \mu^-$  final state.

| VALUE                                  | DOCUMENT ID         | TECN | COMMENT                                    |
|--|---------------------|------|--|
| $(1.8 \pm 0.5 \pm 0.6) \times 10^{-6}$ | <sup>1</sup> ABAZOV | 08D  | D0 $p\bar{p}$ , $E_{\text{cm}} = 1.96$ TeV |

<sup>1</sup>This ABAZOV 08D value is consistent with the known  $D^+ \rightarrow \phi \pi^+$  and  $\phi \rightarrow \mu^+ \mu^-$  fractions.

$\Gamma(\rho^+ \mu^+ \mu^-) / \Gamma_{\text{total}}$   $\Gamma_{131} / \Gamma$

A test for the  $\Delta C = 1$  weak neutral current. Allowed by higher-order electroweak interactions.

| VALUE                  | CL% | DOCUMENT ID | TECN | COMMENT                       |
|------------------------|-----|-------------|------|-------------------------------|
| $< 5.6 \times 10^{-4}$ | 90  | KODAMA      | 95   | E653 $\pi^-$ emulsion 600 GeV |

$\Gamma(K^+ e^+ e^-) / \Gamma_{\text{total}}$   $\Gamma_{132} / \Gamma$

Both quarks would have to change flavor for this decay to occur.

| VALUE                  | CL% | DOCUMENT ID | TECN | COMMENT                             |
|------------------------|-----|-------------|------|-------------------------------------|
| $< 1.0 \times 10^{-6}$ | 90  | LEES        | 11G  | BABR $e^+ e^- \approx \Upsilon(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                       |    |          |     |      |   |
|-----------------------|----|----------|-----|------|---|
| $<3.0 \times 10^{-6}$ | 90 | RUBIN    | 10  | CLEO | $e^+e^-$ at $\psi(3770)$                      |
| $<6.2 \times 10^{-6}$ | 90 | HE       | 05A | CLEO | See RUBIN 10                                  |
| $<2.0 \times 10^{-4}$ | 90 | AITALA   | 99G | E791 | $\pi^- N$ 500 GeV                             |
| $<2.0 \times 10^{-4}$ | 90 | FRABETTI | 97B | E687 | $\gamma$ Be, $\bar{E}_\gamma \approx 220$ GeV |
| $<4.8 \times 10^{-3}$ | 90 | WEIR     | 90B | MRK2 | $e^+e^-$ 29 GeV                               |

**$\Gamma(K^+\pi^0 e^+ e^-)/\Gamma_{\text{total}}$**   **$\Gamma_{133}/\Gamma$**

| <u>VALUE</u>          | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>           |
|-----------------------|------------|--------------------|-------------|--------------------------|
| $<1.5 \times 10^{-5}$ | 90         | ABLIKIM            | 18P         | BES3 $e^+e^-$ , 3773 MeV |

**$\Gamma(K_S^0 \pi^+ e^+ e^-)/\Gamma_{\text{total}}$**   **$\Gamma_{134}/\Gamma$**

| <u>VALUE</u>          | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>           |
|-----------------------|------------|--------------------|-------------|--------------------------|
| $<2.6 \times 10^{-5}$ | 90         | ABLIKIM            | 18P         | BES3 $e^+e^-$ , 3773 MeV |

**$\Gamma(K_S^0 K^+ e^+ e^-)/\Gamma_{\text{total}}$**   **$\Gamma_{135}/\Gamma$**

| <u>VALUE</u>          | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>           |
|-----------------------|------------|--------------------|-------------|--------------------------|
| $<1.1 \times 10^{-5}$ | 90         | ABLIKIM            | 18P         | BES3 $e^+e^-$ , 3773 MeV |

**$\Gamma(K^+ \mu^+ \mu^-)/\Gamma_{\text{total}}$**   **$\Gamma_{136}/\Gamma$**

Both quarks would have to change flavor for this decay to occur.

| <u>VALUE</u>          | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>                     |
|-----------------------|------------|--------------------|-------------|------------------------------------|
| $<4.3 \times 10^{-6}$ | 90         | LEES               | 11G         | BABR $e^+e^- \approx \Upsilon(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                       |    |          |     |      |   |
|-----------------------|----|----------|-----|------|---|
| $<9.2 \times 10^{-6}$ | 90 | LINK     | 03F | FOCS | $\gamma$ A, $\bar{E}_\gamma \approx 180$ GeV  |
| $<4.4 \times 10^{-5}$ | 90 | AITALA   | 99G | E791 | $\pi^- N$ 500 GeV                             |
| $<9.7 \times 10^{-5}$ | 90 | FRABETTI | 97B | E687 | $\gamma$ Be, $\bar{E}_\gamma \approx 220$ GeV |
| $<3.2 \times 10^{-4}$ | 90 | KODAMA   | 95  | E653 | $\pi^-$ emulsion 600 GeV                      |
| $<9.2 \times 10^{-3}$ | 90 | WEIR     | 90B | MRK2 | $e^+e^-$ 29 GeV                               |

**$\Gamma(\pi^+ e^+ \mu^-)/\Gamma_{\text{total}}$**   **$\Gamma_{137}/\Gamma$**

A test of lepton-family-number conservation.

| <u>VALUE</u>          | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>                     |
|-----------------------|------------|--------------------|-------------|------------------------------------|
| $<2.9 \times 10^{-6}$ | 90         | LEES               | 11G         | BABR $e^+e^- \approx \Upsilon(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                       |    |          |     |      |   |
|-----------------------|----|----------|-----|------|---|
| $<1.1 \times 10^{-4}$ | 90 | FRABETTI | 97B | E687 | $\gamma$ Be, $\bar{E}_\gamma \approx 220$ GeV |
| $<3.3 \times 10^{-3}$ | 90 | WEIR     | 90B | MRK2 | $e^+e^-$ 29 GeV                               |

**$\Gamma(\pi^+ e^- \mu^+)/\Gamma_{\text{total}}$**   **$\Gamma_{138}/\Gamma$**

A test of lepton-family-number conservation.

| <u>VALUE</u>          | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u>                     |
|-----------------------|------------|--------------------|-------------|------------------------------------|
| $<3.6 \times 10^{-6}$ | 90         | LEES               | 11G         | BABR $e^+e^- \approx \Upsilon(4S)$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                       |    |          |     |      |   |
|-----------------------|----|----------|-----|------|---|
| $<1.3 \times 10^{-4}$ | 90 | FRABETTI | 97B | E687 | $\gamma$ Be, $\bar{E}_\gamma \approx 220$ GeV |
| $<3.3 \times 10^{-3}$ | 90 | WEIR     | 90B | MRK2 | $e^+e^-$ 29 GeV                               |

$\Gamma(K^+ e^+ \mu^-)/\Gamma_{\text{total}}$   $\Gamma_{139}/\Gamma$

A test of lepton-family-number conservation.

| VALUE   | CL% | DOCUMENT ID | TECN     | COMMENT                                       |
|---|-----|-------------|----------|---|
| $<1.2 \times 10^{-6}$   | 90  | LEES        | 11G BABR | $e^+ e^- \approx \Upsilon(4S)$                |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● |     |             |          |   |
| $<1.3 \times 10^{-4}$   | 90  | FRABETTI    | 97B E687 | $\gamma$ Be, $\bar{E}_\gamma \approx 220$ GeV |
| $<3.4 \times 10^{-3}$   | 90  | WEIR        | 90B MRK2 | $e^+ e^-$ 29 GeV                              |

$\Gamma(K^+ e^- \mu^+)/\Gamma_{\text{total}}$   $\Gamma_{140}/\Gamma$

A test of lepton-family-number conservation.

| VALUE   | CL% | DOCUMENT ID | TECN     | COMMENT                                       |
|---|-----|-------------|----------|---|
| $<2.8 \times 10^{-6}$   | 90  | LEES        | 11G BABR | $e^+ e^- \approx \Upsilon(4S)$                |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● |     |             |          |   |
| $<1.2 \times 10^{-4}$   | 90  | FRABETTI    | 97B E687 | $\gamma$ Be, $\bar{E}_\gamma \approx 220$ GeV |
| $<3.4 \times 10^{-3}$   | 90  | WEIR        | 90B MRK2 | $e^+ e^-$ 29 GeV                              |

$\Gamma(\pi^- 2e^+)/\Gamma_{\text{total}}$   $\Gamma_{141}/\Gamma$

A test of lepton-number conservation.

| VALUE   | CL% | DOCUMENT ID | TECN     | COMMENT                                       |
|---|-----|-------------|----------|---|
| $<1.1 \times 10^{-6}$   | 90  | RUBIN       | 10 CLEO  | $e^+ e^-$ at $\psi(3770)$                     |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● |     |             |          |   |
| $<1.9 \times 10^{-6}$   | 90  | LEES        | 11G BABR | $e^+ e^- \approx \Upsilon(4S)$                |
| $<3.6 \times 10^{-6}$   | 90  | HE          | 05A CLEO | See RUBIN 10                                  |
| $<9.6 \times 10^{-5}$   | 90  | AITALA      | 99G E791 | $\pi^- N$ 500 GeV                             |
| $<1.1 \times 10^{-4}$   | 90  | FRABETTI    | 97B E687 | $\gamma$ Be, $\bar{E}_\gamma \approx 220$ GeV |
| $<4.8 \times 10^{-3}$   | 90  | WEIR        | 90B MRK2 | $e^+ e^-$ 29 GeV                              |

$\Gamma(\pi^- 2\mu^+)/\Gamma_{\text{total}}$   $\Gamma_{142}/\Gamma$

A test of lepton-number conservation.

| VALUE   | CL% | DOCUMENT ID | TECN      | COMMENT                                       |
|---|-----|-------------|-----------|---|
| $<2.2 \times 10^{-8}$   | 90  | AAIJ        | 13AF LHCB | $pp$ at 7 TeV                                 |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● |     |             |           |   |
| $<2.0 \times 10^{-6}$   | 90  | LEES        | 11G BABR  | $e^+ e^- \approx \Upsilon(4S)$                |
| $<4.8 \times 10^{-6}$   | 90  | LINK        | 03F FOCS  | $\gamma$ A, $\bar{E}_\gamma \approx 180$ GeV  |
| $<1.7 \times 10^{-5}$   | 90  | AITALA      | 99G E791  | $\pi^- N$ 500 GeV                             |
| $<8.7 \times 10^{-5}$   | 90  | FRABETTI    | 97B E687  | $\gamma$ Be, $\bar{E}_\gamma \approx 220$ GeV |
| $<2.2 \times 10^{-4}$   | 90  | KODAMA      | 95 E653   | $\pi^-$ emulsion 600 GeV                      |
| $<6.8 \times 10^{-3}$   | 90  | WEIR        | 90B MRK2  | $e^+ e^-$ 29 GeV                              |

$\Gamma(\pi^- e^+ \mu^+)/\Gamma_{\text{total}}$   $\Gamma_{143}/\Gamma$

A test of lepton-number conservation.

| VALUE   | CL% | DOCUMENT ID | TECN     | COMMENT                                       |
|---|-----|-------------|----------|---|
| $<2.0 \times 10^{-6}$   | 90  | LEES        | 11G BABR | $e^+ e^- \approx \Upsilon(4S)$                |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● |     |             |          |   |
| $<5.0 \times 10^{-5}$   | 90  | AITALA      | 99G E791 | $\pi^- N$ 500 GeV                             |
| $<1.1 \times 10^{-4}$   | 90  | FRABETTI    | 97B E687 | $\gamma$ Be, $\bar{E}_\gamma \approx 220$ GeV |
| $<3.7 \times 10^{-3}$   | 90  | WEIR        | 90B MRK2 | $e^+ e^-$ 29 GeV                              |

$\Gamma(\rho^- 2\mu^+)/\Gamma_{\text{total}}$   $\Gamma_{144}/\Gamma$

A test of lepton-number conservation.

| VALUE                 | CL% | DOCUMENT ID | TECN    | COMMENT                  |
|-----------------------|-----|-------------|---------|--------------------------|
| $<5.6 \times 10^{-4}$ | 90  | KODAMA      | 95 E653 | $\pi^-$ emulsion 600 GeV |

$\Gamma(K^- 2e+)/\Gamma_{\text{total}}$   $\Gamma_{145}/\Gamma$

A test of lepton-number conservation.

| VALUE   | CL% | DOCUMENT ID | TECN     | COMMENT                                       |
|---|-----|-------------|----------|---|
| $<0.9 \times 10^{-6}$   | 90  | LEES        | 11G BABR | $e^+ e^- \approx \Upsilon(4S)$                |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |     |             |          |   |
| $<3.5 \times 10^{-6}$   | 90  | RUBIN       | 10 CLEO  | $e^+ e^-$ at $\psi(3770)$                     |
| $<4.5 \times 10^{-6}$   | 90  | HE          | 05A CLEO | See RUBIN 10                                  |
| $<1.2 \times 10^{-4}$   | 90  | FRABETTI    | 97B E687 | $\gamma$ Be, $\bar{E}_\gamma \approx 220$ GeV |
| $<9.1 \times 10^{-3}$   | 90  | WEIR        | 90B MRK2 | $e^+ e^-$ 29 GeV                              |

$\Gamma(K^- 2\mu^+)/\Gamma_{\text{total}}$   $\Gamma_{146}/\Gamma$

A test of lepton-number conservation.

| VALUE   | CL% | DOCUMENT ID | TECN     | COMMENT                                       |
|---|-----|-------------|----------|---|
| $<10 \times 10^{-6}$  | 90  | LEES        | 11G BABR | $e^+ e^- \approx \Upsilon(4S)$                |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |     |             |          |   |
| $<1.3 \times 10^{-5}$   | 90  | LINK        | 03F FOCS | $\gamma$ A, $\bar{E}_\gamma \approx 180$ GeV  |
| $<1.2 \times 10^{-4}$   | 90  | FRABETTI    | 97B E687 | $\gamma$ Be, $\bar{E}_\gamma \approx 220$ GeV |
| $<3.2 \times 10^{-4}$   | 90  | KODAMA      | 95 E653  | $\pi^-$ emulsion 600 GeV                      |
| $<4.3 \times 10^{-3}$   | 90  | WEIR        | 90B MRK2 | $e^+ e^-$ 29 GeV                              |

$\Gamma(K^- e^+ \mu^+)/\Gamma_{\text{total}}$   $\Gamma_{147}/\Gamma$

A test of lepton-number conservation.

| VALUE   | CL% | DOCUMENT ID | TECN     | COMMENT                                       |
|---|-----|-------------|----------|---|
| $<1.9 \times 10^{-6}$   | 90  | LEES        | 11G BABR | $e^+ e^- \approx \Upsilon(4S)$                |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |     |             |          |   |
| $<1.3 \times 10^{-4}$   | 90  | FRABETTI    | 97B E687 | $\gamma$ Be, $\bar{E}_\gamma \approx 220$ GeV |
| $<4.0 \times 10^{-3}$   | 90  | WEIR        | 90B MRK2 | $e^+ e^-$ 29 GeV                              |

$\Gamma(K^*(892)^- 2\mu^+)/\Gamma_{\text{total}}$   $\Gamma_{148}/\Gamma$

A test of lepton-number conservation.

| VALUE                 | CL% | DOCUMENT ID | TECN    | COMMENT                  |
|-----------------------|-----|-------------|---------|--------------------------|
| $<8.5 \times 10^{-4}$ | 90  | KODAMA      | 95 E653 | $\pi^-$ emulsion 600 GeV |

**$D^\pm$  CP-VIOLATING DECAY-RATE ASYMMETRIES**

This is the difference between  $D^+$  and  $D^-$  partial widths for the decay to state  $f$ , divided by the sum of the widths:

$$A_{CP}(f) = [\Gamma(D^+ \rightarrow f) - \Gamma(D^- \rightarrow \bar{f})] / [\Gamma(D^+ \rightarrow f) + \Gamma(D^- \rightarrow \bar{f})].$$

**$A_{CP}(\mu^\pm \nu)$  in  $D^+ \rightarrow \mu^+ \nu_\mu$ ,  $D^- \rightarrow \mu^- \bar{\nu}_\mu$**

| VALUE (%)  | DOCUMENT ID | TECN    | COMMENT                   |
|------------|-------------|---------|---------------------------|
| $+8 \pm 8$ | EISENSTEIN  | 08 CLEO | $e^+ e^-$ at $\psi(3770)$ |

**$A_{CP}(K_L^0 e^\pm \nu)$  in  $D^+ \rightarrow K_L^0 e^+ \nu_e$ ,  $D^- \rightarrow K_L^0 e^- \bar{\nu}_e$**

| VALUE (%)                                   | DOCUMENT ID | TECN      | COMMENT            |
|---|-------------|-----------|--------------------|
| <b><math>-0.59 \pm 0.60 \pm 1.48</math></b> | ABLIKIM     | 15AF BES3 | $e^+ e^-$ 3773 MeV |

**$A_{CP}(K_S^0 \pi^\pm)$  in  $D^\pm \rightarrow K_S^0 \pi^\pm$**

| VALUE (%)                                      | EVTs  | DOCUMENT ID       | TECN     | COMMENT  |
|--|-------|-------------------|----------|--|
| <b><math>-0.41 \pm 0.09</math> OUR AVERAGE</b> |       |                   |          |  |
| $-1.1 \pm 0.6 \pm 0.2$                         |       | BONVICINI         | 14 CLEO  | All CLEO-c runs                                    |
| $-0.363 \pm 0.094 \pm 0.067$                   | 1738k | <sup>1</sup> KO   | 12A BELL | $e^+ e^- \approx \Upsilon(nS)$                     |
| $-0.44 \pm 0.13 \pm 0.10$                      | 807k  | DEL-AMO-SA..11H   | BABR     | $e^+ e^- \approx \Upsilon(4S)$                     |
| $-1.6 \pm 1.5 \pm 0.9$                         | 10.6k | <sup>2</sup> LINK | 02B FOCS | $\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                           |     |        |         |                  |
|---------------------------|-----|--------|---------|------------------|
| $-0.71 \pm 0.19 \pm 0.20$ |     | KO     | 10 BELL | See KO 12A       |
| $-1.3 \pm 0.7 \pm 0.3$    | 30k | MENDEZ | 10 CLEO | See BONVICINI 14 |
| $-0.6 \pm 1.0 \pm 0.3$    |     | DOBBS  | 07 CLEO | See MENDEZ 10    |

<sup>1</sup> KO 12A finds that after subtracting the contribution due to  $K^0 - \bar{K}^0$  mixing, the CP asymmetry due to the change of charm is  $(-0.024 \pm 0.094 \pm 0.067)\%$ , consistent with zero.

<sup>2</sup> LINK 02B measures  $N(D^+ \rightarrow K_S^0 \pi^+)/N(D^+ \rightarrow K^- \pi^+ \pi^+)$ , the ratio of numbers of events observed, and similarly for the  $D^-$ .

**$A_{CP}(K^\mp 2\pi^\pm)$  in  $D^+ \rightarrow K^- 2\pi^+$ ,  $D^- \rightarrow K^+ 2\pi^-$**

| VALUE (%)                                      | EVTs | DOCUMENT ID | TECN | COMMENT |
|--|------|-------------|------|---------|
| <b><math>-0.18 \pm 0.16</math> OUR AVERAGE</b> |      |             |      |         |

|                           |      |           |         |                                    |
|---------------------------|------|-----------|---------|------------------------------------|
| $-0.16 \pm 0.15 \pm 0.09$ | 2.3M | ABAZOV    | 14L D0  | $p\bar{p}$ , $\sqrt{s} = 1.96$ TeV |
| $-0.3 \pm 0.2 \pm 0.4$    |      | BONVICINI | 14 CLEO | All CLEO-c runs                    |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                        |      |        |         |                  |
|------------------------|------|--------|---------|------------------|
| $-0.1 \pm 0.4 \pm 0.9$ | 231k | MENDEZ | 10 CLEO | See BONVICINI 14 |
| $-0.5 \pm 0.4 \pm 0.9$ |      | DOBBS  | 07 CLEO | See MENDEZ 10    |

**$A_{CP}(K^\mp \pi^\pm \pi^\pm \pi^0)$  in  $D^+ \rightarrow K^- \pi^+ \pi^+ \pi^0$ ,  $D^- \rightarrow K^+ \pi^- \pi^- \pi^0$**

| VALUE (%)                                | DOCUMENT ID | TECN    | COMMENT         |
|--|-------------|---------|-----------------|
| <b><math>-0.3 \pm 0.6 \pm 0.4</math></b> | BONVICINI   | 14 CLEO | All CLEO-c runs |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                       |       |         |                  |
|-----------------------|-------|---------|------------------|
| $1.0 \pm 0.9 \pm 0.9$ | DOBBS | 07 CLEO | See BONVICINI 14 |
|-----------------------|-------|---------|------------------|

**$A_{CP}(K_S^0 \pi^\pm \pi^0)$  in  $D^+ \rightarrow K_S^0 \pi^+ \pi^0$ ,  $D^- \rightarrow K_S^0 \pi^- \pi^0$**

| VALUE (%)                                | DOCUMENT ID | TECN    | COMMENT         |
|--|-------------|---------|-----------------|
| <b><math>-0.1 \pm 0.7 \pm 0.2</math></b> | BONVICINI   | 14 CLEO | All CLEO-c runs |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                       |       |         |                  |
|-----------------------|-------|---------|------------------|
| $0.3 \pm 0.9 \pm 0.3$ | DOBBS | 07 CLEO | See BONVICINI 14 |
|-----------------------|-------|---------|------------------|

**$A_{CP}(K_S^0 \pi^\pm \pi^+ \pi^-)$  in  $D^+ \rightarrow K_S^0 \pi^+ \pi^+ \pi^-$ ,  $D^- \rightarrow K_S^0 \pi^- \pi^- \pi^+$**

| VALUE (%)                               | DOCUMENT ID | TECN    | COMMENT         |
|---|-------------|---------|-----------------|
| <b><math>0.0 \pm 1.2 \pm 0.3</math></b> | BONVICINI   | 14 CLEO | All CLEO-c runs |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                       |       |         |                  |
|-----------------------|-------|---------|------------------|
| $0.1 \pm 1.1 \pm 0.6$ | DOBBS | 07 CLEO | See BONVICINI 14 |
|-----------------------|-------|---------|------------------|

### $A_{CP}(\pi^\pm \pi^0)$ in $D^\pm \rightarrow \pi^\pm \pi^0$

| VALUE (%)                    | EVTS | DOCUMENT ID | TECN    | COMMENT                                 |
|------------------------------|------|-------------|---------|---|
| <b>2.4 ± 1.2 OUR AVERAGE</b> |      |             |         |   |
| 2.31 ± 1.24 ± 0.23           | 108k | BABU        | 18 BELL | At/near $\Upsilon(4S)$ , $\Upsilon(5S)$ |
| 2.9 ± 2.9 ± 0.3              | 2.6k | MENDEZ      | 10 CLEO | $e^+e^-$ at 3774 MeV                    |

### $A_{CP}(\pi^\pm \eta)$ in $D^\pm \rightarrow \pi^\pm \eta$

| VALUE (%)  | EVTS | DOCUMENT ID | TECN    | COMMENT                       |
|--|------|-------------|---------|-------------------------------|
| <b>1.0 ± 1.5 OUR AVERAGE</b> Error includes scale factor of 1.4. |      |             |         |                               |
| +1.74 ± 1.13 ± 0.19  |      | WON         | 11 BELL | $e^+e^- \approx \Upsilon(4S)$ |
| -2.0 ± 2.3 ± 0.3   | 2.9k | MENDEZ      | 10 CLEO | $e^+e^-$ at 3774 MeV          |

### $A_{CP}(\pi^\pm \eta'(958))$ in $D^\pm \rightarrow \pi^\pm \eta'(958)$

| VALUE (%)                     | EVTS | DOCUMENT ID | TECN      | COMMENT                       |
|-------------------------------|------|-------------|-----------|-------------------------------|
| <b>-0.6 ± 0.7 OUR AVERAGE</b> |      |             |           |                               |
| -0.61 ± 0.72 ± 0.54           | 63k  | AAIJ        | 17AF LHCb | $pp$ at 7, 8 TeV              |
| -0.12 ± 1.12 ± 0.17           |      | WON         | 11 BELL   | $e^+e^- \approx \Upsilon(4S)$ |
| -4.0 ± 3.4 ± 0.3              | 1.0k | MENDEZ      | 10 CLEO   | $e^+e^-$ at 3774 MeV          |

### $A_{CP}(\bar{K}^0/K^0 K^\pm)$ in $D^+ \rightarrow \bar{K}^0 K^+$ , $D^- \rightarrow K^0 K^-$

| VALUE (%)                      | EVTS | DOCUMENT ID       | TECN      | COMMENT                    |
|--------------------------------|------|-------------------|-----------|----------------------------|
| <b>0.11 ± 0.17 OUR AVERAGE</b> |      |                   |           |                            |
| 0.03 ± 0.17 ± 0.14             | 1.0M | <sup>1</sup> AAIJ | 14BD LHCb | $pp$ at 7, 8 TeV           |
| 0.08 ± 0.28 ± 0.14             | 277k | KO                | 13 BELL   | $e^+e^-$ at $\Upsilon(4S)$ |
| 0.46 ± 0.36 ± 0.25             | 159k | LEES              | 13E BABR  | $e^+e^-$ at $\Upsilon(4S)$ |

<sup>1</sup>AAIJ 14BD reports its result as  $A_{CP}(D^\pm \rightarrow K_S^0 \pi^\pm)$  with  $CP$ -violation effects in the  $K^0 - \bar{K}^0$  system subtracted. It also measures  $A_{CP}(D^\pm \rightarrow \bar{K}^0/K^0 K^\pm) + A_{CP}(D_S^\pm \rightarrow \bar{K}^0/K^0 \pi^\pm) = (0.41 \pm 0.49 \pm 0.26)\%$ .

### $A_{CP}(K_S^0 K^\pm)$ in $D^\pm \rightarrow K_S^0 K^\pm$

| VALUE (%)                       | EVTS | DOCUMENT ID       | TECN     | COMMENT  |
|---------------------------------|------|-------------------|----------|--|
| <b>-0.11 ± 0.25 OUR AVERAGE</b> |      |                   |          |  |
| -0.25 ± 0.28 ± 0.14             | 277k | KO                | 13 BELL  | $e^+e^-$ at $\Upsilon(nS)$                         |
| 0.13 ± 0.36 ± 0.25              | 159k | LEES              | 13E BABR | $e^+e^-$ at $\Upsilon(4S)$                         |
| -0.2 ± 1.5 ± 0.9                | 5.2k | MENDEZ            | 10 CLEO  | $e^+e^-$ at 3774 MeV                               |
| 7.1 ± 6.1 ± 1.2                 | 949  | <sup>1</sup> LINK | 02B FOCS | $\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                     |     |                   |          |  |
|---------------------|-----|-------------------|----------|--|
| -0.16 ± 0.58 ± 0.25 |     | KO                | 10 BELL  | $e^+e^- \approx \Upsilon(4S)$                      |
| 6.9 ± 6.0 ± 1.5     | 949 | <sup>2</sup> LINK | 02B FOCS | $\gamma$ nucleus, $\bar{E}_\gamma \approx 180$ GeV |

<sup>1</sup>LINK 02B measures  $N(D^+ \rightarrow K_S^0 K^+)/N(D^+ \rightarrow K_S^0 \pi^+)$ , the ratio of numbers of events observed, and similarly for the  $D^-$ .

<sup>2</sup>LINK 02B measures  $N(D^+ \rightarrow K_S^0 K^+)/N(D^+ \rightarrow K^- \pi^+ \pi^+)$ , the ratio of numbers of events observed, and similarly for the  $D^-$ .



**$A_{CP}(K^+ K^- \pi^\pm)$  in  $D^\pm \rightarrow K^+ K^- \pi^\pm$** 

See also AAIJ 11G for a search for  $CP$  asymmetry in the  $D^\pm \rightarrow K^+ K^- \pi^\pm$  Dalitz plots using 370k decays and four different binning schemes. No evidence for  $CP$  asymmetry was found.

| <u>VALUE (%)</u>  | <u>EVTS</u> | <u>DOCUMENT ID</u>     | <u>TECN</u> | <u>COMMENT</u>                      |
|---|-------------|------------------------|-------------|-------------------------------------|
| <b>0.37±0.29 OUR AVERAGE</b>  |             |                        |             |                                     |
| 0.37±0.30±0.15  | 224k        | <sup>1</sup> LEES      | 13F BABR    | $e^+ e^-$ at $\Upsilon(4S)$         |
| -0.03±0.84±0.29   |             | RUBIN                  | 08 CLEO     | $e^+ e^-$ at 3774 MeV               |
| 1.4 ±1.0 ±0.8   | 43k         | <sup>2</sup> AUBERT    | 05S BABR    | $e^+ e^-$ at $\Upsilon(4S)$         |
| 0.6 ±1.1 ±0.5   | 14k         | <sup>3</sup> LINK      | 00B FOCS    |                                     |
| -1.4 ±2.9   |             | <sup>3</sup> AITALA    | 97B E791    | $-0.062 < A_{CP} < +0.034$ (90% CL) |
| -3.1 ±6.8   |             | <sup>3</sup> FRABETTI  | 94I E687    | $-0.14 < A_{CP} < +0.081$ (90% CL)  |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● |             |                        |             |                                     |
| -0.1 ±0.9 ±0.4  |             | <sup>4</sup> BONVICINI | 14 CLEO     | See RUBIN 08                        |
| -0.1 ±1.5 ±0.8  |             | DOBBS                  | 07 CLEO     | See BONVICINI 14 and RUBIN 08       |

<sup>1</sup> This is the integrated  $CP$  asymmetry. LEES 13F also searches for  $CP$  asymmetries in four regions of the Dalitz plots (two of which are listed below); in comparisons of binned  $D^+$  and  $D^-$  Dalitz plots; in parametrized fits to those plots, including 2-body submodes; and in comparisons of Legendre-polynomial distributions for the  $K^+ K^-$  and  $K^- \pi^+$  systems.

<sup>2</sup> AUBERT 05S measures  $N(D^+ \rightarrow K^+ K^- \pi^+)/N(D_s^+ \rightarrow K^+ K^- \pi^+)$ , the ratio of the numbers of events observed, and similarly for the  $D^-$ .

<sup>3</sup> FRABETTI 94I, AITALA 98C, and LINK 00B measure  $N(D^+ \rightarrow K^- K^+ \pi^+)/N(D^+ \rightarrow K^- \pi^+ \pi^+)$ , the ratio of numbers of events observed, and similarly for the  $D^-$ .

<sup>4</sup> RUBIN 08 performs a dedicated analysis of this decay mode on the same dataset, with slightly better precision. We therefore take it that BONVICINI 14 does not supersede RUBIN 08's  $A_{CP}$  result.

 **$A_{CP}(K^\pm K^{*0})$  in  $D^+ \rightarrow K^+ \bar{K}^{*0}$ ,  $D^- \rightarrow K^- K^{*0}$** 

| <u>VALUE (%)</u>              | <u>EVTS</u> | <u>DOCUMENT ID</u>    | <u>TECN</u> | <u>COMMENT</u>                      |
|-------------------------------|-------------|-----------------------|-------------|-------------------------------------|
| <b>- 0.3± 0.4 OUR AVERAGE</b> |             |                       |             |                                     |
| - 0.3± 0.4±0.2                | 73k         | <sup>1</sup> LEES     | 13F BABR    | $e^+ e^-$ at $\Upsilon(4S)$         |
| - 0.4± 2.0±0.6                |             | RUBIN                 | 08 CLEO     | Fit-fraction asymmetry              |
| + 0.9± 1.7±0.7                | 11k         | <sup>2</sup> AUBERT   | 05S BABR    | $e^+ e^-$ at $\Upsilon(4S)$         |
| - 1.0± 5.0                    |             | <sup>3</sup> AITALA   | 97B E791    | $-0.092 < A_{CP} < +0.072$ (90% CL) |
| -12 ±13                       |             | <sup>3</sup> FRABETTI | 94I E687    | $-0.33 < A_{CP} < +0.094$ (90% CL)  |

<sup>1</sup> This LEES 13F result is for the  $K^\mp \pi^\pm$  mass-squared between 0.4 and 1.0  $\text{GeV}^2$ , and does not actually separate out the  $K^*$ .

<sup>2</sup> AUBERT 05S measures  $N(D^+ \rightarrow K^+ \bar{K}^{*0})/N(D_s^+ \rightarrow K^+ K^- \pi^+)$ , the ratio of the numbers of events observed, and similarly for the  $D^-$ .

<sup>3</sup> FRABETTI 94I and AITALA 97B measure  $N(D^+ \rightarrow K^+ \bar{K}^*(892)^0)/N(D^+ \rightarrow K^- \pi^+ \pi^+)$ , the ratio of numbers of events observed, and similarly for the  $D^-$ .

### $A_{CP}(\phi\pi^\pm)$ in $D^\pm \rightarrow \phi\pi^\pm$

| VALUE (%)   | EVTS               | DOCUMENT ID                         | TECN | COMMENT                                  |
|---|--------------------|-------------------------------------|------|--|
| <b><math>0.09 \pm 0.19</math></b>                                 | <b>OUR AVERAGE</b> | Error includes scale factor of 1.2. |      |  |
| $-0.04 \pm 0.14 \pm 0.14$   | 1.58M              | AAIJ                                | 13W  | LHCB $pp$ at 7 TeV                       |
| $-0.3 \pm 0.3 \pm 0.5$  | 97k                | <sup>1</sup> LEES                   | 13F  | BABR $e^+e^-$ at $\Upsilon(4S)$          |
| $+0.51 \pm 0.28 \pm 0.05$   | 237k               | STARIC                              | 12   | BELL Mainly at $\Upsilon(4S)$            |
| $-1.8 \pm 1.6 \begin{smallmatrix} +0.2 \\ -0.4 \end{smallmatrix}$ |                    | RUBIN                               | 08   | CLEO Fit-fraction asymmetry              |
| $+0.2 \pm 1.5 \pm 0.6$  | 10k                | <sup>2</sup> AUBERT                 | 05S  | BABR $e^+e^-$ at $\Upsilon(4S)$          |
| $-2.8 \pm 3.6$  |                    | <sup>3</sup> AITALA                 | 97B  | E791 $-0.087 < A_{CP} < +0.031$ (90% CL) |
| $+6.6 \pm 8.6$  |                    | <sup>3</sup> FRABETTI               | 94I  | E687 $-0.075 < A_{CP} < +0.21$ (90% CL)  |

<sup>1</sup> This LEES 13F result is for the  $K^+K^-$  mass-squared less than  $1.3 \text{ GeV}^2$  and the  $K^\mp\pi^\pm$  mass-squared above  $1.0 \text{ GeV}^2$ , and does not actually separate out the  $\phi$ .

<sup>2</sup> AUBERT 05S measures  $N(D^+ \rightarrow \phi\pi^+)/N(D_s^+ \rightarrow K^+K^-\pi^+)$ , the ratio of the numbers of events observed, and similarly for the  $D^-$ .

<sup>3</sup> FRABETTI 94I and AITALA 97B measure  $N(D^+ \rightarrow \phi\pi^+)/N(D^+ \rightarrow K^-\pi^+\pi^+)$ , the ratio of numbers of events observed, and similarly for the  $D^-$ .

### $A_{CP}(K^\pm K_0^*(1430)^0)$ in $D^+ \rightarrow K^+\bar{K}_0^*(1430)^0$ , $D^- \rightarrow K^-K_0^*(1430)^0$

| VALUE (%)   | DOCUMENT ID | TECN | COMMENT                     |
|---|-------------|------|-----------------------------|
| $+8 \pm 6 \begin{smallmatrix} +4 \\ -2 \end{smallmatrix}$ | RUBIN       | 08   | CLEO Fit-fraction asymmetry |

### $A_{CP}(K^\pm K_2^*(1430)^0)$ in $D^+ \rightarrow K^+\bar{K}_2^*(1430)^0$ , $D^- \rightarrow K^-K_2^*(1430)^0$

| VALUE (%)  | DOCUMENT ID | TECN | COMMENT                     |
|--|-------------|------|-----------------------------|
| $+43 \pm 19 \begin{smallmatrix} +5 \\ -18 \end{smallmatrix}$ | RUBIN       | 08   | CLEO Fit-fraction asymmetry |

### $A_{CP}(K^\pm K_0^*(700))$ in $D^+ \rightarrow K^+\bar{K}_0^*(700)$ , $D^- \rightarrow K^-K_0^*(700)$

| VALUE (%)  | DOCUMENT ID | TECN | COMMENT                     |
|--|-------------|------|-----------------------------|
| $-12 \pm 11 \begin{smallmatrix} +14 \\ -6 \end{smallmatrix}$ | RUBIN       | 08   | CLEO Fit-fraction asymmetry |

### $A_{CP}(a_0(1450)^0\pi^\pm)$ in $D^\pm \rightarrow a_0(1450)^0\pi^\pm$

| VALUE (%)  | DOCUMENT ID | TECN | COMMENT                     |
|--|-------------|------|-----------------------------|
| $-19 \pm 12 \begin{smallmatrix} +8 \\ -11 \end{smallmatrix}$ | RUBIN       | 08   | CLEO Fit-fraction asymmetry |

### $A_{CP}(\phi(1680)\pi^\pm)$ in $D^\pm \rightarrow \phi(1680)\pi^\pm$

| VALUE (%)          | DOCUMENT ID | TECN | COMMENT                     |
|--------------------|-------------|------|-----------------------------|
| $-9 \pm 22 \pm 14$ | RUBIN       | 08   | CLEO Fit-fraction asymmetry |

### $A_{CP}(\pi^+\pi^-\pi^\pm)$ in $D^\pm \rightarrow \pi^+\pi^-\pi^\pm$

See also AAJ 14C for a search for  $CP$  violation in  $D^\pm \rightarrow \pi^+\pi^-\pi^\pm$  Dalitz plots using model-independent binned and unbinned methods. No evidence was found.

| VALUE (%)      | DOCUMENT ID         | TECN | COMMENT                                  |
|----------------|---------------------|------|--|
| $-1.7 \pm 4.2$ | <sup>1</sup> AITALA | 97B  | E791 $-0.086 < A_{CP} < +0.052$ (90% CL) |

<sup>1</sup> AITALA 97B measure  $N(D^+ \rightarrow \pi^+\pi^-\pi^+)/N(D^+ \rightarrow K^-\pi^+\pi^+)$ , the ratio of numbers of events observed, and similarly for the  $D^-$ .

### $A_{CP}(K_S^0 K^\pm \pi^+ \pi^-)$ in $D^\pm \rightarrow K_S^0 K^\pm \pi^+ \pi^-$

| VALUE (%)              | EVTS         | DOCUMENT ID | TECN     | COMMENT                                      |
|------------------------|--------------|-------------|----------|--|
| $-4.2 \pm 6.4 \pm 2.2$ | $523 \pm 32$ | LINK        | 05E FOCS | $\gamma$ A, $\bar{E}_\gamma \approx 180$ GeV |

### $A_{CP}(K^\pm \pi^0)$ in $D^\pm \rightarrow K^\pm \pi^0$

| VALUE (%)               | EVTS         | DOCUMENT ID | TECN    | COMMENT               |
|-------------------------|--------------|-------------|---------|-----------------------|
| $-3.5 \pm 10.7 \pm 0.9$ | $343 \pm 37$ | MENDEZ      | 10 CLEO | $e^+ e^-$ at 3774 MeV |

## $D^\pm \chi^2$ TESTS OF CP-VIOLATION (CPV)

We list model-independent searches for local CP violation in phase-space distributions of multi-body decays.

Most of these searches divide phase space (Dalitz plot for 3-body decays, five-dimensional equivalent for 4-body decays) into bins, and perform a  $\chi^2$  test comparing normalised yields  $N_i, \bar{N}_i$  in CP-conjugate bin pairs  $i$ :  $\chi^2 = \sum_i (N_i - \alpha \bar{N}_i) / \sigma (N_i - \alpha \bar{N}_i)$ . The factor  $\alpha = (\sum_i N_i) / (\sum_i \bar{N}_i)$  removes the dependence on phase-space-integrated rate asymmetries. The result is used to obtain the probability (p-value) to obtain the measured  $\chi^2$  or larger under the assumption of CP conservation [AUBERT 08AO, BEDIAGA 09]. Alternative methods obtain p-values from other test variables based on unbinned analyses [WILLIAMS 11, AAIJ 14C]. Results can be combined using Fisher's method [MOSTELLER 48].

### Local CPV in $D^\pm \rightarrow \pi^+ \pi^- \pi^\pm$

| p-value (%) | EVTS | DOCUMENT ID       | TECN     | COMMENT  |
|-------------|------|-------------------|----------|----------|
| <b>78.1</b> | 3.1M | <sup>1</sup> AAIJ | 14C LHCB | $\chi^2$ |

<sup>1</sup> AAIJ 14C uses binned and unbinned methods, and finds slightly better sensitivity with the former. We took the first value in the table of results for the binned method.

### Local CPV in $D^\pm \rightarrow K^+ K^- \pi^\pm$

| p-value (%)              | EVTS | DOCUMENT ID       | TECN     | COMMENT  |
|--------------------------|------|-------------------|----------|----------|
| <b>31 OUR EVALUATION</b> |      |                   |          |          |
| 72                       | 224k | LEES              | 13F BABR | $\chi^2$ |
| 12.7                     | 370k | <sup>1</sup> AAIJ | 11G LHCB | $\chi^2$ |

<sup>1</sup> AAIJ 11G publishes results for several binning schemes. We picked the first value in their table of results.

## CP VIOLATING ASYMMETRIES OF P-ODD (T-ODD) MOMENTS

### $A_{Tviol}(K_S^0 K^\pm \pi^+ \pi^-)$ in $D^\pm \rightarrow K_S^0 K^\pm \pi^+ \pi^-$

$C_T \equiv \vec{p}_{K^+} \cdot (\vec{p}_{\pi^+} \times \vec{p}_{\pi^-})$  is a parity-odd correlation of the  $K^+$ ,  $\pi^+$ , and  $\pi^-$  momenta for the  $D^+$ .  $\bar{C}_T \equiv \vec{p}_{K^-} \cdot (\vec{p}_{\pi^-} \times \vec{p}_{\pi^+})$  is the corresponding quantity for the  $D^-$ . Then

$$\frac{A_T}{\bar{A}_T} \equiv \frac{[\Gamma(C_T > 0) - \Gamma(C_T < 0)]}{[\Gamma(C_T > 0) + \Gamma(C_T < 0)]}, \text{ and}$$

$$\frac{\bar{A}_T}{A_T} \equiv \frac{[\Gamma(-\bar{C}_T > 0) - \Gamma(-\bar{C}_T < 0)]}{[\Gamma(-\bar{C}_T > 0) + \Gamma(-\bar{C}_T < 0)]}, \text{ and}$$

$A_{Tviol} \equiv \frac{1}{2}(A_T - \bar{A}_T)$ .  $C_T$  and  $\bar{C}_T$  are commonly referred to as  $T$ -odd moments, because they are odd under  $T$  reversal. However, the  $T$ -conjugate process  $K_S^0 K^\pm \pi^+ \pi^- \rightarrow D^\pm$  is not accessible, while the  $P$ -conjugate process is.

| VALUE (units $10^{-3}$ )  | EVTS            | DOCUMENT ID | TECN     | COMMENT  |
|---|-----------------|-------------|----------|--|
| <b><math>-12.0 \pm 10.0 \pm 4.6</math></b>                                    | $21.2 \pm 0.4k$ | LEES        | 11E BABR | $e^+ e^- \approx \Upsilon(4S)$                     |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                 |             |          |  |
| $23 \pm 62 \pm 22$  | $523 \pm 32$    | LINK        | 05E FOCS | $\gamma A, \bar{E}_\gamma \approx 180 \text{ GeV}$ |


### $D^+ \rightarrow (\bar{K}^0/\pi^0/\eta/\omega/\rho^0/\bar{K}^{*0})\ell^+\nu_\ell$ FORM FACTORS

$f_+(0)|V_{cs}|$  in  $D^+ \rightarrow \bar{K}^0 \ell^+ \nu_\ell$

| VALUE   | DOCUMENT ID   | TECN      | COMMENT                           |
|---|---|-----------|-----------------------------------|
| <b><math>0.719 \pm 0.011</math> OUR AVERAGE</b> | Error includes scale factor of 1.6. See the ideogram below. |           |                                   |
| $0.7053 \pm 0.0040 \pm 0.0112$                  | ABLIKIM   | 17S BES3  | $K_S^0 e^+ \nu_e$ 2-parameter fit |
| $0.737 \pm 0.006 \pm 0.009$                     | <sup>1</sup> ABLIKIM  | 15AF BES3 | $K_L e^+ \nu_e$ 3-parameter fit   |
| $0.707 \pm 0.010 \pm 0.009$                     | <sup>2</sup> BESSON   | 09 CLEO   | $K_S e^+ \nu_e$ 3-parameter fit   |

<sup>1</sup> ABLIKIM 15AF finds  $0.728 \pm 0.006 \pm 0.011$  for a 2-parameter fit.

<sup>2</sup> BESSON 09 finds  $0.716 \pm 0.007 \pm 0.009$  for a 2-parameter fit.



$r_1 \equiv a_1/a_0$  in  $D^+ \rightarrow \bar{K}^0 \ell^+ \nu_\ell$

| VALUE  | EVTS | DOCUMENT ID          | TECN      | COMMENT                           |
|--|------|----------------------|-----------|-----------------------------------|
| <b><math>-2.13 \pm 0.14</math> OUR AVERAGE</b>   |      |                      |           |                                   |
| $-2.18 \pm 0.14 \pm 0.05$  |      | ABLIKIM              | 17S BES3  | $K_S^0 e^+ \nu_e$ 2-parameter fit |
| $-2.23 \pm 0.42 \pm 0.53$  | 40k  | <sup>1</sup> ABLIKIM | 15AF BES3 | $K_L e^+ \nu_e$ 3-parameter fit   |
| $-1.66 \pm 0.44 \pm 0.10$  |      | <sup>2</sup> BESSON  | 09 CLEO   | $K_S e^+ \nu_e$ 3-parameter fit   |
| <sup>1</sup> ABLIKIM 15AF finds $r_1 = -1.91 \pm 0.33 \pm 0.28$ for a 2-parameter fit. |      |                      |           |                                   |
| <sup>2</sup> BESSON 09 finds $r_1 = -2.10 \pm 0.25 \pm 0.08$ for 2-parameter fit.      |      |                      |           |                                   |

$r_2 \equiv a_2/a_0$  in  $D^+ \rightarrow \bar{K}^0 \ell^+ \nu_\ell$

| VALUE   | EVTS | DOCUMENT ID | TECN      | COMMENT                         |
|---|------|-------------|-----------|---------------------------------|
| <b><math>-3 \pm 12</math> OUR AVERAGE</b> Error includes scale factor of 1.5. |      |             |           |                                 |
| $+11 \pm 9 \pm 9$   | 40k  | ABLIKIM     | 15AF BES3 | $K_L e^+ \nu_e$ 3-parameter fit |
| $-14 \pm 11 \pm 1$  |      | BESSON      | 09 CLEO   | $K_S e^+ \nu_e$ 3-parameter fit |

$f_+(0)|V_{cd}|$  in  $D^+ \rightarrow \pi^0 \ell^+ \nu_\ell$

| VALUE   | DOCUMENT ID | TECN     | COMMENT                           |
|---|-------------|----------|-----------------------------------|
| <b><math>0.1407 \pm 0.0025</math> OUR AVERAGE</b> |             |          |                                   |
| $0.1400 \pm 0.0026 \pm 0.0007$                    | ABLIKIM     | 17S BES3 | $\pi^0 e^+ \nu_e$ 2-parameter fit |
| $0.146 \pm 0.007 \pm 0.002$                       | BESSON      | 09 CLEO  | $\pi^0 e^+ \nu_e$ 3-parameter fit |

$r_1 \equiv a_1/a_0$  in  $D^+ \rightarrow \pi^0 \ell^+ \nu_\ell$

| VALUE  | DOCUMENT ID | TECN     | COMMENT                           |
|--|-------------|----------|-----------------------------------|
| <b><math>-2.00 \pm 0.13</math> OUR AVERAGE</b> |             |          |                                   |
| $-2.01 \pm 0.13 \pm 0.02$                      | ABLIKIM     | 17S BES3 | $\pi^0 e^+ \nu_e$ 2-parameter fit |
| $-1.37 \pm 0.88 \pm 0.24$                      | BESSON      | 09 CLEO  | $\pi^0 e^+ \nu_e$ 3-parameter fit |

$r_2 \equiv a_2/a_0$  in  $D^+ \rightarrow \pi^0 \ell^+ \nu_\ell$

| VALUE                              | DOCUMENT ID | TECN    | COMMENT                           |
|------------------------------------|-------------|---------|-----------------------------------|
| <b><math>-4 \pm 5 \pm 1</math></b> | BESSON      | 09 CLEO | $\pi^0 e^+ \nu_e$ 3-parameter fit |

$f_+(0)|V_{cd}|$  in  $D^+ \rightarrow \eta e^+ \nu_e$

| VALUE (units $10^{-2}$ )                    | EVTS | DOCUMENT ID | TECN     | COMMENT     |
|---|------|-------------|----------|-------------|
| <b><math>8.3 \pm 0.5</math> OUR AVERAGE</b> |      |             |          |             |
| $7.86 \pm 0.64 \pm 0.21$                    | 373  | ABLIKIM     | 18R BES3 | z expansion |
| $8.6 \pm 0.6 \pm 0.1$                       |      | YELTON      | 11 CLEO  | z expansion |

$r_1 \equiv a_1/a_0$  in  $D^+ \rightarrow \eta e^+ \nu_e$

| VALUE  | EVTS | DOCUMENT ID | TECN     | COMMENT     |
|--|------|-------------|----------|-------------|
| <b><math>-5.3 \pm 2.7</math> OUR AVERAGE</b> Error includes scale factor of 1.9. |      |             |          |             |
| $-7.33 \pm 1.69 \pm 0.40$  | 373  | ABLIKIM     | 18R BES3 | z expansion |
| $-1.83 \pm 2.23 \pm 0.28$  |      | YELTON      | 11 CLEO  | z expansion |

$r_V \equiv V(0)/A_1(0)$  in  $D^+ \rightarrow \omega e^+ \nu_e$

| VALUE                                      | DOCUMENT ID | TECN     | COMMENT                          |
|--|-------------|----------|----------------------------------|
| <b><math>1.24 \pm 0.09 \pm 0.06</math></b> | ABLIKIM     | 15W BES3 | $292 \text{ fb}^{-1}$ , 3773 MeV |

$r_2 \equiv A_2(0)/A_1(0)$  in  $D^+ \rightarrow \omega e^+ \nu_e$

| VALUE                                      | DOCUMENT ID | TECN     | COMMENT                          |
|--|-------------|----------|----------------------------------|
| <b><math>1.06 \pm 0.15 \pm 0.05</math></b> | ABLIKIM     | 15W BES3 | $292 \text{ fb}^{-1}$ , 3773 MeV |

$r_V \equiv V(0)/A_1(0)$  in  $D^+, D^0 \rightarrow \rho e^+ \nu_e$

| VALUE                     | DOCUMENT ID        | TECN | COMMENT                        |
|---------------------------|--------------------|------|--------------------------------|
| <b>1.48 ± 0.15 ± 0.05</b> | <sup>1</sup> DOBBS | 13   | CLEO $e^+ e^-$ at $\psi(3770)$ |

<sup>1</sup> Uses both  $D^+$  and  $D^0$  events. Using PDG 10 values of  $V_{cd}$  and lifetimes, DOBBS 13 gets  $A_1(0) = 0.56 \pm 0.01^{+0.02}_{-0.03}$ ,  $A_2(0) = 0.47 \pm 0.06 \pm 0.04$ , and  $V(0) = 0.84 \pm 0.09^{+0.05}_{-0.06}$ .

$r_2 \equiv A_2(0)/A_1(0)$  in  $D^+, D^0 \rightarrow \rho e^+ \nu_e$

| VALUE                     | DOCUMENT ID        | TECN | COMMENT                        |
|---------------------------|--------------------|------|--------------------------------|
| <b>0.83 ± 0.11 ± 0.04</b> | <sup>1</sup> DOBBS | 13   | CLEO $e^+ e^-$ at $\psi(3770)$ |

<sup>1</sup> Uses both  $D^+$  and  $D^0$  events. Using PDG 10 values of  $V_{cd}$  and lifetimes, DOBBS 13 gets  $A_1(0) = 0.56 \pm 0.01^{+0.02}_{-0.03}$ ,  $A_2(0) = 0.47 \pm 0.06 \pm 0.04$ , and  $V(0) = 0.84 \pm 0.09^{+0.05}_{-0.06}$ .

$r_V \equiv V(0)/A_1(0)$  in  $D^+ \rightarrow \bar{K}^*(892)^0 \ell^+ \nu_\ell$

See also BRIERE 10 for  $\bar{K}^* \ell^+ \nu_\ell$  helicity-basis form-factor measurements.

| VALUE   | EVTS  | DOCUMENT ID                         | TECN | COMMENT                               |
|---|-------|-------------------------------------|------|---------------------------------------|
| <b>1.49 ± 0.05 OUR AVERAGE</b>                |       | Error includes scale factor of 2.1. |      | See the ideogram below.               |
| 1.411 ± 0.058 ± 0.007                         | 16.2k | ABLIKIM                             | 16F  | BES3 $\bar{K}^*(892)^0 e^+ \nu_e$     |
| 1.463 ± 0.017 ± 0.031                         |       | <sup>1</sup> DEL-AMO-SA..11I        |      | BABR                                  |
| 1.504 ± 0.057 ± 0.039                         | 15k   | <sup>2</sup> LINK                   | 02L  | FOCS $\bar{K}^*(892)^0 \mu^+ \nu_\mu$ |
| 1.45 ± 0.23 ± 0.07                            | 763   | ADAMOVICH                           | 99   | BEAT $\bar{K}^*(892)^0 \mu^+ \nu_\mu$ |
| 1.90 ± 0.11 ± 0.09                            | 3000  | <sup>3</sup> AITALA                 | 98B  | E791 $\bar{K}^*(892)^0 e^+ \nu_e$     |
| 1.84 ± 0.11 ± 0.09                            | 3034  | AITALA                              | 98F  | E791 $\bar{K}^*(892)^0 \mu^+ \nu_\mu$ |
| 1.74 ± 0.27 ± 0.28                            | 874   | FRABETTI                            | 93E  | E687 $\bar{K}^*(892)^0 \mu^+ \nu_\mu$ |
| 2.00 <sup>+0.34</sup> <sub>-0.32</sub> ± 0.16 | 305   | KODAMA                              | 92   | E653 $\bar{K}^*(892)^0 \mu^+ \nu_\mu$ |


• • • We do not use the following data for averages, fits, limits, etc. • • •

|                 |     |       |     |                                   |
|-----------------|-----|-------|-----|-----------------------------------|
| 2.0 ± 0.6 ± 0.3 | 183 | ANJOS | 90E | E691 $\bar{K}^*(892)^0 e^+ \nu_e$ |
|-----------------|-----|-------|-----|-----------------------------------|

<sup>1</sup> DEL-AMO-SANCHEZ 11I finds the pole mass  $m_A = (2.63 \pm 0.10 \pm 0.13)$  GeV ( $m_V$  is fixed at 2 GeV).

<sup>2</sup> LINK 02L includes the effects of interference with an  $S$ -wave background. This much improves the goodness of fit, but does not much shift the values of the form factors.

<sup>3</sup> This is slightly different from the AITALA 98B value: see ref. [5] in AITALA 98F.



$$r_V \equiv V(0)/A_1(0) \text{ in } D^+ \rightarrow \bar{K}^*(892)^0 \ell^+ \nu_\ell$$

$$r_2 \equiv A_2(0)/A_1(0) \text{ in } D^+ \rightarrow \bar{K}^*(892)^0 \ell^+ \nu_\ell$$

See also BRIERE 10 for  $\bar{K}^* \ell^+ \nu_\ell$  helicity-basis form-factor measurements.

| VALUE                               | EVTS               | DOCUMENT ID                   | TECN     | COMMENT                          |
|-------------------------------------|--------------------|-------------------------------|----------|----------------------------------|
| <b><math>0.802 \pm 0.021</math></b> | <b>OUR AVERAGE</b> |                               |          |                                  |
| $0.788 \pm 0.042 \pm 0.008$         | 16.2k              | ABLIKIM                       | 16F BES3 | $\bar{K}^*(892)^0 e^+ \nu_e$     |
| $0.801 \pm 0.020 \pm 0.020$         |                    | <sup>1</sup> DEL-AMO-SA...11I | BABR     |                                  |
| $0.875 \pm 0.049 \pm 0.064$         | 15k                | <sup>2</sup> LINK             | 02L FOCS | $\bar{K}^*(892)^0 \mu^+ \nu_\mu$ |
| $1.00 \pm 0.15 \pm 0.03$            | 763                | ADAMOVICH                     | 99 BEAT  | $\bar{K}^*(892)^0 \mu^+ \nu_\mu$ |
| $0.71 \pm 0.08 \pm 0.09$            | 3000               | AITALA                        | 98B E791 | $\bar{K}^*(892)^0 e^+ \nu_e$     |
| $0.75 \pm 0.08 \pm 0.09$            | 3034               | AITALA                        | 98F E791 | $\bar{K}^*(892)^0 \mu^+ \nu_\mu$ |
| $0.78 \pm 0.18 \pm 0.10$            | 874                | FRABETTI                      | 93E E687 | $\bar{K}^*(892)^0 \mu^+ \nu_\mu$ |
| $0.82^{+0.22}_{-0.23} \pm 0.11$     | 305                | KODAMA                        | 92 E653  | $\bar{K}^*(892)^0 \mu^+ \nu_\mu$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                       |     |       |          |                              |
|-----------------------|-----|-------|----------|------------------------------|
| $0.0 \pm 0.5 \pm 0.2$ | 183 | ANJOS | 90E E691 | $\bar{K}^*(892)^0 e^+ \nu_e$ |
|-----------------------|-----|-------|----------|------------------------------|

<sup>1</sup> DEL-AMO-SANCHEZ 11I finds the pole mass  $m_A = (2.63 \pm 0.10 \pm 0.13)$  GeV ( $m_V$  is fixed at 2 GeV).

<sup>2</sup> LINK 02L includes the effects of interference with an  $S$ -wave background. This much improves the goodness of fit, but does not much shift the values of the form factors.

$$r_3 \equiv A_3(0)/A_1(0) \text{ in } D^+ \rightarrow \bar{K}^*(892)^0 \ell^+ \nu_\ell$$

See also BRIERE 10 for  $\bar{K}^* \ell^+ \nu_\ell$  helicity-basis form-factor measurements.

| VALUE                                      | EVTS | DOCUMENT ID | TECN     | COMMENT                          |
|--|------|-------------|----------|----------------------------------|
| <b><math>0.04 \pm 0.33 \pm 0.29</math></b> | 3034 | AITALA      | 98F E791 | $\bar{K}^*(892)^0 \mu^+ \nu_\mu$ |

### $\Gamma_L/\Gamma_T$ in $D^+ \rightarrow \bar{K}^*(892)^0 \ell^+ \nu_\ell$

See also BRIERE 10 for  $\bar{K}^* \ell^+ \nu_\ell$  helicity-basis form-factor measurements.

| VALUE   | EVTS | DOCUMENT ID  | TECN     | COMMENT                            |
|---|------|--------------|----------|------------------------------------|
| <b>1.13±0.08 OUR AVERAGE</b>  |      |              |          |                                    |
| 1.09±0.10±0.02  | 763  | ADAMOVICH 99 | BEAT     | $\bar{K}^*(892)^0_{\mu^+ \nu_\mu}$ |
| 1.20±0.13±0.13  | 874  | FRABETTI 93E | E687     | $\bar{K}^*(892)^0_{\mu^+ \nu_\mu}$ |
| 1.18±0.18±0.08  | 305  | KODAMA 92    | E653     | $\bar{K}^*(892)^0_{\mu^+ \nu_\mu}$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |      |              |          |                                    |
| 1.8 <sup>+0.6</sup> <sub>-0.4</sub> ±0.3                                      | 183  | ANJOS        | 90E E691 | $\bar{K}^*(892)^0_{e^+ \nu_e}$     |

### $\Gamma_+/\Gamma_-$ in $D^+ \rightarrow \bar{K}^*(892)^0 \ell^+ \nu_\ell$

See also BRIERE 10 for  $\bar{K}^* \ell^+ \nu_\ell$  helicity-basis form-factor measurements.

| VALUE   | EVTS | DOCUMENT ID  | TECN     | COMMENT                            |
|---|------|--------------|----------|------------------------------------|
| <b>0.22±0.06 OUR AVERAGE</b> Error includes scale factor of 1.6.              |      |              |          |                                    |
| 0.28±0.05±0.02  | 763  | ADAMOVICH 99 | BEAT     | $\bar{K}^*(892)^0_{\mu^+ \nu_\mu}$ |
| 0.16±0.05±0.02  | 305  | KODAMA 92    | E653     | $\bar{K}^*(892)^0_{\mu^+ \nu_\mu}$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |      |              |          |                                    |
| 0.15 <sup>+0.07</sup> <sub>-0.05</sub> ±0.03                                  | 183  | ANJOS        | 90E E691 | $\bar{K}^*(892)^0_{e^+ \nu_e}$     |

## $D^\pm$ REFERENCES

|                   |                       |                                  |                   |
|-------------------|-----------------------|----------------------------------|-------------------|
| ABLIKIM 18AC      | PR D98 092009         | M. Ablikim <i>et al.</i>         | (BES III Collab.) |
| ABLIKIM 18AE      | PRL 121 171803        | M. Ablikim <i>et al.</i>         | (BES III Collab.) |
| ABLIKIM 18F       | PRL 121 081802        | M. Ablikim <i>et al.</i>         | (BES III Collab.) |
| ABLIKIM 18P       | PR D97 072015         | M. Ablikim <i>et al.</i>         | (BES III Collab.) |
| ABLIKIM 18R       | PR D97 092009         | M. Ablikim <i>et al.</i>         | (BES III Collab.) |
| ABLIKIM 18W       | PR D97 072004         | M. Ablikim <i>et al.</i>         | (BES III Collab.) |
| BABU 18           | PR D97 011101         | V. Babu <i>et al.</i>            | (BELLE Collab.)   |
| AAIJ 17AF         | PL B771 21            | R. Aaij <i>et al.</i>            | (LHCb Collab.)    |
| ABLIKIM 17A       | PL B765 231           | M. Ablikim <i>et al.</i>         | (BES III Collab.) |
| ABLIKIM 17AD      | PR D96 092002         | M. Ablikim <i>et al.</i>         | (BES III Collab.) |
| ABLIKIM 17M       | PR D95 071102         | M. Ablikim <i>et al.</i>         | (BES III Collab.) |
| ABLIKIM 17S       | PR D96 012002         | M. Ablikim <i>et al.</i>         | (BES III Collab.) |
| ABLIKIM 16D       | PRL 116 082001        | M. Ablikim <i>et al.</i>         | (BES III Collab.) |
| ABLIKIM 16F       | PR D94 032001         | M. Ablikim <i>et al.</i>         | (BES III Collab.) |
| ABLIKIM 16G       | EPJ C76 369           | M. Ablikim <i>et al.</i>         | (BES III Collab.) |
| ABLIKIM 16V       | CP C40 113001         | M. Ablikim <i>et al.</i>         | (BES III Collab.) |
| ABLIKIM 15AF      | PR D92 112008         | M. Ablikim <i>et al.</i>         | (BES III Collab.) |
| ABLIKIM 15W       | PR D92 071101         | M. Ablikim <i>et al.</i>         | (BES III Collab.) |
| AAIJ 14BD         | JHEP 1410 025         | R. Aaij <i>et al.</i>            | (LHCb Collab.)    |
| AAIJ 14C          | PL B728 585           | R. Aaij <i>et al.</i>            | (LHCb Collab.)    |
| ABAZOV 14L        | PR D90 111102         | V.M. Abazov <i>et al.</i>        | (D0 Collab.)      |
| ABLIKIM 14E       | PR D89 052001         | M. Ablikim <i>et al.</i>         | (BES III Collab.) |
| ABLIKIM 14F       | PR D89 051104         | M. Ablikim <i>et al.</i>         | (BES III Collab.) |
| BONVICINI 14      | PR D89 072002         | G. Bonvicini <i>et al.</i>       | (CLEO Collab.)    |
| AAIJ 13AF         | PL B724 203           | R. Aaij <i>et al.</i>            | (LHCb Collab.)    |
| AAIJ 13W          | JHEP 1306 112         | R. Aaij <i>et al.</i>            | (LHCb Collab.)    |
| DOBBS 13          | PRL 110 131802        | S. Dobbs <i>et al.</i>           | (CLEO Collab.)    |
| KO 13             | JHEP 1302 098         | B.R. Ko <i>et al.</i>            | (BELLE Collab.)   |
| LEES 13E          | PR D87 052012         | J.P. Lees <i>et al.</i>          | (BABAR Collab.)   |
| LEES 13F          | PR D87 052010         | J.P. Lees <i>et al.</i>          | (BABAR Collab.)   |
| KO 12A            | PRL 109 119903 (err.) | B.R. Ko <i>et al.</i>            | (BELLE Collab.)   |
| Also              | PRL 109 021601        | B.R. Ko <i>et al.</i>            | (BELLE Collab.)   |
| STARIC 12         | PRL 108 071801        | M. Staric <i>et al.</i>          | (BELLE Collab.)   |
| AAIJ 11G          | PR D84 112008         | R. Aaij <i>et al.</i>            | (LHCb Collab.)    |
| DEL-AMO-SA... 11H | PR D83 071103         | P. del Amo Sanchez <i>et al.</i> | (BABAR Collab.)   |
| DEL-AMO-SA... 11I | PR D83 072001         | P. del Amo Sanchez <i>et al.</i> | (BABAR Collab.)   |
| LEES 11E          | PR D84 031103         | J.P. Lees <i>et al.</i>          | (BABAR Collab.)   |



|                |      |                         |                                  |                        |
|----------------|------|-------------------------|----------------------------------|------------------------|
| LEES           | 11G  | PR D84 072006           | J.P. Lees <i>et al.</i>          | (BABAR Collab.)        |
| WILLIAMS       | 11   | PR D84 054015           | M. Williams                      | (LOIC)                 |
| WON            | 11   | PRL 107 221801          | E. Won <i>et al.</i>             | (BELLE Collab.)        |
| YELTON         | 11   | PR D84 032001           | J. Yelton <i>et al.</i>          | (CLEO Collab.)         |
| ANASHIN        | 10A  | PL B686 84              | V.V. Anashin <i>et al.</i>       | (VEPP-4M KEDR Collab.) |
| ASNER          | 10   | PR D81 052007           | D.M. Asner <i>et al.</i>         | (CLEO Collab.)         |
| BRIERE         | 10   | PR D81 112001           | R.A. Briere <i>et al.</i>        | (CLEO Collab.)         |
| KO             | 10   | PRL 104 181602          | B.R. Ko <i>et al.</i>            | (BELLE Collab.)        |
| MENDEZ         | 10   | PR D81 052013           | H. Mendez <i>et al.</i>          | (CLEO Collab.)         |
| PDG            | 10   | JP G37 075021           | K. Nakamura <i>et al.</i>        | (PDG Collab.)          |
| RUBIN          | 10   | PR D82 092007           | P. Rubin <i>et al.</i>           | (CLEO Collab.)         |
| BEDIAGA        | 09   | PR D80 096006           | I. Bediaga <i>et al.</i>         | (CBPF, NDAM)           |
| BESSON         | 09   | PR D80 032005           | D. Besson <i>et al.</i>          | (CLEO Collab.)         |
| Also           |      | PR D79 052010           | J.Y. Ge <i>et al.</i>            | (CLEO Collab.)         |
| KO             | 09   | PRL 102 221802          | B.R. Ko <i>et al.</i>            | (BELLE Collab.)        |
| LINK           | 09   | PL B681 14              | J.M. Link <i>et al.</i>          | (FNAL FOCUS Collab.)   |
| MITCHELL       | 09B  | PRL 102 081801          | R.E. Mitchell <i>et al.</i>      | (CLEO Collab.)         |
| WON            | 09   | PR D80 111101           | E. Won <i>et al.</i>             | (BELLE Collab.)        |
| ABAZOV         | 08D  | PRL 100 101801          | V.M. Abazov <i>et al.</i>        | (D0 Collab.)           |
| ABLIKIM        | 08L  | PL B665 16              | M. Ablikim <i>et al.</i>         | (BES Collab.)          |
| ARTUSO         | 08   | PR D77 092003           | M. Artuso <i>et al.</i>          | (CLEO Collab.)         |
| AUBERT         | 08AO | PR D78 051102           | B. Aubert <i>et al.</i>          | (BABAR Collab.)        |
| BONVICINI      | 08   | PR D77 091106           | G. Bonvicini <i>et al.</i>       | (CLEO Collab.)         |
| BONVICINI      | 08A  | PR D78 052001           | G. Bonvicini <i>et al.</i>       | (CLEO Collab.)         |
| DOBBS          | 08   | PR D77 112005           | S. Dobbs <i>et al.</i>           | (CLEO Collab.)         |
| Also           |      | PRL 100 251802          | D. Cronin-Hennessy <i>et al.</i> | (CLEO Collab.)         |
| EISENSTEIN     | 08   | PR D78 052003           | B.I. Eisenstein <i>et al.</i>    | (CLEO Collab.)         |
| HE             | 08   | PRL 100 091801          | Q. He <i>et al.</i>              | (CLEO Collab.)         |
| PDG            | 08   | PL B667 1               | C. Amsler <i>et al.</i>          | (PDG Collab.)          |
| RUBIN          | 08   | PR D78 072003           | P. Rubin <i>et al.</i>           | (CLEO Collab.)         |
| ABLIKIM        | 07   | PL B644 20              | M. Ablikim <i>et al.</i>         | (BES Collab.)          |
| ABLIKIM        | 07G  | PL B658 1               | M. Ablikim <i>et al.</i>         | (BES Collab.)          |
| BONVICINI      | 07   | PR D76 012001           | G. Bonvicini <i>et al.</i>       | (CLEO Collab.)         |
| DOBBS          | 07   | PR D76 112001           | S. Dobbs <i>et al.</i>           | (CLEO Collab.)         |
| LINK           | 07B  | PL B653 1               | J.M. Link <i>et al.</i>          | (FNAL FOCUS Collab.)   |
| ABLIKIM        | 06O  | EPJ C47 31              | M. Ablikim <i>et al.</i>         | (BES Collab.)          |
| ABLIKIM        | 06P  | EPJ C47 39              | M. Ablikim <i>et al.</i>         | (BES Collab.)          |
| ABLIKIM        | 06U  | PL B643 246             | M. Ablikim <i>et al.</i>         | (BES Collab.)          |
| ADAM           | 06A  | PRL 97 251801           | N.E. Adam <i>et al.</i>          | (CLEO Collab.)         |
| AITALA         | 06   | PR D73 032004           | E.M. Aitala <i>et al.</i>        | (FNAL E791 Collab.)    |
| Also           |      | PR D74 059901 (errata.) | E.M. Aitala <i>et al.</i>        | (FNAL E791 Collab.)    |
| AUBERT,B       | 06F  | PR D74 011107           | B. Aubert <i>et al.</i>          | (BABAR Collab.)        |
| DYTMAN         | 06   | PR D74 071102           | S.A. Dytman <i>et al.</i>        | (CLEO Collab.)         |
| HUANG          | 06B  | PR D74 112005           | G.S. Huang <i>et al.</i>         | (CLEO Collab.)         |
| LINK           | 06B  | PL B637 32              | J.M. Link <i>et al.</i>          | (FNAL FOCUS Collab.)   |
| RUBIN          | 06   | PRL 96 081802           | P. Rubin <i>et al.</i>           | (CLEO Collab.)         |
| RUBIN          | 06A  | PR D73 112005           | P. Rubin <i>et al.</i>           | (CLEO Collab.)         |
| ABLIKIM        | 05A  | PL B608 24              | M. Ablikim <i>et al.</i>         | (BES Collab.)          |
| ABLIKIM        | 05D  | PL B610 183             | M. Ablikim <i>et al.</i>         | (BES Collab.)          |
| ABLIKIM        | 05F  | PL B622 6               | M. Ablikim <i>et al.</i>         | (BES Collab.)          |
| ABLIKIM        | 05P  | PL B625 196             | M. Ablikim <i>et al.</i>         | (BES Collab.)          |
| ARTUSO         | 05A  | PRL 95 251801           | M. Artuso <i>et al.</i>          | (CLEO Collab.)         |
| AUBERT         | 05S  | PR D71 091101           | B. Aubert <i>et al.</i>          | (BABAR Collab.)        |
| HE             | 05   | PRL 95 121801           | Q. He <i>et al.</i>              | (CLEO Collab.)         |
| Also           |      | PRL 96 199903 (errata.) | Q. He <i>et al.</i>              | (CLEO Collab.)         |
| HE             | 05A  | PRL 95 221802           | Q. He <i>et al.</i>              | (CLEO Collab.)         |
| HUANG          | 05B  | PRL 95 181801           | G.S. Huang <i>et al.</i>         | (CLEO Collab.)         |
| KAYIS-TOPAK... | 05   | PL B626 24              | A. Kayis-Topaksu <i>et al.</i>   | (CERN CHORUS Collab.)  |
| LINK           | 05E  | PL B622 239             | J.M. Link <i>et al.</i>          | (FNAL FOCUS Collab.)   |
| LINK           | 05I  | PL B621 72              | J.M. Link <i>et al.</i>          | (FNAL FOCUS Collab.)   |
| ABLIKIM        | 04C  | PL B597 39              | M. Ablikim <i>et al.</i>         | (BEPC BES Collab.)     |
| ARMS           | 04   | PR D69 071102           | K. Arms <i>et al.</i>            | (CLEO Collab.)         |
| BONVICINI      | 04A  | PR D70 112004           | G. Bonvicini <i>et al.</i>       | (CLEO Collab.)         |
| LINK           | 04   | PL B585 200             | J.M. Link <i>et al.</i>          | (FNAL FOCUS Collab.)   |
| LINK           | 04E  | PL B598 33              | J.M. Link <i>et al.</i>          | (FNAL FOCUS Collab.)   |
| LINK           | 04F  | PL B601 10              | J.M. Link <i>et al.</i>          | (FNAL FOCUS Collab.)   |
| ANISOVICH      | 03   | EPJ A16 229             | V.V. Anisovich <i>et al.</i>     |                        |
| LINK           | 03D  | PL B561 225             | J.M. Link <i>et al.</i>          | (FNAL FOCUS Collab.)   |
| LINK           | 03F  | PL B572 21              | J.M. Link <i>et al.</i>          | (FNAL FOCUS Collab.)   |
| AITALA         | 02   | PRL 89 121801           | E.M. Aitala <i>et al.</i>        | (FNAL E791 Collab.)    |
| BRANDENB...    | 02   | PRL 89 222001           | G. Brandenburg <i>et al.</i>     | (CLEO Collab.)         |

|                 |                        |                                |                         |
|-----------------|------------------------|--------------------------------|-------------------------|
| KAYIS-TOPAK..02 | PL B549 48             | A. Kayis-Topaksu <i>et al.</i> | (CERN CHORUS Collab.)   |
| LINK            | 02B PRL 88 041602      | J.M. Link <i>et al.</i>        | (FNAL FOCUS Collab.)    |
| Also            | PRL 88 159903 (errat.) | J.M. Link <i>et al.</i>        | (FNAL FOCUS Collab.)    |
| LINK            | 02E PL B535 43         | J.M. Link <i>et al.</i>        | (FNAL FOCUS Collab.)    |
| LINK            | 02F PL B537 192        | J.M. Link <i>et al.</i>        | (FNAL FOCUS Collab.)    |
| LINK            | 02I PL B541 227        | J.M. Link <i>et al.</i>        | (FNAL FOCUS Collab.)    |
| LINK            | 02J PL B541 243        | J.M. Link <i>et al.</i>        | (FNAL FOCUS Collab.)    |
| LINK            | 02L PL B544 89         | J.M. Link <i>et al.</i>        | (FNAL FOCUS Collab.)    |
| AITALA          | 01B PRL 86 770         | E.M. Aitala <i>et al.</i>      | (FNAL E791 Collab.)     |
| LINK            | 01C PRL 87 162001      | J.M. Link <i>et al.</i>        | (FNAL FOCUS Collab.)    |
| ABREU           | 000 EPJ C12 209        | P. Abreu <i>et al.</i>         | (DELPHI Collab.)        |
| ASTIER          | 00D PL B486 35         | P. Astier <i>et al.</i>        | (CERN NOMAD Collab.)    |
| JUN             | 00 PRL 84 1857         | S.Y. Jun <i>et al.</i>         | (FNAL SELEX Collab.)    |
| LINK            | 00B PL B491 232        | J.M. Link <i>et al.</i>        | (FNAL FOCUS Collab.)    |
| Also            | PL B495 443 (errat.)   | J.M. Link <i>et al.</i>        | (FNAL FOCUS Collab.)    |
| ABBIENDI        | 99K EPJ C8 573         | G. Abbiendi <i>et al.</i>      | (OPAL Collab.)          |
| ADAMOVICH       | 99 EPJ C6 35           | M. Adamovich <i>et al.</i>     | (CERN BEATRICE Collab.) |
| AITALA          | 99G PL B462 401        | E.M. Aitala <i>et al.</i>      | (FNAL E791 Collab.)     |
| BONVICINI       | 99 PRL 82 4586         | G. Bonvicini <i>et al.</i>     | (CLEO Collab.)          |
| AITALA          | 98B PRL 80 1393        | E.M. Aitala <i>et al.</i>      | (FNAL E791 Collab.)     |
| AITALA          | 98C PL B421 405        | E.M. Aitala <i>et al.</i>      | (FNAL E791 Collab.)     |
| AITALA          | 98F PL B440 435        | E.M. Aitala <i>et al.</i>      | (FNAL E791 Collab.)     |
| BAI             | 98B PL B429 188        | J.Z. Bai <i>et al.</i>         | (BEPV BES Collab.)      |
| AITALA          | 97 PL B397 325         | E.M. Aitala <i>et al.</i>      | (FNAL E791 Collab.)     |
| AITALA          | 97B PL B403 377        | E.M. Aitala <i>et al.</i>      | (FNAL E791 Collab.)     |
| AITALA          | 97C PL B404 187        | E.M. Aitala <i>et al.</i>      | (FNAL E791 Collab.)     |
| BISHAI          | 97 PRL 78 3261         | M. Bishai <i>et al.</i>        | (CLEO Collab.)          |
| FRABETTI        | 97 PL B391 235         | P.L. Frabetti <i>et al.</i>    | (FNAL E687 Collab.)     |
| FRABETTI        | 97B PL B398 239        | P.L. Frabetti <i>et al.</i>    | (FNAL E687 Collab.)     |
| FRABETTI        | 97C PL B401 131        | P.L. Frabetti <i>et al.</i>    | (FNAL E687 Collab.)     |
| FRABETTI        | 97D PL B407 79         | P.L. Frabetti <i>et al.</i>    | (FNAL E687 Collab.)     |
| AITALA          | 96 PRL 76 364          | E.M. Aitala <i>et al.</i>      | (FNAL E791 Collab.)     |
| FRABETTI        | 95 PL B346 199         | P.L. Frabetti <i>et al.</i>    | (FNAL E687 Collab.)     |
| FRABETTI        | 95B PL B351 591        | P.L. Frabetti <i>et al.</i>    | (FNAL E687 Collab.)     |
| FRABETTI        | 95E PL B359 403        | P.L. Frabetti <i>et al.</i>    | (FNAL E687 Collab.)     |
| KODAMA          | 95 PL B345 85          | K. Kodama <i>et al.</i>        | (FNAL E653 Collab.)     |
| ALBRECHT        | 94I ZPHY C64 375       | H. Albrecht <i>et al.</i>      | (ARGUS Collab.)         |
| BALEST          | 94 PRL 72 2328         | R. Balest <i>et al.</i>        | (CLEO Collab.)          |
| FRABETTI        | 94D PL B323 459        | P.L. Frabetti <i>et al.</i>    | (FNAL E687 Collab.)     |
| FRABETTI        | 94G PL B331 217        | P.L. Frabetti <i>et al.</i>    | (FNAL E687 Collab.)     |
| FRABETTI        | 94I PR D50 2953        | P.L. Frabetti <i>et al.</i>    | (FNAL E687 Collab.)     |
| AKERIB          | 93 PRL 71 3070         | D.S. Akerib <i>et al.</i>      | (CLEO Collab.)          |
| ANJOS           | 93 PR D48 56           | J.C. Anjos <i>et al.</i>       | (FNAL E691 Collab.)     |
| FRABETTI        | 93E PL B307 262        | P.L. Frabetti <i>et al.</i>    | (FNAL E687 Collab.)     |
| ALBRECHT        | 92F PL B278 202        | H. Albrecht <i>et al.</i>      | (ARGUS Collab.)         |
| ANJOS           | 92C PR D46 1941        | J.C. Anjos <i>et al.</i>       | (FNAL E691 Collab.)     |
| BARLAG          | 92C ZPHY C55 383       | S. Barlag <i>et al.</i>        | (ACCMOR Collab.)        |
| Also            | ZPHY C48 29            | S. Barlag <i>et al.</i>        | (ACCMOR Collab.)        |
| COFFMAN         | 92B PR D45 2196        | D.M. Coffman <i>et al.</i>     | (Mark III Collab.)      |
| DAOUDI          | 92 PR D45 3965         | M. Daoudi <i>et al.</i>        | (CLEO Collab.)          |
| KODAMA          | 92 PL B274 246         | K. Kodama <i>et al.</i>        | (FNAL E653 Collab.)     |
| KODAMA          | 92C PL B286 187        | K. Kodama <i>et al.</i>        | (FNAL E653 Collab.)     |
| ADAMOVICH       | 91 PL B268 142         | M.I. Adamovich <i>et al.</i>   | (WA82 Collab.)          |
| ALBRECHT        | 91 PL B255 634         | H. Albrecht <i>et al.</i>      | (ARGUS Collab.)         |
| ALVAREZ         | 91B ZPHY C50 11        | M.P. Alvarez <i>et al.</i>     | (CERN NA14/2 Collab.)   |
| AMMAR           | 91 PR D44 3383         | R. Ammar <i>et al.</i>         | (CLEO Collab.)          |
| BAI             | 91 PRL 66 1011         | Z. Bai <i>et al.</i>           | (Mark III Collab.)      |
| COFFMAN         | 91 PL B263 135         | D.M. Coffman <i>et al.</i>     | (Mark III Collab.)      |
| FRABETTI        | 91 PL B263 584         | P.L. Frabetti <i>et al.</i>    | (FNAL E687 Collab.)     |
| ALVAREZ         | 90 ZPHY C47 539        | M.P. Alvarez <i>et al.</i>     | (CERN NA14/2 Collab.)   |
| ANJOS           | 90C PR D41 2705        | J.C. Anjos <i>et al.</i>       | (FNAL E691 Collab.)     |
| ANJOS           | 90D PR D42 2414        | J.C. Anjos <i>et al.</i>       | (FNAL E691 Collab.)     |
| ANJOS           | 90E PRL 65 2630        | J.C. Anjos <i>et al.</i>       | (FNAL E691 Collab.)     |
| BARLAG          | 90C ZPHY C46 563       | S. Barlag <i>et al.</i>        | (ACCMOR Collab.)        |
| WEIR            | 90B PR D41 1384        | A.J. Weir <i>et al.</i>        | (Mark II Collab.)       |
| ANJOS           | 89 PRL 62 125          | J.C. Anjos <i>et al.</i>       | (FNAL E691 Collab.)     |
| ANJOS           | 89B PRL 62 722         | J.C. Anjos <i>et al.</i>       | (FNAL E691 Collab.)     |
| ANJOS           | 89E PL B223 267        | J.C. Anjos <i>et al.</i>       | (FNAL E691 Collab.)     |
| ADLER           | 88C PRL 60 89          | J. Adler <i>et al.</i>         | (Mark III Collab.)      |
| ALBRECHT        | 88I PL B210 267        | H. Albrecht <i>et al.</i>      | (ARGUS Collab.)         |
| HAAS            | 88 PRL 60 1614         | P. Haas <i>et al.</i>          | (CLEO Collab.)          |

|                 |     |                              |                                 |                           |
|-----------------|-----|------------------------------|---------------------------------|---------------------------|
| ONG             | 88  | PRL 60 2587                  | R.A. Ong <i>et al.</i>          | (Mark II Collab.)         |
| RAAB            | 88  | PR D37 2391                  | J.R. Raab <i>et al.</i>         | (FNAL E691 Collab.)       |
| ADAMOVICH       | 87  | EPL 4 887                    | M.I. Adamovich <i>et al.</i>    | (Photon Emulsion Collab.) |
| ADLER           | 87  | PL B196 107                  | J. Adler <i>et al.</i>          | (Mark III Collab.)        |
| BARTEL          | 87  | ZPHY C33 339                 | W. Bartel <i>et al.</i>         | (JADE Collab.)            |
| BALTRUSAITIS... | 86E | PRL 56 2140                  | R.M. Baltrusaitis <i>et al.</i> | (Mark III Collab.)        |
| BALTRUSAITIS... | 85B | PRL 54 1976                  | R.M. Baltrusaitis <i>et al.</i> | (Mark III Collab.)        |
| BALTRUSAITIS... | 85E | PRL 55 150                   | R.M. Baltrusaitis <i>et al.</i> | (Mark III Collab.)        |
| BARTEL          | 85J | PL 163B 277                  | W. Bartel <i>et al.</i>         | (JADE Collab.)            |
| ADAMOVICH       | 84  | PL 140B 119                  | M.I. Adamovich <i>et al.</i>    | (CERN WA58 Collab.)       |
| ALTHOFF         | 84G | ZPHY C22 219                 | M. Althoff <i>et al.</i>        | (TASSO Collab.)           |
| DERRICK         | 84  | PRL 53 1971                  | M. Derrick <i>et al.</i>        | (HRS Collab.)             |
| SCHINDLER       | 81  | PR D24 78                    | R.H. Schindler <i>et al.</i>    | (Mark II Collab.)         |
| TRILLING        | 81  | PRPL 75 57                   | G.H. Trilling                   | (LBL, UCB) J              |
| ZHOLENTZ        | 80  | PL 96B 214                   | A.A. Zholents <i>et al.</i>     | (NOVO)                    |
| Also            |     | SJNP 34 814                  | A.A. Zholents <i>et al.</i>     | (NOVO)                    |
|                 |     | Translated from YAF 34 1471. |                                 |                           |
| GOLDHABER       | 77  | PL 69B 503                   | G. Goldhaber <i>et al.</i>      | (Mark I Collab.)          |
| PERUZZI         | 77  | PRL 39 1301                  | I. Peruzzi <i>et al.</i>        | (LGW Collab.)             |
| PICCOLO         | 77  | PL 70B 260                   | M. Piccolo <i>et al.</i>        | (Mark I Collab.)          |
| PERUZZI         | 76  | PRL 37 569                   | I. Peruzzi <i>et al.</i>        | (Mark I Collab.)          |
| MOSTELLER       | 48  | Am.Stat. 3 No.5 30           | R.A. Fisher, F. Mosteller       |                           |

### OTHER RELATED PAPERS

|         |    |             |                            |              |
|---------|----|-------------|----------------------------|--------------|
| RICHMAN | 95 | RMP 67 893  | J.D. Richman, P.R. Burchat | (UCSB, STAN) |
| ROSNER  | 95 | CNPP 21 369 | J. Rosner                  | (CHIC)       |