

**$J/\psi(1S)$** 

$$J^G(J^{PC}) = 0^-(1^{--})$$

 **$J/\psi(1S)$  MASS**

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>3096.900±0.006 OUR AVERAGE</b>				
3096.900±0.002±0.006		<sup>1</sup> ANASHIN	15 KEDR	$e^+e^- \rightarrow \text{hadrons}$
3096.89 ±0.09	502	<sup>2</sup> ARTAMONOV	00 OLYA	$e^+e^- \rightarrow \text{hadrons}$
3096.91 ±0.03 ±0.01		<sup>3</sup> ARMSTRONG	93B E760	$\bar{p}p \rightarrow e^+e^-$
3096.95 ±0.1 ±0.3	193	BAGLIN	87 SPEC	$\bar{p}p \rightarrow e^+e^-X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
3096.66 ±0.19 ±0.02	6.1k	<sup>4</sup> AAIJ	15BI LHCb	$pp \rightarrow J/\psi X$
3096.917±0.010±0.007		AULCHENKO	03 KEDR	$e^+e^- \rightarrow \text{hadrons}$
3097.5 ±0.3		GRIBUSHIN	96 FMPS	515 $\pi^- \text{Be} \rightarrow 2\mu X$
3098.4 ±2.0	38k	LEMOIGNE	82 GOLI	185 $\pi^- \text{Be} \rightarrow \gamma\mu^+\mu^- A$
3096.93 ±0.09	502	<sup>5</sup> ZHOLENTZ	80 REDE	$e^+e^-$
3097.0 ±1		<sup>6</sup> BRANDELIK	79C DASP	$e^+e^-$

<sup>1</sup> Supersedes AULCHENKO 03.<sup>2</sup> Reanalysis of ZHOLENTZ 80 using new electron mass (COHEN 87) and radiative corrections (KURAEV 85).<sup>3</sup> Mass central value and systematic error recalculated by us according to Eq. (16) in ARMSTRONG 93B, using the value for the  $\psi(2S)$  mass from AULCHENKO 03.<sup>4</sup> From a sample of  $\eta_c(1S)$  and  $J/\psi$  produced in  $b$ -hadron decays. Systematic uncertainties not estimated.<sup>5</sup> Superseded by ARTAMONOV 00.<sup>6</sup> From a simultaneous fit to  $e^+e^-$ ,  $\mu^+\mu^-$  and hadronic channels assuming  $\Gamma(e^+e^-) = \Gamma(\mu^+\mu^-)$ . **$J/\psi(1S)$  WIDTH**

<u>VALUE (keV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>92.9 ± 2.8 OUR AVERAGE</b> Error includes scale factor of 1.1.				
96.1 ± 3.2	13k	<sup>1</sup> ADAMS	06A CLEO	$e^+e^- \rightarrow \mu^+\mu^-\gamma$
84.4 ± 8.9		BAI	95B BES	$e^+e^-$
91 ±11 ±6		<sup>2</sup> ARMSTRONG	93B E760	$\bar{p}p \rightarrow e^+e^-$
85.5 $\begin{smallmatrix} +6.1 \\ -5.8 \end{smallmatrix}$		<sup>3</sup> HSUEH	92 RVUE	See $\mathcal{T}$ mini-review
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
92.94± 1.83		<sup>4</sup> ANASHIN	18A KEDR	$e^+e^-$
94.1 ± 2.7		<sup>5</sup> ANASHIN	10 KEDR	3.097 $e^+e^- \rightarrow e^+e^-, \mu^+\mu^-$
93.7 ± 3.5	7.8k	<sup>1</sup> AUBERT	04 BABR	$e^+e^- \rightarrow \mu^+\mu^-\gamma$

<sup>1</sup> Calculated by us from the reported values of  $\Gamma(e^+e^-) \times B(\mu^+\mu^-)$  using  $B(e^+e^-) = (5.94 \pm 0.06)\%$  and  $B(\mu^+\mu^-) = (5.93 \pm 0.06)\%$ .

<sup>2</sup> The initial-state radiation correction reevaluated by ANDREOTTI 07 in its Ref. [4].

<sup>3</sup> Using data from COFFMAN 92, BALDINI-CELIO 75, BOYARSKI 75, ESPOSITO 75B, BRANDELIK 79C.

<sup>4</sup> Using  $\Gamma(e^+e^-)$  from ANASHIN 18A and  $B(J/\psi(1S) \rightarrow e^+e^-) = (5.971 \pm 0.032)\%$  from PDG 16.

<sup>5</sup> Assuming  $\Gamma(e^+e^-) = \Gamma(\mu^+\mu^-)$  and using  $\Gamma(e^+e^-)/\Gamma_{\text{total}} = (5.94 \pm 0.06)\%$ .

## $J/\psi(1S)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor/ Confidence level
$\Gamma_1$ hadrons	(87.7 ± 0.5 ) %	
$\Gamma_2$ virtual $\gamma \rightarrow$ hadrons	(13.50 ± 0.30 ) %	
$\Gamma_3$ $ggg$	(64.1 ± 1.0 ) %	
$\Gamma_4$ $\gamma gg$	( 8.8 ± 1.1 ) %	
$\Gamma_5$ $e^+e^-$	( 5.971 ± 0.032 ) %	
$\Gamma_6$ $e^+e^- \gamma$	[a] ( 8.8 ± 1.4 ) $\times 10^{-3}$	
$\Gamma_7$ $\mu^+\mu^-$	( 5.961 ± 0.033 ) %	

### Decays involving hadronic resonances

$\Gamma_8$ $\rho\pi$	( 1.69 ± 0.15 ) %	S=2.4
$\Gamma_9$ $\rho^0\pi^0$	( 5.6 ± 0.7 ) $\times 10^{-3}$	
$\Gamma_{10}$ $\rho(770)^\mp K^\pm K_S^0$	( 1.9 ± 0.4 ) $\times 10^{-3}$	
$\Gamma_{11}$ $\rho(1450)\pi$		
$\Gamma_{12}$ $\rho(1450)\pi \rightarrow \pi^+\pi^-\pi^0$	( 2.3 ± 0.7 ) $\times 10^{-3}$	
$\Gamma_{13}$ $\rho(1450)^\pm \pi^\mp \rightarrow K_S^0 K^\pm \pi^\mp$	( 3.5 ± 0.6 ) $\times 10^{-4}$	
$\Gamma_{14}$ $\rho(1450)^0 \pi^0 \rightarrow K^+ K^- \pi^0$	( 2.0 ± 0.5 ) $\times 10^{-4}$	
$\Gamma_{15}$ $\rho(1450)\eta'(958) \rightarrow \pi^+\pi^-\eta'(958)$	( 3.3 ± 0.7 ) $\times 10^{-6}$	
$\Gamma_{16}$ $\rho(1700)\pi$		
$\Gamma_{17}$ $\rho(1700)\pi \rightarrow \pi^+\pi^-\pi^0$	( 1.7 ± 1.1 ) $\times 10^{-4}$	
$\Gamma_{18}$ $\rho(2150)\pi$		
$\Gamma_{19}$ $\rho(2150)\pi \rightarrow \pi^+\pi^-\pi^0$	( 8 ± 40 ) $\times 10^{-6}$	
$\Gamma_{20}$ $\rho_3(1690)\pi \rightarrow \pi^+\pi^-\pi^0$		
$\Gamma_{21}$ $a_2(1320)\rho$	( 1.09 ± 0.22 ) %	
$\Gamma_{22}$ $\omega\pi^+\pi^+\pi^-\pi^-$	( 8.5 ± 3.4 ) $\times 10^{-3}$	
$\Gamma_{23}$ $\omega\pi^+\pi^-\pi^0$	( 4.0 ± 0.7 ) $\times 10^{-3}$	
$\Gamma_{24}$ $\omega\pi^+\pi^-$	( 7.2 ± 1.0 ) $\times 10^{-3}$	
$\Gamma_{25}$ $\omega f_2(1270)$	( 4.3 ± 0.6 ) $\times 10^{-3}$	
$\Gamma_{26}$ $K^*(892)^0 \bar{K}^*(892)^0$	( 2.3 ± 0.6 ) $\times 10^{-4}$	
$\Gamma_{27}$ $K^*(892)^\pm K^*(892)^\mp$	( 1.00 $\begin{smallmatrix} + \\ - \end{smallmatrix}$ $\begin{smallmatrix} 0.22 \\ 0.40 \end{smallmatrix}$ ) $\times 10^{-3}$	
$\Gamma_{28}$ $K^*(892)^\pm K^*(700)^\mp$	( 1.1 $\begin{smallmatrix} + \\ - \end{smallmatrix}$ $\begin{smallmatrix} 1.0 \\ 0.6 \end{smallmatrix}$ ) $\times 10^{-3}$	
$\Gamma_{29}$ $K_S^0 \pi^- K^*(892)^+ + \text{c.c.}$	( 2.0 ± 0.5 ) $\times 10^{-3}$	

$\Gamma_{30}$	$K_S^0 \pi^- K^*(892)^+ + \text{c.c.} \rightarrow$ $K_S^0 K_S^0 \pi^+ \pi^-$	$( 6.7 \pm 2.2 ) \times 10^{-4}$
$\Gamma_{31}$	$K_S^0 K^*(892)^0 \rightarrow \gamma K_S^0 K_S^0$	$( 6.3 \pm 0.6 ) \times 10^{-6}$
$\Gamma_{32}$	$\eta K^*(892)^0 \bar{K}^*(892)^0$	$( 1.15 \pm 0.26 ) \times 10^{-3}$
$\Gamma_{33}$	$\eta' K^{*\pm} K^\mp$	$( 1.48 \pm 0.13 ) \times 10^{-3}$
$\Gamma_{34}$	$\eta' K^{*0} \bar{K}^0 + \text{c.c.}$	$( 1.66 \pm 0.21 ) \times 10^{-3}$
$\Gamma_{35}$	$\eta' h_1(1415) \rightarrow \eta' K^* \bar{K} + \text{c.c.}$	$( 2.16 \pm 0.31 ) \times 10^{-4}$
$\Gamma_{36}$	$\eta' h_1(1415) \rightarrow \eta' K^{*\pm} K^\mp$	$( 1.51 \pm 0.23 ) \times 10^{-4}$
$\Gamma_{37}$	$K^*(1410) \bar{K} + \text{c.c.}$	
$\Gamma_{38}$	$K^*(1410) \bar{K} + \text{c.c.} \rightarrow$ $K^\pm K^\mp \pi^0$	$( 4.9 \pm 2.8 ) \times 10^{-5}$
$\Gamma_{39}$	$K^*(1410) \bar{K} + \text{c.c.} \rightarrow$ $K_S^0 K^\pm \pi^\mp$	$( 8 \pm 6 ) \times 10^{-5}$
$\Gamma_{40}$	$K_2^*(1430) \bar{K} + \text{c.c.}$	
$\Gamma_{41}$	$K_2^*(1430) \bar{K} + \text{c.c.} \rightarrow$ $K^\pm K^\mp \pi^0$	$( 7.5 \pm 3.5 ) \times 10^{-5}$
$\Gamma_{42}$	$K_2^*(1430) \bar{K} + \text{c.c.} \rightarrow$ $K_S^0 K^\pm \pi^\mp$	$( 4.0 \pm 1.0 ) \times 10^{-4}$
$\Gamma_{43}$	$K^*(892)^0 \bar{K}_2^*(1430)^0 + \text{c.c.}$	$( 4.66 \pm 0.31 ) \times 10^{-3}$
$\Gamma_{44}$	$K^*(892)^+ K_2^*(1430)^- + \text{c.c.}$	$( 3.4 \pm 2.9 ) \times 10^{-3}$
$\Gamma_{45}$	$K^*(892)^+ K_2^*(1430)^- + \text{c.c.} \rightarrow$ $K^*(892)^+ K_S^0 \pi^- + \text{c.c.}$	$( 4 \pm 4 ) \times 10^{-4}$
$\Gamma_{46}$	$K^*(892)^0 \bar{K}_2(1770)^0 + \text{c.c.} \rightarrow$ $K^*(892)^0 K^- \pi^+ + \text{c.c.}$	$( 6.9 \pm 0.9 ) \times 10^{-4}$
$\Gamma_{47}$	$\omega K^*(892) \bar{K} + \text{c.c.}$	$( 6.1 \pm 0.9 ) \times 10^{-3}$
$\Gamma_{48}$	$\bar{K} K^*(892) + \text{c.c.}$	
$\Gamma_{49}$	$\bar{K} K^*(892) + \text{c.c.} \rightarrow$ $K_S^0 K^\pm \pi^\mp$	$( 5.0 \pm 0.5 ) \times 10^{-3}$
$\Gamma_{50}$	$K^+ K^*(892)^- + \text{c.c.}$	$( 5.0 \pm 0.4 ) \times 10^{-3}$
$\Gamma_{51}$	$K^+ K^*(892)^- + \text{c.c.} \rightarrow$ $K^+ K^- \pi^0$	$( 1.98 \pm 0.21 ) \times 10^{-3}$
$\Gamma_{52}$	$K^+ K^*(892)^- + \text{c.c.} \rightarrow$ $K^0 K^\pm \pi^\mp + \text{c.c.}$	$( 3.0 \pm 0.4 ) \times 10^{-3}$
$\Gamma_{53}$	$K^0 \bar{K}^*(892)^0 + \text{c.c.}$	$( 4.2 \pm 0.4 ) \times 10^{-3}$
$\Gamma_{54}$	$K^0 \bar{K}^*(892)^0 + \text{c.c.} \rightarrow$ $K^0 K^\pm \pi^\mp + \text{c.c.}$	$( 3.2 \pm 0.4 ) \times 10^{-3}$
$\Gamma_{55}$	$K_1(1400)^\pm K^\mp$	$( 3.8 \pm 1.4 ) \times 10^{-3}$
$\Gamma_{56}$	$\bar{K}^*(892)^0 K^+ \pi^- + \text{c.c.}$	$( 7.7 \pm 1.6 ) \times 10^{-3}$
$\Gamma_{57}$	$K^*(892)^\pm K^\mp \pi^0$	$( 4.1 \pm 1.3 ) \times 10^{-3}$
$\Gamma_{58}$	$K^*(892)^0 K_S^0 \pi^0$	$( 6 \pm 4 ) \times 10^{-4}$
$\Gamma_{59}$	$\omega \pi^0 \pi^0$	$( 3.4 \pm 0.8 ) \times 10^{-3}$
$\Gamma_{60}$	$\omega \pi^0 \eta$	$( 3.4 \pm 1.7 ) \times 10^{-4}$
$\Gamma_{61}$	$b_1(1235)^\pm \pi^\mp$	[b] $( 3.0 \pm 0.5 ) \times 10^{-3}$

$\Gamma_{62}$	$\omega K^\pm K_S^0 \pi^\mp$	[b]	$( 3.4 \pm 0.5 ) \times 10^{-3}$	
$\Gamma_{63}$	$b_1(1235)^0 \pi^0$		$( 2.3 \pm 0.6 ) \times 10^{-3}$	
$\Gamma_{64}$	$\eta K^\pm K_S^0 \pi^\mp$	[b]	$( 2.2 \pm 0.4 ) \times 10^{-3}$	
$\Gamma_{65}$	$\phi K^*(892) \bar{K} + \text{c.c.}$		$( 2.18 \pm 0.23 ) \times 10^{-3}$	
$\Gamma_{66}$	$\omega K \bar{K}$		$( 1.9 \pm 0.4 ) \times 10^{-3}$	
$\Gamma_{67}$	$\omega f_0(1710) \rightarrow \omega K \bar{K}$		$( 4.8 \pm 1.1 ) \times 10^{-4}$	
$\Gamma_{68}$	$\phi 2(\pi^+ \pi^-)$		$( 1.60 \pm 0.32 ) \times 10^{-3}$	
$\Gamma_{69}$	$\Delta(1232)^{++} \bar{p} \pi^-$		$( 1.6 \pm 0.5 ) \times 10^{-3}$	
$\Gamma_{70}$	$\omega \eta$		$( 1.74 \pm 0.20 ) \times 10^{-3}$	S=1.6
$\Gamma_{71}$	$\phi K \bar{K}$		$( 1.77 \pm 0.16 ) \times 10^{-3}$	S=1.3
$\Gamma_{72}$	$\phi K_S^0 K_S^0$		$( 5.9 \pm 1.5 ) \times 10^{-4}$	
$\Gamma_{73}$	$\phi f_0(1710) \rightarrow \phi K \bar{K}$		$( 3.6 \pm 0.6 ) \times 10^{-4}$	
$\Gamma_{74}$	$\phi K^+ K^-$		$( 8.3 \pm 1.2 ) \times 10^{-4}$	
$\Gamma_{75}$	$\phi f_2(1270)$		$( 3.2 \pm 0.6 ) \times 10^{-4}$	
$\Gamma_{76}$	$\Delta(1232)^{++} \bar{\Delta}(1232)^{--}$		$( 1.10 \pm 0.29 ) \times 10^{-3}$	
$\Gamma_{77}$	$\Sigma(1385)^- \bar{\Sigma}(1385)^+ (\text{or c.c.})$	[b]	$( 1.16 \pm 0.05 ) \times 10^{-3}$	
$\Gamma_{78}$	$\Sigma(1385)^0 \bar{\Sigma}(1385)^0$		$( 1.07 \pm 0.08 ) \times 10^{-3}$	
$\Gamma_{79}$	$K^+ K^- f_2'(1525)$		$( 1.04 \pm 0.35 ) \times 10^{-3}$	
$\Gamma_{80}$	$\phi f_2'(1525)$		$( 8 \pm 4 ) \times 10^{-4}$	S=2.7
$\Gamma_{81}$	$\phi \pi^+ \pi^-$		$( 9.4 \pm 1.5 ) \times 10^{-4}$	S=1.7
$\Gamma_{82}$	$\phi \pi^0 \pi^0$		$( 5.0 \pm 1.0 ) \times 10^{-4}$	
$\Gamma_{83}$	$\phi K^\pm K_S^0 \pi^\mp$	[b]	$( 7.2 \pm 0.8 ) \times 10^{-4}$	
$\Gamma_{84}$	$\omega f_1(1420)$		$( 6.8 \pm 2.4 ) \times 10^{-4}$	
$\Gamma_{85}$	$\phi \eta$		$( 7.4 \pm 0.8 ) \times 10^{-4}$	S=1.5
$\Gamma_{86}$	$\Xi^0 \Xi^0$		$( 1.17 \pm 0.04 ) \times 10^{-3}$	
$\Gamma_{87}$	$\Xi(1530)^- \Xi^+$		$( 5.9 \pm 1.5 ) \times 10^{-4}$	
$\Gamma_{88}$	$p K^- \bar{\Sigma}(1385)^0$		$( 5.1 \pm 3.2 ) \times 10^{-4}$	
$\Gamma_{89}$	$\omega \pi^0$		$( 4.5 \pm 0.5 ) \times 10^{-4}$	S=1.4
$\Gamma_{90}$	$\omega \pi^0 \rightarrow \pi^+ \pi^- \pi^0$		$( 1.7 \pm 0.8 ) \times 10^{-5}$	
$\Gamma_{91}$	$\phi \eta'(958)$		$( 4.6 \pm 0.5 ) \times 10^{-4}$	S=2.2
$\Gamma_{92}$	$\phi f_0(980)$		$( 3.2 \pm 0.9 ) \times 10^{-4}$	S=1.9
$\Gamma_{93}$	$\phi f_0(980) \rightarrow \phi \pi^+ \pi^-$		$( 2.59 \pm 0.34 ) \times 10^{-4}$	
$\Gamma_{94}$	$\phi f_0(980) \rightarrow \phi \pi^0 \pi^0$		$( 1.8 \pm 0.5 ) \times 10^{-4}$	
$\Gamma_{95}$	$\phi \pi^0 f_0(980) \rightarrow \phi \pi^0 \pi^+ \pi^-$		$( 4.5 \pm 1.0 ) \times 10^{-6}$	
$\Gamma_{96}$	$\phi \pi^0 f_0(980) \rightarrow \phi \pi^0 p^0 \pi^0$		$( 1.7 \pm 0.6 ) \times 10^{-6}$	
$\Gamma_{97}$	$\eta \phi f_0(980) \rightarrow \eta \phi \pi^+ \pi^-$		$( 3.2 \pm 1.0 ) \times 10^{-4}$	
$\Gamma_{98}$	$\phi a_0(980)^0 \rightarrow \phi \eta \pi^0$		$( 4.4 \pm 1.4 ) \times 10^{-6}$	
$\Gamma_{99}$	$\Xi(1530)^0 \Xi^0$		$( 3.2 \pm 1.4 ) \times 10^{-4}$	
$\Gamma_{100}$	$\Sigma(1385)^- \bar{\Sigma}^+ (\text{or c.c.})$	[b]	$( 3.1 \pm 0.5 ) \times 10^{-4}$	
$\Gamma_{101}$	$\phi f_1(1285)$		$( 2.6 \pm 0.5 ) \times 10^{-4}$	
$\Gamma_{102}$	$\phi f_1(1285) \rightarrow \phi \pi^0 f_0(980) \rightarrow$ $\phi \pi^0 \pi^+ \pi^-$		$( 9.4 \pm 2.8 ) \times 10^{-7}$	
$\Gamma_{103}$	$\phi f_1(1285) \rightarrow \phi \pi^0 f_0(980) \rightarrow$ $\phi \pi^0 \pi^0 \pi^0$		$( 2.1 \pm 2.2 ) \times 10^{-7}$	

$\Gamma_{104}$	$\eta\pi^+\pi^-$	$(4.2 \pm 0.8) \times 10^{-4}$	
$\Gamma_{105}$	$\eta\rho$	$(1.93 \pm 0.23) \times 10^{-4}$	
$\Gamma_{106}$	$\omega\eta'(958)$	$(1.89 \pm 0.18) \times 10^{-4}$	
$\Gamma_{107}$	$\omega f_0(980)$	$(1.4 \pm 0.5) \times 10^{-4}$	
$\Gamma_{108}$	$\rho\eta'(958)$	$(8.1 \pm 0.8) \times 10^{-5}$	S=1.6
$\Gamma_{109}$	$a_2(1320)^\pm\pi^\mp$	[b] < 4.3	$\times 10^{-3}$ CL=90%
$\Gamma_{110}$	$K\bar{K}_2^*(1430) + \text{c.c.}$	< 4.0	$\times 10^{-3}$ CL=90%
$\Gamma_{111}$	$K_1(1270)^\pm K^\mp$	< 3.0	$\times 10^{-3}$ CL=90%
$\Gamma_{112}$	$K_1(1270)K_S^0 \rightarrow \gamma K_S^0 K_S^0$	$(8.5 \pm 2.5) \times 10^{-7}$	
$\Gamma_{113}$	$K_S^0\pi^- K_2^*(1430)^+ + \text{c.c.}$	$(3.6 \pm 1.8) \times 10^{-3}$	
$\Gamma_{114}$	$K_2^*(1430)^0\bar{K}_2^*(1430)^0$	< 2.9	$\times 10^{-3}$ CL=90%
$\Gamma_{115}$	$\phi\pi^0$	$3 \times 10^{-6}$ or $1 \times 10^{-7}$	
$\Gamma_{116}$	$\phi\eta(1405) \rightarrow \phi\eta\pi^+\pi^-$	$(2.0 \pm 1.0) \times 10^{-5}$	
$\Gamma_{117}$	$\omega f_2'(1525)$	< 2.2	$\times 10^{-4}$ CL=90%
$\Gamma_{118}$	$\omega X(1835) \rightarrow \omega p\bar{p}$	< 3.9	$\times 10^{-6}$ CL=95%
$\Gamma_{119}$	$\phi X(1835) \rightarrow \phi p\bar{p}$	< 2.1	$\times 10^{-7}$ CL=90%
$\Gamma_{120}$	$\phi X(1835) \rightarrow \phi\eta\pi^+\pi^-$	< 2.8	$\times 10^{-4}$ CL=90%
$\Gamma_{121}$	$\phi X(1870) \rightarrow \phi\eta\pi^+\pi^-$	< 6.13	$\times 10^{-5}$ CL=90%
$\Gamma_{122}$	$\eta\phi(2170) \rightarrow \eta\phi f_0(980) \rightarrow \eta\phi\pi^+\pi^-$	$(1.2 \pm 0.4) \times 10^{-4}$	
$\Gamma_{123}$	$\eta\phi(2170) \rightarrow \eta K^*(892)^0\bar{K}^*(892)^0$	< 2.52	$\times 10^{-4}$ CL=90%
$\Gamma_{124}$	$\Sigma(1385)^0\bar{\Lambda} + \text{c.c.}$	< 8.2	$\times 10^{-6}$ CL=90%
$\Gamma_{125}$	$\Delta(1232)^+\bar{p}$	< 1	$\times 10^{-4}$ CL=90%
$\Gamma_{126}$	$\Lambda(1520)\bar{\Lambda} + \text{c.c.} \rightarrow \gamma\Lambda\bar{\Lambda}$	< 4.1	$\times 10^{-6}$ CL=90%
$\Gamma_{127}$	$\bar{\Lambda}(1520)\Lambda + \text{c.c.}$	< 1.80	$\times 10^{-3}$ CL=90%
$\Gamma_{128}$	$\Theta(1540)\bar{\Theta}(1540) \rightarrow K_S^0 p K^- \bar{n} + \text{c.c.}$	< 1.1	$\times 10^{-5}$ CL=90%
$\Gamma_{129}$	$\Theta(1540)K^- \bar{n} \rightarrow K_S^0 p K^- \bar{n}$	< 2.1	$\times 10^{-5}$ CL=90%
$\Gamma_{130}$	$\Theta(1540)K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n$	< 1.6	$\times 10^{-5}$ CL=90%
$\Gamma_{131}$	$\bar{\Theta}(1540)K^+ n \rightarrow K_S^0 \bar{p} K^+ n$	< 5.6	$\times 10^{-5}$ CL=90%
$\Gamma_{132}$	$\bar{\Theta}(1540)K_S^0 p \rightarrow K_S^0 p K^- \bar{n}$	< 1.1	$\times 10^{-5}$ CL=90%

### Decays into stable hadrons

$\Gamma_{133}$	$2(\pi^+\pi^-)\pi^0$	$(3.37 \pm 0.26) \%$	
$\Gamma_{134}$	$3(\pi^+\pi^-)\pi^0$	$(2.9 \pm 0.6) \%$	
$\Gamma_{135}$	$\pi^+\pi^-\pi^0$	$(2.10 \pm 0.08) \%$	S=1.6
$\Gamma_{136}$	$\pi^+\pi^-\pi^0\pi^0\pi^0$	$(2.71 \pm 0.29) \%$	
$\Gamma_{137}$	$\rho^\pm\pi^\mp\pi^0\pi^0$	$(1.41 \pm 0.22) \%$	
$\Gamma_{138}$	$\rho^+\rho^-\pi^0$	$(6.0 \pm 1.1) \times 10^{-3}$	
$\Gamma_{139}$	$\pi^+\pi^-\pi^0 K^+ K^-$	$(1.20 \pm 0.30) \%$	
$\Gamma_{140}$	$4(\pi^+\pi^-)\pi^0$	$(9.0 \pm 3.0) \times 10^{-3}$	
$\Gamma_{141}$	$\pi^+\pi^- K^+ K^-$	$(6.84 \pm 0.32) \times 10^{-3}$	
$\Gamma_{142}$	$\pi^+\pi^- K_S^0 K_L^0$	$(3.8 \pm 0.6) \times 10^{-3}$	

$\Gamma_{143}$	$\pi^+\pi^-K_S^0K_S^0$	$(1.68 \pm 0.19) \times 10^{-3}$	
$\Gamma_{144}$	$\pi^\pm\pi^0K^\mp K_S^0$	$(5.7 \pm 0.5) \times 10^{-3}$	
$\Gamma_{145}$	$K^+K^-K_S^0K_S^0$	$(4.1 \pm 0.8) \times 10^{-4}$	
$\Gamma_{146}$	$\pi^+\pi^-K^+K^-\eta$	$(4.7 \pm 0.7) \times 10^{-3}$	
$\Gamma_{147}$	$\pi^0\pi^0K^+K^-$	$(2.12 \pm 0.23) \times 10^{-3}$	
$\Gamma_{148}$	$\pi^0\pi^0K_S^0K_L^0$	$(1.9 \pm 0.4) \times 10^{-3}$	
$\Gamma_{149}$	$K\bar{K}\pi$	$(6.1 \pm 1.0) \times 10^{-3}$	
$\Gamma_{150}$	$K^+K^-\pi^0$	$(2.14 \pm 0.24) \times 10^{-3}$	
$\Gamma_{151}$	$K_S^0K^\pm\pi^\mp$	$(5.6 \pm 0.5) \times 10^{-3}$	
$\Gamma_{152}$	$K_S^0K_L^0\pi^0$	$(2.06 \pm 0.27) \times 10^{-3}$	
$\Gamma_{153}$	$K^*(892)^0\bar{K}^0 + \text{c.c.} \rightarrow$ $K_S^0K_L^0\pi^0$	$(1.21 \pm 0.18) \times 10^{-3}$	
$\Gamma_{154}$	$K_2^*(1430)^0\bar{K}^0 + \text{c.c.} \rightarrow$ $K_S^0K_L^0\pi^0$	$(4.3 \pm 1.3) \times 10^{-4}$	
$\Gamma_{155}$	$K_S^0K_L^0\eta$	$(1.44 \pm 0.34) \times 10^{-3}$	
$\Gamma_{156}$	$2(\pi^+\pi^-)$	$(3.57 \pm 0.30) \times 10^{-3}$	
$\Gamma_{157}$	$3(\pi^+\pi^-)$	$(4.3 \pm 0.4) \times 10^{-3}$	
$\Gamma_{158}$	$2(\pi^+\pi^-\pi^0)$	$(1.61 \pm 0.21) \%$	
$\Gamma_{159}$	$2(\pi^+\pi^-\eta)$	$(2.26 \pm 0.28) \times 10^{-3}$	
$\Gamma_{160}$	$3(\pi^+\pi^-\eta)$	$(7.2 \pm 1.5) \times 10^{-4}$	
$\Gamma_{161}$	$\pi^+\pi^-\pi^0\pi^0\eta$	$(2.3 \pm 0.5) \times 10^{-3}$	
$\Gamma_{162}$	$\rho^\pm\pi^\mp\pi^0\eta$	$(1.9 \pm 0.8) \times 10^{-3}$	
$\Gamma_{163}$	$\rho\bar{\rho}$	$(2.121 \pm 0.029) \times 10^{-3}$	
$\Gamma_{164}$	$\rho\bar{\rho}\pi^0$	$(1.19 \pm 0.08) \times 10^{-3}$	S=1.1
$\Gamma_{165}$	$\rho\bar{\rho}\pi^+\pi^-$	$(6.0 \pm 0.5) \times 10^{-3}$	S=1.3
$\Gamma_{166}$	$\rho\bar{\rho}\pi^+\pi^-\pi^0$	[c] $(2.3 \pm 0.9) \times 10^{-3}$	S=1.9
$\Gamma_{167}$	$\rho\bar{\rho}\eta$	$(2.00 \pm 0.12) \times 10^{-3}$	
$\Gamma_{168}$	$\rho\bar{\rho}\rho$	< 3.1 $\times 10^{-4}$	CL=90%
$\Gamma_{169}$	$\rho\bar{\rho}\omega$	$(9.8 \pm 1.0) \times 10^{-4}$	S=1.3
$\Gamma_{170}$	$\rho\bar{\rho}\eta'(958)$	$(1.29 \pm 0.14) \times 10^{-4}$	S=2.0
$\Gamma_{171}$	$\rho\bar{\rho}a_0(980) \rightarrow \rho\bar{\rho}\pi^0\eta$	$(6.8 \pm 1.8) \times 10^{-5}$	
$\Gamma_{172}$	$\rho\bar{\rho}\phi$	$(5.19 \pm 0.33) \times 10^{-5}$	
$\Gamma_{173}$	$n\bar{n}$	$(2.09 \pm 0.16) \times 10^{-3}$	
$\Gamma_{174}$	$n\bar{n}\pi^+\pi^-$	$(4 \pm 4) \times 10^{-3}$	
$\Gamma_{175}$	$\Sigma^+\bar{\Sigma}^-$	$(1.50 \pm 0.24) \times 10^{-3}$	
$\Gamma_{176}$	$\Sigma^0\bar{\Sigma}^0$	$(1.172 \pm 0.032) \times 10^{-3}$	S=1.4
$\Gamma_{177}$	$2(\pi^+\pi^-)K^+K^-$	$(3.1 \pm 1.3) \times 10^{-3}$	
$\Gamma_{178}$	$\rho\bar{n}\pi^-$	$(2.12 \pm 0.09) \times 10^{-3}$	
$\Gamma_{179}$	$nN(1440)$	seen	
$\Gamma_{180}$	$nN(1520)$	seen	
$\Gamma_{181}$	$nN(1535)$	seen	
$\Gamma_{182}$	$\Xi^-\bar{\Xi}^+$	$(9.7 \pm 0.8) \times 10^{-4}$	S=1.4
$\Gamma_{183}$	$\Lambda\bar{\Lambda}$	$(1.89 \pm 0.09) \times 10^{-3}$	S=2.8

$\Gamma_{184}$	$\Lambda\bar{\Sigma}^-\pi^+$ (or c.c.)	[b]	$(8.3 \pm 0.7) \times 10^{-4}$	S=1.2
$\Gamma_{185}$	$\rho K^-\bar{\Lambda} + \text{c.c.}$		$(8.7 \pm 1.1) \times 10^{-4}$	
$\Gamma_{186}$	$2(K^+K^-)$		$(7.2 \pm 0.8) \times 10^{-4}$	
$\Gamma_{187}$	$\rho K^-\bar{\Sigma}^0$		$(2.9 \pm 0.8) \times 10^{-4}$	
$\Gamma_{188}$	$K^+K^-$		$(2.86 \pm 0.21) \times 10^{-4}$	
$\Gamma_{189}$	$K_S^0 K_L^0$		$(1.95 \pm 0.11) \times 10^{-4}$	S=2.4
$\Gamma_{190}$	$\Lambda\bar{\Lambda}\pi^+\pi^-$		$(4.3 \pm 1.0) \times 10^{-3}$	
$\Gamma_{191}$	$\Lambda\bar{\Lambda}\eta$		$(1.62 \pm 0.17) \times 10^{-4}$	
$\Gamma_{192}$	$\Lambda\bar{\Lambda}\pi^0$		$(3.8 \pm 0.4) \times 10^{-5}$	
$\Gamma_{193}$	$\bar{\Lambda}nK_S^0 + \text{c.c.}$		$(6.5 \pm 1.1) \times 10^{-4}$	
$\Gamma_{194}$	$\pi^+\pi^-$		$(1.47 \pm 0.14) \times 10^{-4}$	
$\Gamma_{195}$	$\Lambda\bar{\Sigma} + \text{c.c.}$		$(2.83 \pm 0.23) \times 10^{-5}$	
$\Gamma_{196}$	$K_S^0 K_S^0$		$< 1.4 \times 10^{-8}$	CL=95%

### Radiative decays

$\Gamma_{197}$	$3\gamma$		$(1.16 \pm 0.22) \times 10^{-5}$	
$\Gamma_{198}$	$4\gamma$		$< 9 \times 10^{-6}$	CL=90%
$\Gamma_{199}$	$5\gamma$		$< 1.5 \times 10^{-5}$	CL=90%
$\Gamma_{200}$	$\gamma\pi^0\pi^0$		$(1.15 \pm 0.05) \times 10^{-3}$	
$\Gamma_{201}$	$\gamma\eta\pi^0$		$(2.14 \pm 0.31) \times 10^{-5}$	
$\Gamma_{202}$	$\gamma a_0(980)^0 \rightarrow \gamma\eta\pi^0$		$< 2.5 \times 10^{-6}$	CL=95%
$\Gamma_{203}$	$\gamma a_2(1320)^0 \rightarrow \gamma\eta\pi^0$		$< 6.6 \times 10^{-6}$	CL=95%
$\Gamma_{204}$	$\gamma K_S^0 K_S^0$		$(8.1 \pm 0.4) \times 10^{-4}$	
$\Gamma_{205}$	$\gamma\eta_c(1S)$		$(1.7 \pm 0.4) \%$	S=1.5
$\Gamma_{206}$	$\gamma\eta_c(1S) \rightarrow 3\gamma$		$(3.8 \pm 1.3 \pm 1.0) \times 10^{-6}$	S=1.1
$\Gamma_{207}$	$\gamma\pi^+\pi^-2\pi^0$		$(8.3 \pm 3.1) \times 10^{-3}$	
$\Gamma_{208}$	$\gamma\eta\pi\pi$		$(6.1 \pm 1.0) \times 10^{-3}$	
$\Gamma_{209}$	$\gamma\eta_2(1870) \rightarrow \gamma\eta\pi^+\pi^-$		$(6.2 \pm 2.4) \times 10^{-4}$	
$\Gamma_{210}$	$\gamma\eta(1405/1475) \rightarrow \gamma K\bar{K}\pi$	[d]	$(2.8 \pm 0.6) \times 10^{-3}$	S=1.6
$\Gamma_{211}$	$\gamma\eta(1405/1475) \rightarrow \gamma\gamma\rho^0$		$(7.8 \pm 2.0) \times 10^{-5}$	S=1.8
$\Gamma_{212}$	$\gamma\eta(1405/1475) \rightarrow \gamma\eta\pi^+\pi^-$		$(3.0 \pm 0.5) \times 10^{-4}$	
$\Gamma_{213}$	$\gamma\eta(1405/1475) \rightarrow \gamma\gamma\phi$		$< 8.2 \times 10^{-5}$	CL=95%
$\Gamma_{214}$	$\gamma\eta(1405) \rightarrow \gamma\gamma\gamma$		$< 2.63 \times 10^{-6}$	CL=90%
$\Gamma_{215}$	$\gamma\eta(1475) \rightarrow \gamma\gamma\gamma$		$< 1.86 \times 10^{-6}$	CL=90%
$\Gamma_{216}$	$\gamma\rho\rho$		$(4.5 \pm 0.8) \times 10^{-3}$	
$\Gamma_{217}$	$\gamma\rho\omega$		$< 5.4 \times 10^{-4}$	CL=90%
$\Gamma_{218}$	$\gamma\rho\phi$		$< 8.8 \times 10^{-5}$	CL=90%
$\Gamma_{219}$	$\gamma\eta'(958)$		$(5.21 \pm 0.17) \times 10^{-3}$	S=1.4
$\Gamma_{220}$	$\gamma 2\pi^+ 2\pi^-$		$(2.8 \pm 0.5) \times 10^{-3}$	S=1.9
$\Gamma_{221}$	$\gamma f_2(1270) f_2(1270)$		$(9.5 \pm 1.7) \times 10^{-4}$	
$\Gamma_{222}$	$\gamma f_2(1270) f_2(1270)$ (non resonant)		$(8.2 \pm 1.9) \times 10^{-4}$	
$\Gamma_{223}$	$\gamma K^+ K^- \pi^+ \pi^-$		$(2.1 \pm 0.6) \times 10^{-3}$	
$\Gamma_{224}$	$\gamma f_4(2050)$		$(2.7 \pm 0.7) \times 10^{-3}$	

$\Gamma_{225}$	$\gamma\omega\omega$	$(1.61 \pm 0.33) \times 10^{-3}$	
$\Gamma_{226}$	$\gamma\eta(1405/1475) \rightarrow \gamma\rho^0\rho^0$	$(1.7 \pm 0.4) \times 10^{-3}$	S=1.3
$\Gamma_{227}$	$\gamma f_2(1270)$	$(1.64 \pm 0.12) \times 10^{-3}$	S=1.3
$\Gamma_{228}$	$\gamma f_2(1270) \rightarrow \gamma K_S^0 K_S^0$	$(2.58 \pm_{-0.22}^{0.60}) \times 10^{-5}$	
$\Gamma_{229}$	$\gamma f_0(1370) \rightarrow \gamma K\bar{K}$	$(4.2 \pm 1.5) \times 10^{-4}$	
$\Gamma_{230}$	$\gamma f_0(1370) \rightarrow \gamma K_S^0 K_S^0$	$(1.1 \pm 0.4) \times 10^{-5}$	
$\Gamma_{231}$	$\gamma f_0(1500) \rightarrow \gamma K_S^0 K_S^0$	$(1.59 \pm_{-0.60}^{0.24}) \times 10^{-5}$	
$\Gamma_{232}$	$\gamma f_0(1710) \rightarrow \gamma K\bar{K}$	$(9.5 \pm_{-0.5}^{1.0}) \times 10^{-4}$	S=1.5
$\Gamma_{233}$	$\gamma f_0(1710) \rightarrow \gamma\pi\pi$	$(3.8 \pm 0.5) \times 10^{-4}$	
$\Gamma_{234}$	$\gamma f_0(1710) \rightarrow \gamma\omega\omega$	$(3.1 \pm 1.0) \times 10^{-4}$	
$\Gamma_{235}$	$\gamma f_0(1710) \rightarrow \gamma\eta\eta$	$(2.4 \pm_{-0.7}^{1.2}) \times 10^{-4}$	
$\Gamma_{236}$	$\gamma\eta$	$(1.108 \pm 0.027) \times 10^{-3}$	
$\Gamma_{237}$	$\gamma f_1(1420) \rightarrow \gamma K\bar{K}\pi$	$(7.9 \pm 1.3) \times 10^{-4}$	
$\Gamma_{238}$	$\gamma f_1(1285)$	$(6.1 \pm 0.8) \times 10^{-4}$	
$\Gamma_{239}$	$\gamma f_1(1510) \rightarrow \gamma\eta\pi^+\pi^-$	$(4.5 \pm 1.2) \times 10^{-4}$	
$\Gamma_{240}$	$\gamma f_2'(1525)$	$(5.7 \pm_{-0.5}^{0.8}) \times 10^{-4}$	S=1.5
$\Gamma_{241}$	$\gamma f_2'(1525) \rightarrow \gamma K_S^0 K_S^0$	$(8.0 \pm_{-0.5}^{0.7}) \times 10^{-5}$	
$\Gamma_{242}$	$\gamma f_2'(1525) \rightarrow \gamma\eta\eta$	$(3.4 \pm 1.4) \times 10^{-5}$	
$\Gamma_{243}$	$\gamma f_2(1640) \rightarrow \gamma\omega\omega$	$(2.8 \pm 1.8) \times 10^{-4}$	
$\Gamma_{244}$	$\gamma f_2(1910) \rightarrow \gamma\omega\omega$	$(2.0 \pm 1.4) \times 10^{-4}$	
$\Gamma_{245}$	$\gamma f_0(1750) \rightarrow \gamma K_S^0 K_S^0$	$(1.11 \pm_{-0.33}^{0.20}) \times 10^{-5}$	
$\Gamma_{246}$	$\gamma f_0(1800) \rightarrow \gamma\omega\phi$	$(2.5 \pm 0.6) \times 10^{-4}$	
$\Gamma_{247}$	$\gamma f_2(1810) \rightarrow \gamma\eta\eta$	$(5.4 \pm_{-2.4}^{3.5}) \times 10^{-5}$	
$\Gamma_{248}$	$\gamma f_2(1950) \rightarrow$ $\gamma K^*(892)\bar{K}^*(892)$	$(7.0 \pm 2.2) \times 10^{-4}$	
$\Gamma_{249}$	$\gamma K^*(892)\bar{K}^*(892)$	$(4.0 \pm 1.3) \times 10^{-3}$	
$\Gamma_{250}$	$\gamma\phi\phi$	$(4.0 \pm 1.2) \times 10^{-4}$	S=2.1
$\Gamma_{251}$	$\gamma p\bar{p}$	$(3.8 \pm 1.0) \times 10^{-4}$	
$\Gamma_{252}$	$\gamma\eta(2225)$	$(3.14 \pm_{-0.19}^{0.50}) \times 10^{-4}$	
$\Gamma_{253}$	$\gamma\eta(1760) \rightarrow \gamma\rho^0\rho^0$	$(1.3 \pm 0.9) \times 10^{-4}$	
$\Gamma_{254}$	$\gamma\eta(1760) \rightarrow \gamma\omega\omega$	$(1.98 \pm 0.33) \times 10^{-3}$	
$\Gamma_{255}$	$\gamma\eta(1760) \rightarrow \gamma\gamma\gamma$	$< 4.80 \times 10^{-6}$	CL=90%
$\Gamma_{256}$	$\gamma X(1835) \rightarrow \gamma\pi^+\pi^-\eta'$	$(2.77 \pm_{-0.40}^{0.34}) \times 10^{-4}$	S=1.1
$\Gamma_{257}$	$\gamma X(1835) \rightarrow \gamma p\bar{p}$	$(7.7 \pm_{-0.9}^{1.5}) \times 10^{-5}$	
$\Gamma_{258}$	$\gamma X(1835) \rightarrow \gamma K_S^0 K_S^0 \eta$	$(3.3 \pm_{-1.3}^{2.0}) \times 10^{-5}$	
$\Gamma_{259}$	$\gamma X(1835) \rightarrow \gamma\gamma\phi(1020)$		
$\Gamma_{260}$	$\gamma X(1835) \rightarrow \gamma\gamma\gamma$	$< 3.56 \times 10^{-6}$	CL=90%



$\Gamma_{261}$	$\gamma X(1840) \rightarrow \gamma 3(\pi^+ \pi^-)$	$( 2.4 \begin{smallmatrix} + 0.7 \\ - 0.8 \end{smallmatrix} ) \times 10^{-5}$	
$\Gamma_{262}$	$\gamma(K\bar{K}\pi) [J^{PC} = 0^{-+}]$	$( 7 \pm 4 ) \times 10^{-4}$	S=2.1
$\Gamma_{263}$	$\gamma\pi^0$	$( 3.56 \pm 0.17 ) \times 10^{-5}$	
$\Gamma_{264}$	$\gamma p\bar{p}\pi^+\pi^-$	$< 7.9 \times 10^{-4}$	CL=90%
$\Gamma_{265}$	$\gamma\Lambda\bar{\Lambda}$	$< 1.3 \times 10^{-4}$	CL=90%
$\Gamma_{266}$	$\gamma f_0(2100) \rightarrow \gamma\eta\eta$	$( 1.13 \begin{smallmatrix} + 0.60 \\ - 0.30 \end{smallmatrix} ) \times 10^{-4}$	
$\Gamma_{267}$	$\gamma f_0(2100) \rightarrow \gamma\pi\pi$	$( 6.2 \pm 1.0 ) \times 10^{-4}$	
$\Gamma_{268}$	$\gamma f_0(2200)$		
$\Gamma_{269}$	$\gamma f_0(2200) \rightarrow \gamma K\bar{K}$	$( 5.9 \pm 1.3 ) \times 10^{-4}$	
$\Gamma_{270}$	$\gamma f_0(2200) \rightarrow \gamma K_S^0 K_S^0$	$( 2.72 \begin{smallmatrix} + 0.19 \\ - 0.50 \end{smallmatrix} ) \times 10^{-4}$	
$\Gamma_{271}$	$\gamma f_J(2220)$		
$\Gamma_{272}$	$\gamma f_J(2220) \rightarrow \gamma\pi\pi$	$< 3.9 \times 10^{-5}$	CL=90%
$\Gamma_{273}$	$\gamma f_J(2220) \rightarrow \gamma K\bar{K}$	$< 4.1 \times 10^{-5}$	CL=90%
$\Gamma_{274}$	$\gamma f_J(2220) \rightarrow \gamma p\bar{p}$	$( 1.5 \pm 0.8 ) \times 10^{-5}$	
$\Gamma_{275}$	$\gamma f_0(2330) \rightarrow \gamma K_S^0 K_S^0$	$( 4.9 \pm 0.7 ) \times 10^{-5}$	
$\Gamma_{276}$	$\gamma f_2(2340) \rightarrow \gamma\eta\eta$	$( 5.6 \begin{smallmatrix} + 2.4 \\ - 2.2 \end{smallmatrix} ) \times 10^{-5}$	
$\Gamma_{277}$	$\gamma f_2(2340) \rightarrow \gamma K_S^0 K_S^0$	$( 5.5 \begin{smallmatrix} + 4.0 \\ - 1.5 \end{smallmatrix} ) \times 10^{-5}$	
$\Gamma_{278}$	$\gamma f_0(1500) \rightarrow \gamma\pi\pi$	$( 1.09 \pm 0.24 ) \times 10^{-4}$	
$\Gamma_{279}$	$\gamma f_0(1500) \rightarrow \gamma\eta\eta$	$( 1.7 \begin{smallmatrix} + 0.6 \\ - 1.4 \end{smallmatrix} ) \times 10^{-5}$	
$\Gamma_{280}$	$\gamma A \rightarrow \gamma \text{invisible}$	$[e] < 6.3 \times 10^{-6}$	CL=90%
$\Gamma_{281}$	$\gamma A^0 \rightarrow \gamma\mu^+\mu^-$	$[f] < 5 \times 10^{-6}$	CL=90%

### Dalitz decays

$\Gamma_{282}$	$\pi^0 e^+ e^-$	$( 7.6 \pm 1.4 ) \times 10^{-7}$	
$\Gamma_{283}$	$\eta e^+ e^-$	$( 1.43 \pm 0.07 ) \times 10^{-5}$	
$\Gamma_{284}$	$\eta'(958) e^+ e^-$	$( 6.59 \pm 0.18 ) \times 10^{-5}$	
$\Gamma_{285}$	$\eta U \rightarrow \eta e^+ e^-$	$< 9.11 \times 10^{-7}$	CL=90%
$\Gamma_{286}$	$\eta'(958) U \rightarrow \eta'(958) e^+ e^-$	$< 2.0 \times 10^{-7}$	CL=90%

### Weak decays

$\Gamma_{287}$	$D^- e^+ \nu_e + \text{c.c.}$	$< 1.2 \times 10^{-5}$	CL=90%
$\Gamma_{288}$	$\bar{D}^0 e^+ e^- + \text{c.c.}$	$< 8.5 \times 10^{-8}$	CL=90%
$\Gamma_{289}$	$D_s^- e^+ \nu_e + \text{c.c.}$	$< 1.3 \times 10^{-6}$	CL=90%
$\Gamma_{290}$	$D_s^{*-} e^+ \nu_e + \text{c.c.}$	$< 1.8 \times 10^{-6}$	CL=90%
$\Gamma_{291}$	$D^- \pi^+ + \text{c.c.}$	$< 7.5 \times 10^{-5}$	CL=90%
$\Gamma_{292}$	$\bar{D}^0 \bar{K}^0 + \text{c.c.}$	$< 1.7 \times 10^{-4}$	CL=90%
$\Gamma_{293}$	$\bar{D}^0 \bar{K}^{*0} + \text{c.c.}$	$< 2.5 \times 10^{-6}$	CL=90%
$\Gamma_{294}$	$D_s^- \pi^+ + \text{c.c.}$	$< 1.3 \times 10^{-4}$	CL=90%
$\Gamma_{295}$	$D_s^- \rho^+ + \text{c.c.}$	$< 1.3 \times 10^{-5}$	CL=90%

### Charge conjugation (*C*), Parity (*P*), Lepton Family number (*LF*) violating modes

$\Gamma_{296}$	$\gamma\gamma$	<i>C</i>	< 2.7	$\times 10^{-7}$	CL=90%
$\Gamma_{297}$	$\gamma\phi$	<i>C</i>	< 1.4	$\times 10^{-6}$	CL=90%
$\Gamma_{298}$	$e^\pm\mu^\mp$	<i>LF</i>	< 1.6	$\times 10^{-7}$	CL=90%
$\Gamma_{299}$	$e^\pm\tau^\mp$	<i>LF</i>	< 8.3	$\times 10^{-6}$	CL=90%
$\Gamma_{300}$	$\mu^\pm\tau^\mp$	<i>LF</i>	< 2.0	$\times 10^{-6}$	CL=90%

### Other decays

$\Gamma_{301}$	invisible		< 7	$\times 10^{-4}$	CL=90%
----------------	-----------	--	-----	------------------	--------

[a] For  $E_\gamma > 100$  MeV.

[b] The value is for the sum of the charge states or particle/antiparticle states indicated.

[c] Includes  $p\bar{p}\pi^+\pi^-\gamma$  and excludes  $p\bar{p}\eta$ ,  $p\bar{p}\omega$ ,  $p\bar{p}\eta'$ .

[d] See the "Note on the  $\eta(1405)$ " in the  $\eta(1405)$  Particle Listings.

[e] For a narrow state *A* with mass less than 960 MeV.

[f] For a narrow scalar or pseudoscalar  $A^0$  with mass 0.21–3.0 GeV.

## $J/\psi(1S)$ PARTIAL WIDTHS

### $\Gamma(\text{hadrons})$

$\Gamma_1$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$74.1 \pm 8.1$	BAI	95B	BES $e^+e^-$
$59 \pm 24$	BALDINI-...	75	FRAG $e^+e^-$
$59 \pm 14$	BOYARSKI	75	MRK1 $e^+e^-$
$50 \pm 25$	ESPOSITO	75B	FRAM $e^+e^-$

### $\Gamma(e^+e^-)$

$\Gamma_5$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>5.53 ± 0.10 OUR AVERAGE</b>				
$5.550 \pm 0.056 \pm 0.089$		1 ANASHIN	18A	KEDR $e^+e^-$
$5.36^{+0.29}_{-0.28}$		2 HSUEH	92	RVUE See $\Upsilon$ mini-review
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$5.58 \pm 0.05 \pm 0.08$		3 ABLIKIM	16Q	BES3 $3.773 e^+e^- \rightarrow \mu^+\mu^-\gamma$
$5.71 \pm 0.16$	13k	4 ADAMS	06A	CLEO $e^+e^- \rightarrow \mu^+\mu^-\gamma$
$5.57 \pm 0.19$	7.8k	4 AUBERT	04	BABR $e^+e^- \rightarrow \mu^+\mu^-\gamma$
$5.14 \pm 0.39$		BAI	95B	BES $e^+e^-$
$4.72 \pm 0.35$		ALEXANDER	89	RVUE See $\Upsilon$ mini-review
$4.4 \pm 0.6$		2 BRANDELIK	79C	DASP $e^+e^-$
$4.6 \pm 0.8$		5 BALDINI-...	75	FRAG $e^+e^-$
$4.8 \pm 0.6$		BOYARSKI	75	MRK1 $e^+e^-$
$4.6 \pm 1.0$		ESPOSITO	75B	FRAM $e^+e^-$

<sup>1</sup> From the cross sections of  $e^+e^- \rightarrow e^+e^-$  and  $e^+e^- \rightarrow$  hadrons near the  $J/\psi(1S)$  peak.

<sup>2</sup> From a simultaneous fit to  $e^+e^-$ ,  $\mu^+\mu^-$ , and hadronic channels assuming  $\Gamma(e^+e^-) = \Gamma(\mu^+\mu^-)$ .

<sup>3</sup> Using  $B(J/\psi \rightarrow \mu^+\mu^-) = (5.973 \pm 0.007 \pm 0.037)\%$  from ABLIKIM 13R.

<sup>4</sup> Calculated by us from the reported values of  $\Gamma(e^+e^-) \times B(\mu^+\mu^-)$  using  $B(\mu^+\mu^-) = (5.93 \pm 0.06)\%$ .

<sup>5</sup> Assuming equal partial widths for  $e^+e^-$  and  $\mu^+\mu^-$ .

### $\Gamma(\mu^+\mu^-)$ $\Gamma_7$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$5.13 \pm 0.52$	BAI	95B	BES $e^+e^-$
$4.8 \pm 0.6$	BOYARSKI	75	MRK1 $e^+e^-$
$5 \pm 1$	ESPOSITO	75B	FRAM $e^+e^-$

### $\Gamma(\gamma\gamma)$ $\Gamma_{296}$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;5.4</b>	90	BRANDELIK	79C	DASP $e^+e^-$

### $J/\psi(1S) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

This combination of a partial width with the partial width into  $e^+e^-$  and with the total width is obtained from the integrated cross section into channel  $i$  in the  $e^+e^-$  annihilation.

### $\Gamma(\text{hadrons}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_1\Gamma_5/\Gamma$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
<b><math>4.884 \pm 0.048 \pm 0.078</math></b>	<sup>1</sup> ANASHIN	18A	KEDR $e^+e^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$4 \pm 0.8$	<sup>2</sup> BALDINI-...	75	FRAG $e^+e^-$
$3.9 \pm 0.8$	<sup>2</sup> ESPOSITO	75B	FRAM $e^+e^-$

<sup>1</sup> From the cross sections of  $e^+e^- \rightarrow e^+e^-$  and  $e^+e^- \rightarrow$  hadrons near the  $J/\psi(1S)$  peak.

<sup>2</sup> Data redundant with branching ratios or partial widths above.

### $\Gamma(e^+e^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_5\Gamma_5/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
<b><math>333.1 \pm 6.6 \pm 4.0</math></b>	<sup>1</sup> ANASHIN	18A	KEDR $e^+e^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$332.3 \pm 6.4 \pm 4.8$	ANASHIN	10	KEDR $3.097 e^+e^- \rightarrow e^+e^-$
$350 \pm 20$	BRANDELIK	79C	DASP $e^+e^-$
$320 \pm 70$	<sup>2</sup> BALDINI-...	75	FRAG $e^+e^-$
$340 \pm 90$	<sup>2</sup> ESPOSITO	75B	FRAM $e^+e^-$
$360 \pm 100$	<sup>2</sup> FORD	75	SPEC $e^+e^-$

<sup>1</sup> From the cross sections of  $e^+e^- \rightarrow e^+e^-$  and  $e^+e^- \rightarrow$  hadrons near the  $J/\psi(1S)$  peak.

<sup>2</sup> Data redundant with branching ratios or partial widths above.

$\Gamma(\mu^+\mu^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_7\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

**333 ± 4 OUR AVERAGE**

333.4 ± 2.5 ± 4.4		ABLIKIM	16Q BES3	3.773 $e^+e^- \rightarrow \mu^+\mu^-\gamma$
331.8 ± 5.2 ± 6.3		ANASHIN	10 KEDR	3.097 $e^+e^- \rightarrow \mu^+\mu^-$
338.4 ± 5.8 ± 7.1	13k	ADAMS	06A CLEO	$e^+e^- \rightarrow \mu^+\mu^-\gamma$
330.1 ± 7.7 ± 7.3	7.8k	AUBERT	04 BABR	$e^+e^- \rightarrow \mu^+\mu^-\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

510 ± 90		DASP	75 DASP	$e^+e^-$
380 ± 50		<sup>1</sup> ESPOSITO	75B FRAM	$e^+e^-$

<sup>1</sup>Data redundant with branching ratios or partial widths above.

$\Gamma(\rho(770)^\mp K^\pm K_S^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{10}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

<b>10.4 ± 1.0 ± 1.9</b>	130	LEES	17D BABR	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$
-------------------------	-----	------	----------	---

$\Gamma(\omega\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{23}\Gamma_5/\Gamma$

VALUE (10 <sup>-2</sup> keV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------------------------	------	-------------	------	---------

<b>2.2 ± 0.3 ± 0.2</b>	170	AUBERT	06D BABR	10.6 $e^+e^- \rightarrow \omega\pi^+\pi^-\pi^0\gamma$
------------------------	-----	--------	----------	---

$\Gamma(\omega\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{24}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

<b>53.6 ± 5.0 ± 0.4</b>	788	<sup>1</sup> AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$
-------------------------	-----	---------------------	-----------	--

<sup>1</sup>AUBERT 07AU reports [ $\Gamma(J/\psi(1S) \rightarrow \omega\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$ ] × [B( $\omega(782) \rightarrow \pi^+\pi^-\pi^0$ )] = 47.8 ± 3.1 ± 3.2 eV which we divide by our best value B( $\omega(782) \rightarrow \pi^+\pi^-\pi^0$ ) = (89.3 ± 0.6) × 10<sup>-2</sup>. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\omega\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{59}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

<b>27.8 ± 3.5 ± 0.2</b>	398	<sup>1</sup> LEES	18E BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$
-------------------------	-----	-------------------	----------	---

<sup>1</sup>LEES 18E reports [ $\Gamma(J/\psi(1S) \rightarrow \omega\pi^0\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$ ] × [B( $\omega(782) \rightarrow \pi^+\pi^-\pi^0$ )] = 24.8 ± 1.8 ± 2.5 eV which we divide by our best value B( $\omega(782) \rightarrow \pi^+\pi^-\pi^0$ ) = (89.3 ± 0.6) × 10<sup>-2</sup>. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^*(892)^0 \bar{K}^*(892)^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{26}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

<b>1.28 ± 0.34 ± 0.07</b>	47 ± 12	<sup>1</sup> LEES	12F BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$
---------------------------	---------	-------------------	----------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.28 ± 0.40 ± 0.11	25 ± 8	<sup>1,2</sup> AUBERT	07AK BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$
--------------------	--------	-----------------------	-----------	--

<sup>1</sup>Dividing by (2/3)<sup>2</sup> to take twice into account that B( $K^{*0} \rightarrow K^+\pi^-$ ) = 2/3 B( $K^{*0} \rightarrow K\pi$ ).

<sup>2</sup>Superseded by LEES 12F.

$$\Gamma(K^*(892)^\pm K^*(892)^\mp) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{27}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.80±0.48±0.32</b>	1 ± 5	<sup>1</sup> LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$

<sup>1</sup> Dividing by  $(1/4)^2$  to take twice into account  $B(K^*(892) \rightarrow K_S^0 \pi) = 1/4$ .

$$\Gamma(K_S^0 \pi^- K^*(892)^+ + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{29}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>11.0±2.8 OUR AVERAGE</b>				
9.2±1.2±3.2	64	<sup>1</sup> LEES	17D BABR	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$
14.8±4.8±1.2	53	<sup>2</sup> LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$

<sup>1</sup> Dividing by 1/2 to take into account  $B(K^*(892)^\pm \rightarrow K^\pm \pi^\mp) = 1/2$ .

<sup>2</sup> Dividing by 1/4 to take into account  $B(K^*(892) \rightarrow K_S^0 \pi) = 1/4$ .

$$\Gamma(K_S^0 \pi^- K^*(892)^+ + \text{c.c.} \rightarrow K_S^0 K_S^0 \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{30}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.7±1.2±0.3</b>	53	LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$

$$\Gamma(K^*(892)^0 \bar{K}_2^*(1430)^0 + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{43}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>25.8±1.4±0.6</b>	710	<sup>1,2,3</sup> LEES	12F BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

33 ± 4 ± 1	317	<sup>2,4</sup> AUBERT	07AK BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$
------------	-----	-----------------------	-----------	---

<sup>1</sup> LEES 12F reports  $[\Gamma(J/\psi(1S) \rightarrow K^*(892)^0 \bar{K}_2^*(1430)^0 + \text{c.c.}) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(K_2^*(1430) \rightarrow K \pi)] = 12.89 \pm 0.54 \pm 0.41$  eV which we divide by our best value  $B(K_2^*(1430) \rightarrow K \pi) = (49.9 \pm 1.2) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Dividing by 2/3 to take into account that  $B(K^{*0} \rightarrow K^+ \pi^-) = 2/3 B(K^{*0} \rightarrow K \pi)$ .

<sup>3</sup> The  $K_2^*(1430)$  cannot be distinguished from the  $K_0^*(1430)$ .

<sup>4</sup> Superseded by LEES 12F. AUBERT 07AK reports  $[\Gamma(J/\psi(1S) \rightarrow K^*(892)^0 \bar{K}_2^*(1430)^0 + \text{c.c.}) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(K_2^*(1430) \rightarrow K \pi)] = 16.4 \pm 1.1 \pm 1.4$  eV which we divide by our best value  $B(K_2^*(1430) \rightarrow K \pi) = (49.9 \pm 1.2) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(K^*(892)^+ K_2^*(1430)^- + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{44}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>18.6±16.1±0.4</b>	8 ± 8	<sup>1,2</sup> LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$

<sup>1</sup> Dividing by  $(1/4)^2$  to take into account  $B(K^*(892) \rightarrow K_S^0 \pi) = 1/4$  and  $B(K^*(1430) \rightarrow K_S^0 \pi) = 1/4 B(K^*(1430) \rightarrow K \pi)$ .

<sup>2</sup> LEES 14H reports  $[\Gamma(J/\psi(1S) \rightarrow K^*(892)^+ K_2^*(1430)^- + \text{c.c.}) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(K_2^*(1430) \rightarrow K \pi)] = 9.28 \pm 8.0 \pm 0.32$  eV which we divide by our best value  $B(K_2^*(1430) \rightarrow K \pi) = (49.9 \pm 1.2) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(K^*(892)^+ K_2^*(1430)^- + \text{c.c.} \rightarrow K^*(892)^+ K_S^0 \pi^- + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{45} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.32 ± 2.00 ± 0.08</b>	8 ± 8	<sup>1</sup> LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$

<sup>1</sup> Dividing by 1/4 to take into account  $B(K^*(892) \rightarrow K_S^0 \pi) = 1/4$ .

$$\Gamma(K^*(892)^0 \bar{K}_2(1770)^0 + \text{c.c.} \rightarrow K^*(892)^0 K^- \pi^+ + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{46} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.8 ± 0.4 ± 0.3</b>	110 ± 14	<sup>1</sup> AUBERT 07AK	BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

<sup>1</sup> Dividing by 2/3 to take into account that  $B(K^*0 \rightarrow K^+ \pi^-) = 2/3$ .

$$\Gamma(K^+ K^*(892)^- + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{50} \Gamma_5 / \Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
<b>29.0 ± 1.7 ± 1.3</b>	AUBERT 08S	BABR	10.6 $e^+ e^- \rightarrow K^+ K^*(892)^- \gamma$

$$\Gamma(K^+ K^*(892)^- + \text{c.c.} \rightarrow K^+ K^- \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{51} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>10.96 ± 0.85 ± 0.70</b>	155	AUBERT 08S	BABR	10.6 $e^+ e^- \rightarrow K^+ K^- \pi^0 \gamma$

$$\Gamma(K^+ K^*(892)^- + \text{c.c.} \rightarrow K^0 K^\pm \pi^\mp + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{52} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>16.76 ± 1.70 ± 1.00</b>	89	AUBERT 08S	BABR	10.6 $e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \gamma$

$$\Gamma(K^0 \bar{K}^*(892)^0 + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{53} \Gamma_5 / \Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
<b>26.6 ± 2.5 ± 1.5</b>	AUBERT 08S	BABR	10.6 $e^+ e^- \rightarrow K^0 \bar{K}^*(892)^0 \gamma$

$$\Gamma(K^0 \bar{K}^*(892)^0 + \text{c.c.} \rightarrow K^0 K^\pm \pi^\mp + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{54} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>17.70 ± 1.70 ± 1.00</b>	94	AUBERT 08S	BABR	10.6 $e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \gamma$

$$\Gamma(K_S^0 K^*(892)^0 \rightarrow \gamma K_S^0 K_S^0) / \Gamma_{\text{total}} \quad \Gamma_{31} / \Gamma$$

VALUE (units 10 <sup>-6</sup> )	DOCUMENT ID	TECN	COMMENT
<b>6.28<sup>+0.16+0.59</sup><sub>-0.17-0.52</sub></b>	ABLIKIM 18AA	BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

$$\Gamma(\bar{K}^*(892)^0 K^+ \pi^- + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{56} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>42.6 ± 4.8 ± 7.2</b>	99	<sup>1</sup> LEES	17D BABR	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$

<sup>1</sup> Dividing by 1/6 to account for  $B(K^*(892)^0 \rightarrow K_S^0 \pi^0) = 1/6$ .

$$\Gamma(K^*(892)^\pm K^\mp \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{57} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>22.8 ± 2.8 ± 6.8</b>	80	<sup>1</sup> LEES	17D BABR	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$

<sup>1</sup> Dividing by 1/4 to account for  $B(K^*(892)^\pm \rightarrow K_S^0 \pi^\pm) = 1/4$ .

$\Gamma(K^*(892)^0 K_S^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$   $\Gamma_{58} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.60±0.75±2.25</b>	34	<sup>1</sup> LEES	17D BABR	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$

<sup>1</sup> Dividing by 2/3 to account for  $B(K^*(892)^0 \rightarrow K^+ \pi^-) = 2/3$ .

$\Gamma(\eta K^\pm K_S^0 \pi^\mp) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$   $\Gamma_{64} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>7.3±1.4±0.4</b>	44	LEES	17D BABR	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$

$\Gamma(\omega K \bar{K}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$   $\Gamma_{66} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.70±1.98±0.03</b>	24	<sup>1</sup> AUBERT	07AU BABR	$10.6 e^+ e^- \rightarrow \omega K^+ K^- \gamma$

<sup>1</sup> AUBERT 07AU reports  $[\Gamma(J/\psi(1S) \rightarrow \omega K \bar{K}) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0)] = 3.3 \pm 1.3 \pm 1.2$  eV which we divide by our best value  $B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0) = (89.3 \pm 0.6) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi 2(\pi^+ \pi^-)) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$   $\Gamma_{68} \Gamma_5 / \Gamma$

VALUE ( $10^{-2}$ keV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.95±0.19±0.01</b>	35	<sup>1</sup> AUBERT	06D BABR	$10.6 e^+ e^- \rightarrow \phi 2(\pi^+ \pi^-) \gamma$

<sup>1</sup> AUBERT 06D reports  $[\Gamma(J/\psi(1S) \rightarrow \phi 2(\pi^+ \pi^-)) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = (0.47 \pm 0.09 \pm 0.03) \times 10^{-2}$  keV which we divide by our best value  $B(\phi(1020) \rightarrow K^+ K^-) = (49.2 \pm 0.5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi K_S^0 K_S^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$   $\Gamma_{72} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.25±0.84±0.03</b>	29	<sup>1</sup> LEES	14H BABR	$e^+ e^- \rightarrow K_S^0 K_S^0 K^+ K^- \gamma$

<sup>1</sup> LEES 14H reports  $[\Gamma(J/\psi(1S) \rightarrow \phi K_S^0 K_S^0) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 1.6 \pm 0.4 \pm 0.1$  eV which we divide by our best value  $B(\phi(1020) \rightarrow K^+ K^-) = (49.2 \pm 0.5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi K^+ K^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$   $\Gamma_{74} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.59±0.62±0.05</b>	163	<sup>1</sup> LEES	12F BABR	$10.6 e^+ e^- \rightarrow K^+ K^- K^+ K^- \gamma$

<sup>1</sup> LEES 12F reports  $[\Gamma(J/\psi(1S) \rightarrow \phi K^+ K^-) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 2.26 \pm 0.26 \pm 0.16$  eV which we divide by our best value  $B(\phi(1020) \rightarrow K^+ K^-) = (49.2 \pm 0.5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi f_2(1270)) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$   $\Gamma_{75}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

<b>1.79±0.32<sup>+0.02</sup><sub>-0.06</sub></b>	61	1,2,3 LEES	12F BABR	10.6 e <sup>+</sup> e <sup>-</sup> → π <sup>+</sup> π <sup>-</sup> K <sup>+</sup> K <sup>-</sup> γ
--	----	------------	----------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.08±0.73 <sup>+0.04</sup> <sub>-0.14</sub>	44	2,4 AUBERT	07AK BABR	10.6 e <sup>+</sup> e <sup>-</sup> → π <sup>+</sup> π <sup>-</sup> K <sup>+</sup> K <sup>-</sup> γ
---	----	------------	-----------	--

<sup>1</sup> LEES 12F reports  $[\Gamma(J/\psi(1S) \rightarrow \phi f_2(1270)) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = 1.51 \pm 0.25 \pm 0.10$  eV which we divide by our best value  $B(f_2(1270) \rightarrow \pi\pi) = (84.2^{+2.9}_{-0.9}) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Using  $B(\phi \rightarrow K^+ K^-) = (48.9 \pm 0.5)\%$ .

<sup>3</sup> Using π<sup>+</sup>π<sup>-</sup> invariant mass between 1.1 and 1.5 GeV. May include other sources such as f<sub>0</sub>(1370).

<sup>4</sup> Superseded by LEES 12F. AUBERT 07AK reports  $[\Gamma(J/\psi(1S) \rightarrow \phi f_2(1270)) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = 3.44 \pm 0.55 \pm 0.28$  eV which we divide by our best value  $B(f_2(1270) \rightarrow \pi\pi) = (84.2^{+2.9}_{-0.9}) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^+ K^- f'_2(1525)) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$   $\Gamma_{79}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

<b>5.8±1.9±0.1</b>	16	1,2 LEES	14H BABR	e <sup>+</sup> e <sup>-</sup> → K <sub>S</sub> <sup>0</sup> K <sub>S</sub> <sup>0</sup> K <sup>+</sup> K <sup>-</sup> γ
--------------------	----	----------	----------	---

<sup>1</sup> Dividing by 1/4 to take into account  $B(f'_2(1525) \rightarrow K_S^0 K_S^0) = 1/4 B(f'_2(1525) \rightarrow K\bar{K})$ .

<sup>2</sup> LEES 14H reports  $[\Gamma(J/\psi(1S) \rightarrow K^+ K^- f'_2(1525)) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(f'_2(1525) \rightarrow K\bar{K})] = 5.12 \pm 1.68 \pm 0.20$  eV which we divide by our best value  $B(f'_2(1525) \rightarrow K\bar{K}) = (88.7 \pm 2.2) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi f'_2(1525)) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$   $\Gamma_{80}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

<b>8.1±3.2±0.2</b>	11	1,2 LEES	14H BABR	e <sup>+</sup> e <sup>-</sup> → K <sub>S</sub> <sup>0</sup> K <sub>S</sub> <sup>0</sup> K <sup>+</sup> K <sup>-</sup> γ
--------------------	----	----------	----------	---

<sup>1</sup> Dividing by 1/4 to take into account  $B(f'_2(1525) \rightarrow K_S^0 K_S^0) = 1/4 B(f'_2(1525) \rightarrow K\bar{K})$  and using  $B(\phi \rightarrow K^+ K^-) = (48.9 \pm 0.5)\%$ .

<sup>2</sup> LEES 14H reports  $[\Gamma(J/\psi(1S) \rightarrow \phi f'_2(1525)) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(f'_2(1525) \rightarrow K\bar{K})] = 7.2 \pm 2.8 \pm 0.3$  eV which we divide by our best value  $B(f'_2(1525) \rightarrow K\bar{K}) = (88.7 \pm 2.2) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi\pi^+\pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$   $\Gamma_{81}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

**4.47±0.35 OUR AVERAGE**

4.45±0.49±0.05	181	<sup>1</sup> LEES	12F BABR	10.6 e <sup>+</sup> e <sup>-</sup> → K <sup>+</sup> K <sup>-</sup> π <sup>+</sup> π <sup>-</sup> γ
----------------	-----	-------------------	----------	--

4.50±0.48±0.05	254 ± 23	<sup>2</sup> SHEN	09 BELL	10.6 e <sup>+</sup> e <sup>-</sup> → K <sup>+</sup> K <sup>-</sup> π <sup>+</sup> π <sup>-</sup> γ
----------------	----------	-------------------	---------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.3 ± 0.7 ± 0.1	103	<sup>3</sup> AUBERT, BE	06D BABR	10.6 e <sup>+</sup> e <sup>-</sup> → K <sup>+</sup> K <sup>-</sup> π <sup>+</sup> π <sup>-</sup> γ
-----------------	-----	-------------------------	----------	--



<sup>1</sup> LEES 12F reports  $[\Gamma(J/\psi(1S) \rightarrow \phi\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 2.19 \pm 0.23 \pm 0.07$  eV which we divide by our best value  $B(\phi(1020) \rightarrow K^+K^-) = (49.2 \pm 0.5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> SHEN 09 reports  $4.50 \pm 0.41 \pm 0.26$  eV from a measurement of  $[\Gamma(J/\psi(1S) \rightarrow \phi\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)]$  assuming  $B(\phi(1020) \rightarrow K^+K^-) = (49.2 \pm 0.6) \times 10^{-2}$ , which we rescale to our best value  $B(\phi(1020) \rightarrow K^+K^-) = (49.2 \pm 0.5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>3</sup> Superseded by LEES 12F. AUBERT, BE 06D reports  $[\Gamma(J/\psi(1S) \rightarrow \phi\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 2.61 \pm 0.30 \pm 0.18$  eV which we divide by our best value  $B(\phi(1020) \rightarrow K^+K^-) = (49.2 \pm 0.5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

### $\Gamma(\phi\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{82}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.76±0.57±0.03</b>	45	<sup>1</sup> LEES 12F	BABR	10.6 $e^+e^- \rightarrow K^+K^-\pi^0\pi^0\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.13±0.88±0.03	23	<sup>2</sup> AUBERT, BE 06D	BABR	10.6 $e^+e^- \rightarrow K^+K^-\pi^0\pi^0\gamma$
----------------	----	-----------------------------	------	--

<sup>1</sup> LEES 12F reports  $[\Gamma(J/\psi(1S) \rightarrow \phi\pi^0\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 1.36 \pm 0.27 \pm 0.07$  eV which we divide by our best value  $B(\phi(1020) \rightarrow K^+K^-) = (49.2 \pm 0.5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Superseded by LEES 12F. AUBERT, BE 06D reports  $[\Gamma(J/\psi(1S) \rightarrow \phi\pi^0\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 1.54 \pm 0.40 \pm 0.16$  eV which we divide by our best value  $B(\phi(1020) \rightarrow K^+K^-) = (49.2 \pm 0.5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

### $\Gamma(\phi\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{85}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>6.1±2.7±0.4</b>	6	<sup>1</sup> AUBERT 07AU	BABR	10.6 $e^+e^- \rightarrow \phi\eta\gamma$

<sup>1</sup> AUBERT 07AU quotes  $\Gamma_{ee}^{J/\psi} \cdot B(J/\psi \rightarrow \phi\eta) \cdot B(\phi \rightarrow K^+K^-) \cdot B(\eta \rightarrow 3\pi) = 0.84 \pm 0.37 \pm 0.05$  eV.

### $\Gamma(\phi f_0(980) \rightarrow \phi\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{93}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.44±0.19 OUR AVERAGE</b>				

1.40±0.25±0.02	57 ± 9	<sup>1</sup> LEES 12F	BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$
----------------	--------	-----------------------	------	--

1.48±0.27±0.09	60 ± 11	<sup>2</sup> SHEN 09	BELL	10.6 $e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
----------------	---------	----------------------	------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.02±0.24±0.01	20 ± 5	<sup>3</sup> AUBERT 07AK	BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$
----------------	--------	--------------------------	------	--

<sup>1</sup> LEES 12F reports  $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi \pi^+ \pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 0.69 \pm 0.11 \pm 0.05$  eV which we divide by our best value  $B(\phi(1020) \rightarrow K^+ K^-) = (49.2 \pm 0.5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Multiplied by 2/3 to take into account the  $\phi \pi^+ \pi^-$  mode only. Using  $B(\phi \rightarrow K^+ K^-) = (49.2 \pm 0.6)\%$ .

<sup>3</sup> Superseded by LEES 12F. AUBERT 07AK reports  $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi \pi^+ \pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 0.50 \pm 0.11 \pm 0.04$  eV which we divide by our best value  $B(\phi(1020) \rightarrow K^+ K^-) = (49.2 \pm 0.5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

### $\Gamma(\phi f_0(980) \rightarrow \phi \pi^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{94} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.98 ± 0.26 ± 0.01</b>	16 ± 4	<sup>1</sup> LEES 12F	BABR	10.6 $e^+ e^- \rightarrow \pi^0 \pi^0 K^+ K^- \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.95 ± 0.40 ± 0.01 7.0 ± 2.8 <sup>2</sup> AUBERT 07AK BABR 10.6  $e^+ e^- \rightarrow \pi^0 \pi^0 K^+ K^- \gamma$

<sup>1</sup> LEES 12F reports  $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi \pi^0 \pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 0.48 \pm 0.12 \pm 0.05$  eV which we divide by our best value  $B(\phi(1020) \rightarrow K^+ K^-) = (49.2 \pm 0.5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Superseded by LEES 12F. AUBERT 07AK reports  $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi \pi^0 \pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 0.47 \pm 0.19 \pm 0.05$  eV which we divide by our best value  $B(\phi(1020) \rightarrow K^+ K^-) = (49.2 \pm 0.5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

### $\Gamma(\eta \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{104} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.3 ± 0.4 OUR AVERAGE</b>				

2.34 ± 0.43 ± 0.16 49 LEES 18 BABR  $e^+ e^- \rightarrow \eta \pi^+ \pi^- \gamma$

2.23 ± 0.97 ± 0.03 9 <sup>1</sup> AUBERT 07AU BABR 10.6  $e^+ e^- \rightarrow \eta \pi^+ \pi^- \gamma$

<sup>1</sup> AUBERT 07AU reports  $[\Gamma(J/\psi(1S) \rightarrow \eta \pi^+ \pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\eta \rightarrow \pi^+ \pi^- \pi^0)] = 0.51 \pm 0.22 \pm 0.03$  eV which we divide by our best value  $B(\eta \rightarrow \pi^+ \pi^- \pi^0) = (22.92 \pm 0.28) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

### $\Gamma(K_S^0 \pi^- K_2^*(1430)^+ + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{113} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>20.1 ± 9.8 ± 0.5</b>	35	<sup>1,2</sup> LEES 14H	BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$

<sup>1</sup> Dividing by 1/4 to take into account  $B(K^*(1430) \rightarrow K_S^0 \pi) = 1/4 B(K^*(1430) \rightarrow K \pi)$ .

<sup>2</sup> LEES 14H reports  $[\Gamma(J/\psi(1S) \rightarrow K_S^0 \pi^- K_2^*(1430)^+ + \text{c.c.}) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(K_2^*(1430) \rightarrow K \pi)] = 10.0 \pm 4.8 \pm 0.8$  eV which we divide by our best value  $B(K_2^*(1430) \rightarrow K \pi) = (49.9 \pm 1.2) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(2(\pi^+\pi^-\pi^0)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{133}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>303±5±18</b>	4990	AUBERT	07AU BABR	10.6 e <sup>+</sup> e <sup>-</sup> → 2(π <sup>+</sup> π <sup>-</sup> )π <sup>0</sup> γ

$$\Gamma(\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{135}\Gamma_5/\Gamma$$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
<b>0.122±0.005±0.008</b>	AUBERT,B	04N BABR	10.6 e <sup>+</sup> e <sup>-</sup> → π <sup>+</sup> π <sup>-</sup> π <sup>0</sup> γ

$$\Gamma(\pi^+\pi^-\pi^0\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{136}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>150.0±4.0±15.0</b>	2.3k	LEES	18E BABR	10.6 e <sup>+</sup> e <sup>-</sup> → π <sup>+</sup> π <sup>-</sup> 3π <sup>0</sup> γ

$$\Gamma(\rho^\pm\pi^\mp\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{137}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>78.0±9.0±8.0</b>	1.2k	LEES	18E BABR	10.6 e <sup>+</sup> e <sup>-</sup> → π <sup>+</sup> π <sup>-</sup> 3π <sup>0</sup> γ

$$\Gamma(\rho^+\rho^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{138}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>33.0±5.0±3.3</b>	529	LEES	18E BABR	10.6 e <sup>+</sup> e <sup>-</sup> → π <sup>+</sup> π <sup>-</sup> 3π <sup>0</sup> γ

$$\Gamma(\pi^+\pi^-\pi^0K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{139}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>107.0±4.3±6.4</b>	768	AUBERT	07AU BABR	10.6 e <sup>+</sup> e <sup>-</sup> → K <sup>+</sup> K <sup>-</sup> π <sup>+</sup> π <sup>-</sup> π <sup>0</sup> γ

$$\Gamma(\pi^+\pi^-K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{141}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>37.94±0.81±1.10</b>	3.1k	LEES	12F BABR	10.6 e <sup>+</sup> e <sup>-</sup> → π <sup>+</sup> π <sup>-</sup> K <sup>+</sup> K <sup>-</sup> γ

• • • We do not use the following data for averages, fits, limits, etc. • • •

36.3 ±1.3 ±2.1	1.5k	<sup>1</sup> AUBERT	07AK BABR	10.6 e <sup>+</sup> e <sup>-</sup> → π <sup>+</sup> π <sup>-</sup> K <sup>+</sup> K <sup>-</sup> γ
----------------	------	---------------------	-----------	--

33.6 ±2.7 ±2.7	233	<sup>2</sup> AUBERT	05D BABR	10.6 e <sup>+</sup> e <sup>-</sup> → K <sup>+</sup> K <sup>-</sup> π <sup>+</sup> π <sup>-</sup> γ
----------------	-----	---------------------	----------	--

<sup>1</sup> Superseded by LEES 12F.

<sup>2</sup> Superseded by AUBERT 07AK.

$$\Gamma(\pi^+\pi^-K_S^0K_L^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{142}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>20.8±2.3±2.1</b>	248	LEES	14H BABR	e <sup>+</sup> e <sup>-</sup> → π <sup>+</sup> π <sup>-</sup> K <sub>S</sub> <sup>0</sup> K <sub>L</sub> <sup>0</sup> γ

$$\Gamma(\pi^+\pi^-K_S^0K_S^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{143}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>9.3±0.9±0.5</b>	133	LEES	14H BABR	e <sup>+</sup> e <sup>-</sup> → π <sup>+</sup> π <sup>-</sup> K <sub>S</sub> <sup>0</sup> K <sub>S</sub> <sup>0</sup> γ

$$\Gamma(\pi^\pm\pi^0K^\mp K_S^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{144}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>31.7±1.9±1.8</b>	393	LEES	17D BABR	e <sup>+</sup> e <sup>-</sup> → K <sub>S</sub> <sup>0</sup> K <sup>±</sup> π <sup>∓</sup> π <sup>0</sup> γ

$$\Gamma(K^+K^-K_S^0K_S^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{145}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.3±0.4±0.1</b>	29	LEES	14H BABR	e <sup>+</sup> e <sup>-</sup> → K <sub>S</sub> <sup>0</sup> K <sub>S</sub> <sup>0</sup> K <sup>+</sup> K <sup>-</sup> γ

$\Gamma(\pi^+\pi^-K^+K^-\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{146}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

<b>25.9±3.9±0.1</b>	73	<sup>1</sup> AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow K^+K^-\pi^+\pi^-\eta\gamma$
---------------------	----	---------------------	-----------	--

<sup>1</sup> AUBERT 07AU reports  $[\Gamma(J/\psi(1S) \rightarrow \pi^+\pi^-K^+K^-\eta) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] = 10.2 \pm 1.3 \pm 0.8$  eV which we divide by our best value  $B(\eta \rightarrow 2\gamma) = (39.41 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\pi^0\pi^0K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{147}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

<b>11.75±0.81±0.90</b>	388	LEES	12F BABR	10.6 $e^+e^- \rightarrow \pi^0\pi^0K^+K^-\gamma$
------------------------	-----	------	----------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

13.6 ±1.1 ±1.3	203	<sup>1</sup> AUBERT	07AK BABR	10.6 $e^+e^- \rightarrow \pi^0\pi^0K^+K^-\gamma$
----------------	-----	---------------------	-----------	--

<sup>1</sup> Superseded by LEES 12F.

$\Gamma(\pi^0\pi^0K_S^0K_L^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{148}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

<b>10.3±2.3±0.5</b>	47	LEES	17A BABR	$e^+e^- \rightarrow K_S^0K_L^0\pi^0\pi^0\gamma$
---------------------	----	------	----------	---

$\Gamma(K_S^0K_L^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{152}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

<b>11.4±1.3±0.6</b>	182	LEES	17A BABR	$e^+e^- \rightarrow K_S^0K_L^0\pi^0\gamma$
---------------------	-----	------	----------	--

$\Gamma(K^*(892)^0\bar{K}^0 + \text{c.c.} \rightarrow K_S^0K_L^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{153}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

<b>6.7±0.9±0.4</b>	106	LEES	17A BABR	$e^+e^- \rightarrow K_S^0K_L^0\pi^0\gamma$
--------------------	-----	------	----------	--

$\Gamma(K_2^*(1430)^0\bar{K}^0 + \text{c.c.} \rightarrow K_S^0K_L^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{154}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

<b>2.4±0.7±0.1</b>	37	LEES	17A BABR	$e^+e^- \rightarrow K_S^0K_L^0\pi^0\gamma$
--------------------	----	------	----------	--

$\Gamma(K_S^0K_L^0\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{155}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

<b>8.0±1.8±0.4</b>	45	LEES	17A BABR	$e^+e^- \rightarrow K_S^0K_L^0\eta\gamma$
--------------------	----	------	----------	---

$\Gamma(2(\pi^+\pi^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{156}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

<b>20.4±0.9±0.4</b>		LEES	12E BABR	10.6 $e^+e^- \rightarrow 2\pi^+2\pi^-\gamma$
---------------------	--	------	----------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

19.5±1.4±1.3	270	<sup>1</sup> AUBERT	05D BABR	10.6 $e^+e^- \rightarrow 2(\pi^+\pi^-)\gamma$
--------------	-----	---------------------	----------	---

<sup>1</sup> Superseded by LEES 12E.

$\Gamma(3(\pi^+\pi^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{157}\Gamma_5/\Gamma$

VALUE ( $10^{-2}$ keV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------------------	------	-------------	------	---------

<b>2.37±0.16±0.14</b>	496	AUBERT	06D BABR	10.6 $e^+e^- \rightarrow 3(\pi^+\pi^-)\gamma$
-----------------------	-----	--------	----------	---

$\Gamma(2(\pi^+\pi^-\pi^0)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{158}\Gamma_5/\Gamma$

VALUE ( $10^{-2}$ keV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>8.9±0.5±1.0</b>	761	AUBERT	06D BABR	10.6 $e^+e^- \rightarrow 2(\pi^+\pi^-\pi^0)\gamma$

$\Gamma(2(\pi^+\pi^-\eta)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{159}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>13.1±2.4±0.1</b>	85	<sup>1</sup> AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow 2(\pi^+\pi^-\eta)\gamma$

<sup>1</sup> AUBERT 07AU reports  $[\Gamma(J/\psi(1S) \rightarrow 2(\pi^+\pi^-\eta)) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] = 5.16 \pm 0.85 \pm 0.39$  eV which we divide by our best value  $B(\eta \rightarrow 2\gamma) = (39.41 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\pi^+\pi^-\pi^0\pi^0\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{161}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>12.8±1.8±2.0</b>	203	LEES	18E BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0\eta\gamma$

$\Gamma(\omega\pi^0\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{60}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.90±0.96±0.01</b>	27	<sup>1</sup> LEES	18E BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0\eta\gamma$

<sup>1</sup> LEES 18E reports  $[\Gamma(J/\psi(1S) \rightarrow \omega\pi^0\eta) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = 1.7 \pm 0.8 \pm 0.3$  eV which we divide by our best value  $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.3 \pm 0.6) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\rho^\pm\pi^\mp\pi^0\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{162}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>10.5±4.1±1.6</b>	168	LEES	18E BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0\eta\gamma$

$\Gamma(p\bar{p}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{163}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>11.9±0.6 OUR AVERAGE</b>		Error includes scale factor of 1.8. See the ideogram below.		

11.3±0.4±0.3	821	<sup>1</sup> LEES	130 BABR	$e^+e^- \rightarrow p\bar{p}\gamma$
12.9±0.4±0.4	918	<sup>2</sup> LEES	13Y BABR	$e^+e^- \rightarrow p\bar{p}\gamma$
9.7±1.7		<sup>3</sup> ARMSTRONG	93B E760	$\bar{p}p \rightarrow e^+e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

12.0±0.6±0.5	438	<sup>4</sup> AUBERT	06B BABR	$e^+e^- \rightarrow p\bar{p}\gamma$
--------------	-----	---------------------	----------	-------------------------------------


<sup>1</sup> ISR photon reconstructed in the detector

<sup>2</sup> ISR photon undetected

<sup>3</sup> Using  $\Gamma_{\text{total}} = 85.5^{+6.1}_{-5.8}$  MeV.

<sup>4</sup> Superseded by LEES 130

WEIGHTED AVERAGE  
 $11.9 \pm 0.6$  (Error scaled by 1.8)



Values above of weighted average, error, and scale factor are based upon the data in this ideogram only. They are not necessarily the same as our 'best' values, obtained from a least-squares constrained fit utilizing measurements of other (related) quantities as additional information.

			$\chi^2$
LEES	13O	BABR	1.4
LEES	13Y	BABR	3.2
ARMSTRONG	93B	E760	1.7
			6.2
(Confidence Level = 0.044)			

$$\Gamma(p\bar{p}) \times \Gamma(e^+e^-) / \Gamma_{\text{total}} \text{ (eV)}$$

$$\Gamma(\Sigma^0 \bar{\Sigma}^0) \times \Gamma(e^+e^-) / \Gamma_{\text{total}} \quad \Gamma_{176} \Gamma_5 / \Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
<b><math>6.4 \pm 1.2 \pm 0.6</math></b>	AUBERT	07BD BABR	$10.6 e^+ e^- \rightarrow \Sigma^0 \bar{\Sigma}^0 \gamma$

$$\Gamma(2(\pi^+ \pi^-) K^+ K^-) \times \Gamma(e^+e^-) / \Gamma_{\text{total}} \quad \Gamma_{177} \Gamma_5 / \Gamma$$

VALUE ( $10^{-2}$ keV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>2.75 \pm 0.23 \pm 0.17</math></b>	205	AUBERT	06D BABR	$10.6 e^+ e^- \rightarrow K^+ K^- 2(\pi^+ \pi^-) \gamma$

$$\Gamma(\Lambda \bar{\Lambda}) \times \Gamma(e^+e^-) / \Gamma_{\text{total}} \quad \Gamma_{183} \Gamma_5 / \Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
<b><math>10.7 \pm 0.9 \pm 0.7</math></b>	AUBERT	07BD BABR	$10.6 e^+ e^- \rightarrow \Lambda \bar{\Lambda} \gamma$

$$\Gamma(2(K^+ K^-)) \times \Gamma(e^+e^-) / \Gamma_{\text{total}} \quad \Gamma_{186} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>4.00 \pm 0.33 \pm 0.29</math></b>	$287 \pm 24$	LEES	12F BABR	$10.6 e^+ e^- \rightarrow 2(K^+ K^-) \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$4.11 \pm 0.39 \pm 0.30$	$156 \pm 15$	<sup>1</sup> AUBERT	07AK BABR	$10.6 e^+ e^- \rightarrow 2(K^+ K^-) \gamma$
$4.0 \pm 0.7 \pm 0.6$	38	<sup>2</sup> AUBERT	05D BABR	$10.6 e^+ e^- \rightarrow 2(K^+ K^-) \gamma$

<sup>1</sup> Superseded by LEES 12F.

<sup>2</sup> Superseded by AUBERT 07AK.

**$\Gamma(K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$**   **$\Gamma_{188}\Gamma_5/\Gamma$**

<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-------------------	-------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.78±0.11±0.05	462	<sup>1</sup> LEES	15J BABR	$e^+e^- \rightarrow K^+K^-\gamma$
1.94±0.11±0.05	462	<sup>2</sup> LEES	15J BABR	$e^+e^- \rightarrow K^+K^-\gamma$
1.42±0.23±0.08	51	<sup>3</sup> LEES	13Q BABR	$e^+e^- \rightarrow K^+K^-\gamma$

<sup>1</sup>  $\sin\phi > 0$ .

<sup>2</sup>  $\sin\phi < 0$ .

<sup>3</sup> Interference with non-resonant  $K^+K^-$  production not taken into account.

**$J/\psi(1S)$  BRANCHING RATIOS**

For the first four branching ratios, see also the partial widths, and (partial widths)  $\times \Gamma(e^+e^-)/\Gamma_{\text{total}}$  above.

**$\Gamma(\text{hadrons})/\Gamma_{\text{total}}$**   **$\Gamma_1/\Gamma$**

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

**0.877±0.005 OUR AVERAGE**

0.878±0.005	BAI	95B BES	$e^+e^-$
0.86 ±0.02	BOYARSKI	75 MRK1	$e^+e^-$

**$\Gamma(\text{virtual } \gamma \rightarrow \text{hadrons})/\Gamma_{\text{total}}$**   **$\Gamma_2/\Gamma$**

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

**0.135±0.003** <sup>1,2</sup> SETH 04 RVUE  $e^+e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.17 ±0.02	<sup>1</sup> BOYARSKI	75 MRK1	$e^+e^-$
------------	-----------------------	---------	----------

<sup>1</sup> Included in  $\Gamma(\text{hadrons})/\Gamma_{\text{total}}$ .

<sup>2</sup> Using  $B(J/\psi \rightarrow \ell^+\ell^-) = (5.90 \pm 0.09)\%$  from RPP-2002 and  $R = 2.28 \pm 0.04$  determined by a fit to data from BAI 00 and BAI 02C.

**$\Gamma(ggg)/\Gamma_{\text{total}}$**   **$\Gamma_3/\Gamma$**

<u>VALUE (units <math>10^{-2}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

**64.1±1.0** 6 M <sup>1</sup> BESSON 08 CLEO  $\psi(2S) \rightarrow \pi^+\pi^- + \text{hadrons}$

<sup>1</sup> Calculated using the value  $\Gamma(\gamma gg)/\Gamma(ggg) = 0.137 \pm 0.001 \pm 0.016 \pm 0.004$  from BESSON 08 and the PDG 08 values of  $B(\ell^+\ell^-)$ ,  $B(\text{virtual } \gamma \rightarrow \text{hadrons})$ , and  $B(\gamma\eta_c)$ . The statistical error is negligible and the systematic error is partially correlated with that of  $\Gamma(\gamma gg)/\Gamma_{\text{total}}$  measurement of BESSON 08.

**$\Gamma(\gamma gg)/\Gamma_{\text{total}}$**   **$\Gamma_4/\Gamma$**

<u>VALUE (units <math>10^{-2}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

**8.79±1.05** 200 k <sup>1</sup> BESSON 08 CLEO  $\psi(2S) \rightarrow \pi^+\pi^-\gamma + \text{hadrons}$

<sup>1</sup> Calculated using the value  $\Gamma(\gamma gg)/\Gamma(ggg) = 0.137 \pm 0.001 \pm 0.016 \pm 0.004$  from BESSON 08 and the value of  $\Gamma(ggg)/\Gamma_{\text{total}}$ . The statistical error is negligible and the systematic error is partially correlated with that of  $\Gamma(ggg)/\Gamma_{\text{total}}$  measurement of BESSON 08.

### $\Gamma(\gamma g g)/\Gamma(g g g)$

$\Gamma_4/\Gamma_3$

<u>VALUE (units <math>10^{-2}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>13.7±0.1±0.7</b>	6 M	BESSION	08	CLEO $\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$

### $\Gamma(e^+ e^-)/\Gamma_{\text{total}}$

$\Gamma_5/\Gamma$

<u>VALUE (units <math>10^{-2}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>5.971±0.032 OUR AVERAGE</b>				
5.983±0.007±0.037	720k	ABLIKIM	13R	BES3 $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
5.945±0.067±0.042	15k	LI	05C	CLEO $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
5.90 ±0.05 ±0.10		BAI	98D	BES $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
6.09 ±0.33		BAI	95B	BES $e^+ e^-$
5.92 ±0.15 ±0.20		COFFMAN	92	MRK3 $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
6.9 ±0.9		BOYARSKI	75	MRK1 $e^+ e^-$

### $\Gamma(e^+ e^- \gamma)/\Gamma_{\text{total}}$

$\Gamma_6/\Gamma$

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>8.8±1.3±0.4</b>	<sup>1</sup> ARMSTRONG	96	E760 $\bar{p} p \rightarrow e^+ e^- \gamma$

<sup>1</sup> For  $E_\gamma > 100$  MeV.

### $\Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$

$\Gamma_7/\Gamma$

<u>VALUE (units <math>10^{-2}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>5.961±0.033 OUR AVERAGE</b>				
5.973±0.007±0.038	770k	ABLIKIM	13R	BES3 $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
5.960±0.065±0.050	17k	LI	05C	CLEO $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
5.84 ±0.06 ±0.10		BAI	98D	BES $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
6.08 ±0.33		BAI	95B	BES $e^+ e^-$
5.90 ±0.15 ±0.19		COFFMAN	92	MRK3 $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
6.9 ±0.9		BOYARSKI	75	MRK1 $e^+ e^-$

### $\Gamma(e^+ e^-)/\Gamma(\mu^+ \mu^-)$

$\Gamma_5/\Gamma_7$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>1.0016±0.0031 OUR AVERAGE</b>			
1.0022±0.0044±0.0048	<sup>1</sup> AULCHENKO	14	KEDR $3.097 e^+ e^- \rightarrow e^+ e^-, \mu^+ \mu^-$
1.0017±0.0017±0.0033	<sup>2</sup> ABLIKIM	13R	BES3 $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
1.002 ±0.021 ±0.013	<sup>3</sup> ANASHIN	10	KEDR $3.097 e^+ e^- \rightarrow e^+ e^-, \mu^+ \mu^-$
0.997 ±0.012 ±0.006	LI	05C	CLEO $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1.011 ±0.013 ±0.016	BAI	98D	BES $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
1.00 ±0.07	BAI	95B	BES $e^+ e^-$
1.00 ±0.05	BOYARSKI	75	MRK1 $e^+ e^-$
0.91 ±0.15	ESPOSITO	75B	FRAM $e^+ e^-$
0.93 ±0.10	FORD	75	SPEC $e^+ e^-$

<sup>1</sup> From 235.3k  $J/\psi \rightarrow e^+ e^-$  and 156.6k  $J/\psi \rightarrow \mu^+ \mu^-$  observed events.

<sup>2</sup> Not independent of the corresponding measurements of  $\Gamma(e^+ e^-)/\Gamma_{\text{total}}$  and  $\Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$ .

<sup>3</sup> Not independent of the corresponding measurements of  $\Gamma(e^+ e^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$  and  $\Gamma(\mu^+ \mu^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ .




————— HADRONIC DECAYS —————

$\Gamma(\rho\pi)/\Gamma_{\text{total}}$

$\Gamma_8/\Gamma$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.69 ± 0.15 OUR AVERAGE</b>		Error includes scale factor of 2.4. See the ideogram below.		
2.18 ± 0.19		1,2 AUBERT,B	04N BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \gamma$
2.184 ± 0.005 ± 0.201	220k	2,3 BAI	04H BES	$e^+ e^- \rightarrow J/\psi \rightarrow \pi^+ \pi^- \pi^0$
2.091 ± 0.021 ± 0.116		2,4 BAI	04H BES	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$
1.21 ± 0.20		BAI	96D BES	$e^+ e^- \rightarrow \rho\pi$
1.42 ± 0.01 ± 0.19		COFFMAN	88 MRK3	$e^+ e^-$
1.3 ± 0.3	150	FRANKLIN	83 MRK2	$e^+ e^-$
1.6 ± 0.4	183	ALEXANDER	78 PLUT	$e^+ e^-$
1.33 ± 0.21		BRANDELIK	78B DASP	$e^+ e^-$
1.0 ± 0.2	543	BARTEL	76 CNTR	$e^+ e^-$
1.3 ± 0.3	153	JEAN-MARIE	76 MRK1	$e^+ e^-$

- <sup>1</sup> From the ratio of  $\Gamma(e^+ e^-) B(\pi^+ \pi^- \pi^0)$  and  $\Gamma(e^+ e^-) B(\mu^+ \mu^-)$  (AUBERT 04).
- <sup>2</sup> Not independent of their  $B(\pi^+ \pi^- \pi^0)$ .
- <sup>3</sup> From  $J/\psi \rightarrow \pi^+ \pi^- \pi^0$  events directly.
- <sup>4</sup> Obtained comparing the rates for  $\pi^+ \pi^- \pi^0$  and  $\mu^+ \mu^-$ , using  $J/\psi$  events produced via  $\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$  and with  $B(J/\psi \rightarrow \mu^+ \mu^-) = 5.88 \pm 0.10\%$ .



$$\Gamma(\rho\pi)/\Gamma(\pi^+\pi^-\pi^0) \qquad \Gamma_8/\Gamma_{135}$$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>1.142±0.011±0.026</b>	20K	<sup>1</sup> LEES	17C BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.331±0.033	20K	<sup>2</sup> LEES	17C BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$

<sup>1</sup> From a Dalitz plot analysis in an isobar model.<sup>2</sup> From a Dalitz plot analysis in a Veneziano model.
$$\Gamma(\rho^0\pi^0)/\Gamma(\rho\pi) \qquad \Gamma_9/\Gamma_8$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.328±0.005±0.027</b>	COFFMAN	88 MRK3	$e^+e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.35 ±0.08	ALEXANDER	78 PLUT	$e^+e^-$
0.32 ±0.08	BRANDELIK	78B DASP	$e^+e^-$
0.39 ±0.11	BARTEL	76 CNTR	$e^+e^-$
0.37 ±0.09	JEAN-MARIE	76 MRK1	$e^+e^-$

$$\Gamma(\rho(1450)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma(\pi^+\pi^-\pi^0) \qquad \Gamma_{12}/\Gamma_{135}$$

<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>10.9 ±1.7 ±2.7</b>	20K	<sup>1</sup> LEES	17C BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.80±0.27	20K	<sup>2</sup> LEES	17C BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$

<sup>1</sup> From a Dalitz plot analysis in an isobar model.<sup>2</sup> From a Dalitz plot analysis in a Veneziano model.
$$\Gamma(\rho(1450)^\pm\pi^\mp \rightarrow K_S^0 K^\pm\pi^\mp)/\Gamma(K_S^0 K^\pm\pi^\mp) \qquad \Gamma_{13}/\Gamma_{151}$$

<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>6.3±0.8±0.6</b>	4K	<sup>1</sup> LEES	17C BABR	$J/\psi \rightarrow K_S^0 K^\pm\pi^\mp$

<sup>1</sup> From a Dalitz plot analysis in an isobar model.
$$\Gamma(\rho(1450)^0\pi^0 \rightarrow K^+K^-\pi^0)/\Gamma(K^+K^-\pi^0) \qquad \Gamma_{14}/\Gamma_{150}$$

<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>9.3±2.0±0.6</b>	2K	<sup>1</sup> LEES	17C BABR	$J/\psi \rightarrow K^+K^-\pi^0$

<sup>1</sup> From a Dalitz plot analysis in an isobar model.
$$\Gamma(\rho(1450)\eta'(958) \rightarrow \pi^+\pi^-\eta'(958))/\Gamma_{\text{total}} \qquad \Gamma_{15}/\Gamma$$

<u>VALUE (units 10<sup>-6</sup>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>3.28±0.55±0.44</b>	119	<sup>1</sup> ABLIKIM	17AK BES3	$J/\psi \rightarrow \pi^+\pi^-\eta'$

<sup>1</sup> From a partial wave analysis of the decay  $J/\psi \rightarrow \pi^+\pi^-\eta'$ .
$$\Gamma(\rho(1700)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma(\pi^+\pi^-\pi^0) \qquad \Gamma_{17}/\Gamma_{135}$$

<u>VALUE (units 10<sup>-3</sup>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>8±2±5</b>	20K	<sup>1</sup> LEES	17C BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
22±6	20K	<sup>2</sup> LEES	17C BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$

<sup>1</sup> From a Dalitz plot analysis in an isobar model.<sup>2</sup> From a Dalitz plot analysis in a Veneziano model.

$\Gamma(\rho(2150)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma(\pi^+\pi^-\pi^0)$   $\Gamma_{19}/\Gamma_{135}$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

<b>4± 1±20</b>	20K	<sup>1</sup> LEES	17C BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$
----------------	-----	-------------------	----------	--------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

600±250	20K	<sup>2</sup> LEES	17C BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$
---------	-----	-------------------	----------	--------------------------------------

<sup>1</sup> From a Dalitz plot analysis in an isobar model.

<sup>2</sup> From a Dalitz plot analysis in a Veneziano model.

$\Gamma(\rho_3(1690)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma(\pi^+\pi^-\pi^0)$   $\Gamma_{20}/\Gamma_{135}$

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.0±0.8	20K	<sup>1</sup> LEES	17C BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$
---------	-----	-------------------	----------	--------------------------------------

<sup>1</sup> From a Dalitz plot analysis in a Veneziano model.

$\Gamma(a_2(1320)\rho)/\Gamma_{\text{total}}$   $\Gamma_{21}/\Gamma$

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

**10.9±2.2 OUR AVERAGE**

11.7±0.7±2.5	7584	AUGUSTIN	89 DM2	$J/\psi \rightarrow \rho^0\rho^\pm\pi^\mp$
--------------	------	----------	--------	--

8.4±4.5	36	VANNUCCI	77 MRK1	$e^+e^- \rightarrow 2(\pi^+\pi^-)\pi^0$
---------	----	----------	---------	---

$\Gamma(\omega\pi^+\pi^+\pi^-\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{22}/\Gamma$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

<b>85±34</b>	140	VANNUCCI	77 MRK1	$e^+e^- \rightarrow 3(\pi^+\pi^-)\pi^0$
--------------	-----	----------	---------	---

$\Gamma(\omega\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{24}/\Gamma$

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

**7.2±1.0 OUR AVERAGE**

7.0±1.6	18058	AUGUSTIN	89 DM2	$J/\psi \rightarrow 2(\pi^+\pi^-)\pi^0$
---------	-------	----------	--------	---

7.8±1.6	215	BURMESTER	77D PLUT	$e^+e^-$
---------	-----	-----------	----------	----------

6.8±1.9	348	VANNUCCI	77 MRK1	$e^+e^- \rightarrow 2(\pi^+\pi^-)\pi^0$
---------	-----	----------	---------	---

$\Gamma(\omega f_2(1270))/\Gamma_{\text{total}}$   $\Gamma_{25}/\Gamma$

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

**4.3±0.6 OUR AVERAGE**

4.3±0.2±0.6	5860	AUGUSTIN	89 DM2	$e^+e^-$
-------------	------	----------	--------	----------

4.0±1.6	70	BURMESTER	77D PLUT	$e^+e^-$
---------	----	-----------	----------	----------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.9±0.8	81	VANNUCCI	77 MRK1	$e^+e^- \rightarrow 2(\pi^+\pi^-)\pi^0$
---------	----	----------	---------	---

$\Gamma(K^*(892)^0\bar{K}^*(892)^0)/\Gamma_{\text{total}}$   $\Gamma_{26}/\Gamma$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<5	90	VANNUCCI	77 MRK1	$e^+e^- \rightarrow \pi^+\pi^-K^+K^-$
----	----	----------	---------	---------------------------------------

$\Gamma(K^*(892)^\pm K^*(892)^\mp)/\Gamma_{\text{total}}$   $\Gamma_{27}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$1.00 \pm 0.19^{+0.11}_{-0.32}$	323	ABLIKIM	10E BES2	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp \pi^0$

$\Gamma(K^*(892)^\pm K^*(700)^\mp)/\Gamma_{\text{total}}$   $\Gamma_{28}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$1.09 \pm 0.18^{+0.94}_{-0.54}$	655	ABLIKIM	10E BES2	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp \pi^0$

$\Gamma(\eta K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}$   $\Gamma_{32}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$1.15 \pm 0.13 \pm 0.22$	209	ABLIKIM	10C BES2	$J/\psi \rightarrow \eta K^+ \pi^- K^- \pi^+$

$\Gamma(K^*(1410) \bar{K} + \text{c.c.} \rightarrow K^\pm K^\mp \pi^0)/\Gamma(K^+ K^- \pi^0)$   $\Gamma_{38}/\Gamma_{150}$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$2.3 \pm 1.1 \pm 0.7$	2K	<sup>1</sup> LEES	17C BABR	$J/\psi \rightarrow K^+ K^- \pi^0$

<sup>1</sup> From a Dalitz plot analysis in an isobar model.

$\Gamma(K^*(1410) \bar{K} + \text{c.c.} \rightarrow K_S^0 K^\pm \pi^\mp)/\Gamma(K_S^0 K^\pm \pi^\mp)$   $\Gamma_{39}/\Gamma_{151}$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$1.5 \pm 0.5 \pm 0.9$	4K	<sup>1</sup> LEES	17C BABR	$J/\psi \rightarrow K_S^0 K^\pm \pi^\mp$

<sup>1</sup> From a Dalitz plot analysis in an isobar model.

$\Gamma(K_2^*(1430) \bar{K} + \text{c.c.} \rightarrow K^\pm K^\mp \pi^0)/\Gamma(K^+ K^- \pi^0)$   $\Gamma_{41}/\Gamma_{150}$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$3.5 \pm 1.3 \pm 0.9$	2K	<sup>1</sup> LEES	17C BABR	$J/\psi \rightarrow K^+ K^- \pi^0$

<sup>1</sup> From a Dalitz plot analysis in an isobar model.

$\Gamma(K_2^*(1430) \bar{K} + \text{c.c.} \rightarrow K_S^0 K^\pm \pi^\mp)/\Gamma(K_S^0 K^\pm \pi^\mp)$   $\Gamma_{42}/\Gamma_{151}$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$7.1 \pm 1.3 \pm 1.2$	4K	<sup>1</sup> LEES	17C BABR	$J/\psi \rightarrow K_S^0 K^\pm \pi^\mp$

<sup>1</sup> From a Dalitz plot analysis in an isobar model.

$\Gamma(K^*(892)^0 \bar{K}_2^*(1430)^0 + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{43}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$6.7 \pm 2.6$	40	VANNUCCI 77	MRK1	$e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^-$
---------------	----	-------------	------	---

$\Gamma(\omega K^*(892) \bar{K} + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{47}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**61 ± 9 OUR AVERAGE**

$62.0 \pm 6.8 \pm 10.6$	$899 \pm 98$	ABLIKIM	08E BES2	$J/\psi \rightarrow \omega K_S^0 K^\pm \pi^\mp$
$65.3 \pm 10.2 \pm 13.5$	$176 \pm 28$	ABLIKIM	08E BES2	$J/\psi \rightarrow \omega K^+ K^- \pi^0$
$53 \pm 14 \pm 14$	$530 \pm 140$	BECKER	87 MRK3	$e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(\overline{K} K^*(892) + \text{c.c.} \rightarrow K_S^0 K^\pm \pi^\mp) / \Gamma(K_S^0 K^\pm \pi^\mp)$   $\Gamma_{49} / \Gamma_{151}$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>90.5 ± 0.9 ± 3.8</b>	4K	<sup>1</sup> LEES	17C BABR	$J/\psi \rightarrow K_S^0 K^\pm \pi^\mp$

<sup>1</sup>From a Dalitz plot analysis in an isobar model.

$\Gamma(K^+ K^*(892)^- + \text{c.c.}) / \Gamma_{\text{total}}$   $\Gamma_{50} / \Gamma$

VALUE (units 10 <sup>-3</sup> )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>5.0 ± 0.4 OUR AVERAGE</b>				
4.57 ± 0.17 ± 0.70	2285	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
5.26 ± 0.13 ± 0.53		COFFMAN	88 MRK3	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp,$ $K^+ K^- \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.6 ± 0.6	24	FRANKLIN	83 MRK2	$J/\psi \rightarrow K^+ K^- \pi^0$
3.2 ± 0.6	48	VANNUCCI	77 MRK1	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp$
4.1 ± 1.2	39	BRAUNSCH...	76 DASP	$J/\psi \rightarrow K^\pm X$

$\Gamma(K^+ K^*(892)^- + \text{c.c.} \rightarrow K^+ K^- \pi^0) / \Gamma(K^+ K^- \pi^0)$   $\Gamma_{51} / \Gamma_{150}$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>92.4 ± 1.5 ± 3.4</b>	2K	<sup>1</sup> LEES	17C BABR	$J/\psi \rightarrow K^+ K^- \pi^0$

<sup>1</sup>From a Dalitz plot analysis in an isobar model.

$\Gamma(K^0 \overline{K}^*(892)^0 + \text{c.c.}) / \Gamma_{\text{total}}$   $\Gamma_{53} / \Gamma$

VALUE (units 10 <sup>-3</sup> )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.2 ± 0.4 OUR AVERAGE</b>				
3.96 ± 0.15 ± 0.60	1192	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
4.33 ± 0.12 ± 0.45		COFFMAN	88 MRK3	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.7 ± 0.6	45	VANNUCCI	77 MRK1	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp$
-----------	----	----------	---------	--

$\Gamma(K_1(1400)^\pm K^\mp) / \Gamma_{\text{total}}$   $\Gamma_{55} / \Gamma$

VALUE (units 10 <sup>-3</sup> )	DOCUMENT ID	TECN	COMMENT
<b>3.8 ± 0.8 ± 1.2</b>	<sup>1</sup> BAI	99C BES	$e^+ e^-$

<sup>1</sup>Assuming  $B(K_1(1400) \rightarrow K^* \pi) = 0.94 \pm 0.06$

$\Gamma(\overline{K}^*(892)^0 K^+ \pi^- + \text{c.c.}) / \Gamma_{\text{total}}$   $\Gamma_{56} / \Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen <sup>1</sup> ABLIKIM 06C BES2  $J/\psi \rightarrow \overline{K}^*(892)^0 K^+ \pi^-$

<sup>1</sup>A  $K_0^*(700)$  is observed by ABLIKIM 06C in the  $K^+ \pi^-$  mass spectrum of the  $\overline{K}^*(892)^0 K^+ \pi^-$  final state against the  $\overline{K}^*(892)$ . A corresponding branching fraction of the  $J/\psi(1S)$  is not presented.

$\Gamma(\omega \pi^0 \pi^0) / \Gamma_{\text{total}}$   $\Gamma_{59} / \Gamma$

VALUE (units 10 <sup>-3</sup> )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.4 ± 0.3 ± 0.7</b>	509	AUGUSTIN	89 DM2	$J/\psi \rightarrow \pi^+ \pi^- 3\pi^0$

$\Gamma(b_1(1235)^\pm \pi^\mp)/\Gamma_{\text{total}}$   $\Gamma_{61}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>30 ± 5 OUR AVERAGE</b>				
31 ± 6	4600	AUGUSTIN	89 DM2	$J/\psi \rightarrow 2(\pi^+ \pi^-) \pi^0$
29 ± 7	87	BURMESTER	77D PLUT	$e^+ e^-$

$\Gamma(\omega K^\pm K_S^0 \pi^\mp)/\Gamma_{\text{total}}$   $\Gamma_{62}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>34 ± 5 OUR AVERAGE</b>				
37.7 ± 0.8 ± 5.8	1972 ± 41	ABLIKIM	08E BES2	$e^+ e^- \rightarrow J/\psi$
29.5 ± 1.4 ± 7.0	879 ± 41	BECKER	87 MRK3	$e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(b_1(1235)^0 \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{63}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>23 ± 3 ± 5</b>				
	229	AUGUSTIN	89 DM2	$e^+ e^-$

$\Gamma(\eta K^\pm K_S^0 \pi^\mp)/\Gamma_{\text{total}}$   $\Gamma_{64}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>21.8 ± 2.2 ± 3.4</b>				
	232 ± 23	ABLIKIM	08E BES2	$e^+ e^- \rightarrow J/\psi$

$\Gamma(\eta' K^{*0} \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{34}/\Gamma$

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
<b>1.66 ± 0.03 ± 0.21</b>			
	<sup>1</sup> ABLIKIM	18AB BES3	$J/\psi \rightarrow \eta' K^* \bar{K}$

<sup>1</sup> From  $\eta' K_S^0 K^\pm \pi^\mp$ .

$\Gamma(\eta' K^{*\pm} K^\mp)/\Gamma_{\text{total}}$   $\Gamma_{33}/\Gamma$

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
<b>1.48 ± 0.13 OUR AVERAGE</b>			
1.50 ± 0.02 ± 0.19	<sup>1</sup> ABLIKIM	18AB BES3	$J/\psi \rightarrow \eta' K^* \bar{K}$
1.47 ± 0.03 ± 0.17	<sup>2</sup> ABLIKIM	18AB BES3	$J/\psi \rightarrow \eta' K^* \bar{K}$

<sup>1</sup> From  $\eta' K^+ K^- \pi^0$ .  
<sup>2</sup> From  $\eta' K_S^0 K^\pm \pi^\mp$ .

$\Gamma(\eta' h_1(1415) \rightarrow \eta' K^{*\pm} K^\mp)/\Gamma_{\text{total}}$   $\Gamma_{36}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.51 ± 0.09 ± 0.21</b>				
	1.0k	<sup>1</sup> ABLIKIM	18AB BES3	$J/\psi \rightarrow \eta' h_1 \rightarrow \eta' K^* \bar{K}$

<sup>1</sup> From  $\eta' K^+ K^- \pi^0$ .

$\Gamma(\eta' h_1(1415) \rightarrow \eta' K^* \bar{K} + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{35}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.16 ± 0.12 ± 0.29</b>				
	1.1k	<sup>1</sup> ABLIKIM	18AB BES3	$J/\psi \rightarrow \eta' h_1 \rightarrow \eta' K^* \bar{K}$

<sup>1</sup> From  $\eta' K_S^0 K^\pm \pi^\mp$ .

$\Gamma(\phi K^*(892)\bar{K} + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{65}/\Gamma$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>21.8 ± 2.3 OUR AVERAGE</b>				
20.8 ± 2.7 ± 3.9	195 ± 25	ABLIKIM	08E BES2	$J/\psi \rightarrow \phi K_S^0 K^\pm \pi^\mp$
29.6 ± 3.7 ± 4.7	238 ± 30	ABLIKIM	08E BES2	$J/\psi \rightarrow \phi K^+ K^- \pi^0$
20.7 ± 2.4 ± 3.0		FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
20 ± 3 ± 3	155 ± 20	BECKER	87 MRK3	$e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(\omega K\bar{K})/\Gamma_{\text{total}}$   $\Gamma_{66}/\Gamma$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>19 ± 4 OUR AVERAGE</b>				
19.8 ± 2.1 ± 3.9		<sup>1</sup> FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
16 ± 10	22	FELDMAN	77 MRK1	$e^+ e^-$

<sup>1</sup> Addition of  $\omega K^+ K^-$  and  $\omega K^0 \bar{K}^0$  branching ratios.

$\Gamma(\omega f_0(1710) \rightarrow \omega K\bar{K})/\Gamma_{\text{total}}$   $\Gamma_{67}/\Gamma$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>4.8 ± 1.1 ± 0.3</b>	<sup>1,2</sup> FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$

<sup>1</sup> Includes unknown branching fraction  $f_0(1710) \rightarrow K\bar{K}$ .

<sup>2</sup> Addition of  $f_0(1710) \rightarrow K^+ K^-$  and  $f_0(1710) \rightarrow K^0 \bar{K}^0$  branching ratios.

$\Gamma(\phi 2(\pi^+ \pi^-))/\Gamma_{\text{total}}$   $\Gamma_{68}/\Gamma$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>16.0 ± 1.0 ± 3.0</b>	FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$

$\Gamma(\Delta(1232)^{++} \bar{p} \pi^-)/\Gamma_{\text{total}}$   $\Gamma_{69}/\Gamma$

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>1.58 ± 0.23 ± 0.40</b>	332	EATON	84 MRK2	$e^+ e^-$


$\Gamma(\omega \eta)/\Gamma_{\text{total}}$   $\Gamma_{70}/\Gamma$

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>1.74 ± 0.20 OUR AVERAGE</b> Error includes scale factor of 1.6. See the ideogram below.				
2.352 ± 0.273	5k	<sup>1</sup> ABLIKIM	06F BES2	$J/\psi \rightarrow \omega \eta$
1.44 ± 0.40 ± 0.14	13	<sup>2</sup> AUBERT	06D BABR	$10.6 e^+ e^- \rightarrow \omega \eta \gamma$
1.43 ± 0.10 ± 0.21	378	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
1.71 ± 0.08 ± 0.20		COFFMAN	88 MRK3	$e^+ e^- \rightarrow 3\pi \eta$

<sup>1</sup> Using  $B(\eta \rightarrow 2\gamma) = (39.43 \pm 0.26)\%$ ,  $B(\eta \rightarrow \pi^+ \pi^- \pi^0) = 22.6 \pm 0.4\%$ ,  $B(\eta \rightarrow \pi^+ \pi^- \gamma) = 4.68 \pm 0.11\%$ , and  $B(\omega \rightarrow \pi^+ \pi^- \pi^0) = (89.1 \pm 0.7)\%$ .

<sup>2</sup> Using  $\Gamma(J/\psi \rightarrow e^+ e^-) = 5.52 \pm 0.14 \pm 0.04$  keV.

WEIGHTED AVERAGE  
 $1.74 \pm 0.20$  (Error scaled by 1.6)



$\Gamma(\phi K \bar{K})/\Gamma_{\text{total}}$

$\Gamma_{71}/\Gamma$


VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>17.7 \pm 1.6</math></b>	<b>OUR AVERAGE</b>	Error includes scale factor of 1.3. See the ideogram below.		
$16.6 \pm 1.9 \pm 1.2$	$163 \pm 19$	LEES	12F BABR	$10.6 e^+ e^- \rightarrow 2(K^+ K^-) \gamma$
$21.4 \pm 0.4 \pm 2.2$		ABLIKIM	05 BES2	$J/\psi \rightarrow \phi \pi^+ \pi^-$
$48^{+20}_{-16} \pm 6$	$9.0^{+3.7}_{-3.0}$	<sup>1,2</sup> HUANG	03 BELL	$B^+ \rightarrow (\phi K^+ K^-) K^+$
$14.6 \pm 0.8 \pm 2.1$		<sup>3</sup> FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
$18 \pm 8$	14	FELDMAN	77 MRK1	$e^+ e^-$

<sup>1</sup> We have multiplied  $K^+ K^-$  measurement by 2 to obtain  $K \bar{K}$ .

<sup>2</sup> Using  $B(B^+ \rightarrow J/\psi K^+) = (1.01 \pm 0.05) \times 10^{-3}$ .

<sup>3</sup> Addition of  $\phi K^+ K^-$  and  $\phi K^0 \bar{K}^0$  branching ratios.





$\Gamma(\phi f_0(1710) \rightarrow \phi K \bar{K}) / \Gamma_{\text{total}}$   $\Gamma_{73} / \Gamma$

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
<b><math>3.6 \pm 0.2 \pm 0.6</math></b>	1,2 FALVARD	88 DM2	$J/\psi \rightarrow$ hadrons

<sup>1</sup> Including interference with  $f_2'(1525)$ .

<sup>2</sup> Includes unknown branching fraction  $f_0(1710) \rightarrow K \bar{K}$ .

$\Gamma(\phi f_2(1270)) / \Gamma_{\text{total}}$   $\Gamma_{75} / \Gamma$

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
• • •				We do not use the following data for averages, fits, limits, etc. • • •
< 0.45	90	FALVARD	88 DM2	$J/\psi \rightarrow$ hadrons
< 0.37	90	VANNUCCI	77 MRK1	$e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^-$

$\Gamma(\Delta(1232)^{++} \bar{\Delta}(1232)^{--}) / \Gamma_{\text{total}}$   $\Gamma_{76} / \Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>1.10 \pm 0.09 \pm 0.28</math></b>	233	EATON	84 MRK2	$e^+ e^-$

$\Gamma(\Sigma(1385)^- \bar{\Sigma}(1385)^+ \text{ (or c.c.)}) / \Gamma_{\text{total}}$   $\Gamma_{77} / \Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>1.16 \pm 0.05</math> OUR AVERAGE</b>				
$1.096 \pm 0.012 \pm 0.071$	43K	ABLIKIM	16L BES3	$J/\psi \rightarrow \Sigma(1385)^- \bar{\Sigma}(1385)^+$
$1.258 \pm 0.014 \pm 0.078$	53k	ABLIKIM	16L BES3	$J/\psi \rightarrow \Sigma(1385)^+ \bar{\Sigma}(1385)^-$
$1.23 \pm 0.07 \pm 0.30$	0.8k	ABLIKIM	12P BES2	$J/\psi \rightarrow \Sigma(1385)^- \bar{\Sigma}(1385)^+$
$1.50 \pm 0.08 \pm 0.38$	1k	ABLIKIM	12P BES2	$J/\psi \rightarrow \Sigma(1385)^+ \bar{\Sigma}(1385)^-$

1.00 ± 0.04 ± 0.21	0.6k	HENRARD	87	DM2	$e^+e^- \rightarrow \Sigma^{*-}$
1.19 ± 0.04 ± 0.25	0.7k	HENRARD	87	DM2	$e^+e^- \rightarrow \Sigma^{*+}$
0.86 ± 0.18 ± 0.22	56	EATON	84	MRK2	$e^+e^- \rightarrow \Sigma^{*-}$
1.03 ± 0.24 ± 0.25	68	EATON	84	MRK2	$e^+e^- \rightarrow \Sigma^{*+}$

$\Gamma(\Sigma(1385)^0 \bar{\Sigma}(1385)^0) / \Gamma_{\text{total}}$   $\Gamma_{78} / \Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.071 ± 0.009 ± 0.082</b>	103k	ABLIKIM	17E BES3	$e^+e^- \rightarrow J/\psi \rightarrow$ hadrons

$\Gamma(\phi f'_2(1525)) / \Gamma_{\text{total}}$   $\Gamma_{80} / \Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>8 ± 4 OUR AVERAGE</b>				Error includes scale factor of 2.7.
12.3 ± 0.6 ± 2.0		<sup>1,2</sup> FALVARD	88 DM2	$J/\psi \rightarrow$ hadrons
4.8 ± 1.8	46	<sup>1</sup> GIDAL	81 MRK2	$J/\psi \rightarrow K^+ K^- K^+ K^-$
<sup>1</sup> Re-evaluated using $B(f'_2(1525) \rightarrow K\bar{K}) = 0.713$ .				
<sup>2</sup> Including interference with $f_0(1710)$ .				

$\Gamma(\phi \pi^+ \pi^-) / \Gamma_{\text{total}}$   $\Gamma_{81} / \Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.94 ± 0.15 OUR AVERAGE</b>				Error includes scale factor of 1.7.
1.09 ± 0.02 ± 0.13		ABLIKIM	05 BES2	$J/\psi \rightarrow \phi \pi^+ \pi^-$
0.78 ± 0.03 ± 0.12		FALVARD	88 DM2	$J/\psi \rightarrow$ hadrons
2.1 ± 0.9	23	FELDMAN	77 MRK1	$e^+e^-$

$\Gamma(\phi K^\pm K_S^0 \pi^\mp) / \Gamma_{\text{total}}$   $\Gamma_{83} / \Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>7.2 ± 0.8 OUR AVERAGE</b>				
7.4 ± 0.6 ± 1.4	227 ± 19	ABLIKIM	08E BES2	$e^+e^- \rightarrow J/\psi$
7.4 ± 0.9 ± 1.1		FALVARD	88 DM2	$J/\psi \rightarrow$ hadrons
7 ± 0.6 ± 1.0	163 ± 15	BECKER	87 MRK3	$e^+e^- \rightarrow$ hadrons


$\Gamma(\omega f_1(1420)) / \Gamma_{\text{total}}$   $\Gamma_{84} / \Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>6.8<sup>+1.9</sup><sub>-1.6</sub> ± 1.7</b>	111 <sup>+31</sup> <sub>-26</sub>	BECKER	87 MRK3	$e^+e^- \rightarrow$ hadrons

$\Gamma(\phi \eta) / \Gamma_{\text{total}}$   $\Gamma_{85} / \Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.74 ± 0.08 OUR AVERAGE</b>				Error includes scale factor of 1.5. See the ideogram below.
0.898 ± 0.024 ± 0.089		ABLIKIM	05B BES2	$e^+e^- \rightarrow J/\psi \rightarrow$ hadr
0.64 ± 0.04 ± 0.11	346	JOUSSET	90 DM2	$J/\psi \rightarrow$ hadrons
0.661 ± 0.045 ± 0.078		COFFMAN	88 MRK3	$e^+e^- \rightarrow K^+ K^- \eta$

WEIGHTED AVERAGE  
 $0.74 \pm 0.08$  (Error scaled by 1.5)



$\Gamma(\Xi^0 \Xi^0)/\Gamma_{\text{total}}$

VALUE (units $10^{-3}$ )	EVTS
<b>1.17 ± 0.04 OUR AVERAGE</b>	
1.165 ± 0.004 ± 0.043	135K
1.20 ± 0.12 ± 0.21	206

DOCUMENT ID	TECN	COMMENT	$\Gamma_{86}/\Gamma$
ABLIKIM	17E	BES3	$e^+ e^- \rightarrow J/\psi \rightarrow$ hadrons
ABLIKIM	080	BES2	$e^+ e^- \rightarrow J/\psi$

$\Gamma(\Xi(1530)^- \Xi^+)/\Gamma_{\text{total}}$

VALUE (units $10^{-3}$ )	EVTS
<b>0.59 ± 0.09 ± 0.12</b>	75 ± 11

DOCUMENT ID	TECN	COMMENT	$\Gamma_{87}/\Gamma$
HENRARD	87	DM2	$e^+ e^-$

$\Gamma(\rho K^- \bar{\Sigma}(1385)^0)/\Gamma_{\text{total}}$

VALUE (units $10^{-3}$ )	EVTS
<b>0.51 ± 0.26 ± 0.18</b>	89


DOCUMENT ID	TECN	COMMENT	$\Gamma_{88}/\Gamma$
EATON	84	MRK2	$e^+ e^-$

$\Gamma(\omega \pi^0)/\Gamma_{\text{total}}$

VALUE (units $10^{-3}$ )	EVTS
<b>0.45 ± 0.05 OUR AVERAGE</b>	
0.538 ± 0.012 ± 0.065	2090
0.360 ± 0.028 ± 0.054	222
0.482 ± 0.019 ± 0.064	

DOCUMENT ID	TECN	COMMENT	$\Gamma_{89}/\Gamma$
Error includes scale factor of 1.4. See the ideogram below.			
<sup>1</sup> ABLIKIM	06F	BES2	$J/\psi \rightarrow \omega \pi^0$
JOUSSET	90	DM2	$J/\psi \rightarrow$ hadrons
COFFMAN	88	MRK3	$e^+ e^- \rightarrow \pi^0 \pi^+ \pi^- \pi^0$

<sup>1</sup> Using  $B(\omega \rightarrow \pi^+ \pi^- \pi^0) = (89.1 \pm 0.7)\%$ .



$\Gamma(\omega\pi^0 \rightarrow \pi^+ \pi^- \pi^0)/\Gamma(\pi^+ \pi^- \pi^0)$   $\Gamma_{90}/\Gamma_{135}$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>8 \pm 3 \pm 2</math></b>	20K	<sup>1</sup> LEES	17C BABR	$J/\psi \rightarrow \pi^+ \pi^- \pi^0$

<sup>1</sup> From a Dalitz plot analysis in an isobar model and significance  $4.9 \sigma$ .


$\Gamma(\phi\eta'(958))/\Gamma_{\text{total}}$   $\Gamma_{91}/\Gamma$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>4.6 \pm 0.5</math></b>	<b>OUR AVERAGE</b>		Error includes scale factor of 2.2. See the ideogram below.		
$5.10 \pm 0.03 \pm 0.32$		31k	ABLIKIM 16M	BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \text{hadrons}$
$5.46 \pm 0.31 \pm 0.56$			ABLIKIM 05B	BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \text{hadrons}$
$4.1 \pm 0.3 \pm 0.8$		167	JOUSSET 90	DM2	$J/\psi \rightarrow \text{hadrons}$
$3.08 \pm 0.34 \pm 0.36$			COFFMAN 88	MRK3	$e^+ e^- \rightarrow K^+ K^- \eta'$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 13	90	VANNUCCI 77	MRK1	$e^+ e^-$
------	----	-------------	------	-----------

WEIGHTED AVERAGE  
 $4.6 \pm 0.5$  (Error scaled by 2.2)



$\Gamma(\phi f_0(980)) / \Gamma_{\text{total}}$

$\Gamma_{92} / \Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>3.2 \pm 0.9</math> OUR AVERAGE</b>				Error includes scale factor of 1.9.
$4.6 \pm 0.4 \pm 0.8$		<sup>1</sup> FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
$2.6 \pm 0.6$	50	<sup>1</sup> GIDAL	81 MRK2	$J/\psi \rightarrow K^+ K^- K^+ K^-$

<sup>1</sup> Assuming  $B(f_0(980) \rightarrow \pi\pi) = 0.78$ .

$\Gamma(\phi \pi^0 f_0(980) \rightarrow \phi \pi^0 \pi^+ \pi^-) / \Gamma_{\text{total}}$

$\Gamma_{95} / \Gamma$

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>4.50 \pm 0.80 \pm 0.61</math></b>	355	ABLIKIM	15P BES3	$J/\psi \rightarrow K^+ K^- 3\pi$

$\Gamma(\phi \pi^0 f_0(980) \rightarrow \phi \pi^0 \rho^0 \pi^0) / \Gamma_{\text{total}}$

$\Gamma_{96} / \Gamma$

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>1.67 \pm 0.50 \pm 0.24</math></b>	70	ABLIKIM	15P BES3	$J/\psi \rightarrow K^+ K^- 3\pi$

$\Gamma(\eta \phi f_0(980) \rightarrow \eta \phi \pi^+ \pi^-) / \Gamma_{\text{total}}$

$\Gamma_{97} / \Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>3.23 \pm 0.75 \pm 0.73</math></b>	52	ABLIKIM	08F BES	$J/\psi \rightarrow \eta \phi f_0(980)$

$\Gamma(\phi a_0(980)^0 \rightarrow \phi \eta \pi^0) / \Gamma_{\text{total}}$

$\Gamma_{98} / \Gamma$

VALUE (units $10^{-6}$ )	DOCUMENT ID	TECN	COMMENT
<b><math>4.37 \pm 1.35</math></b>	<sup>1</sup> ABLIKIM	18D BES3	$J/\psi \rightarrow \phi \eta \pi^0$
$5.0 \pm 2.7 \pm 2.5$	<sup>2</sup> ABLIKIM	11D BES3	$J/\psi \rightarrow \phi \eta \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup> Assuming constructive interference between  $a_0(980) - f_0(980)$  mixing and electromagnetic decay. Destructive interference gives a value of  $(4.93 \pm 1.77) \times 10^{-6}$  for this branching fraction.

<sup>2</sup> Assuming  $a_0(980) - f_0(980)$  mixing and isospin breaking via  $\gamma^*$  and  $K^* K$  loops.

$\Gamma(\Xi(1530)^0 \Xi^0)/\Gamma_{\text{total}}$   $\Gamma_{99}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.32±0.12±0.07</b>	24 ± 9	HENRARD	87 DM2	$e^+ e^-$

$\Gamma(\Sigma(1385)^- \bar{\Sigma}^+ \text{ (or c.c.)})/\Gamma_{\text{total}}$   $\Gamma_{100}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.31±0.05 OUR AVERAGE</b>				
0.30±0.03±0.07	74 ± 8	HENRARD	87 DM2	$e^+ e^- \rightarrow \Sigma^{*-}$
0.34±0.04±0.07	77 ± 9	HENRARD	87 DM2	$e^+ e^- \rightarrow \Sigma^{*+}$
0.29±0.11±0.10	26	EATON	84 MRK2	$e^+ e^- \rightarrow \Sigma^{*-}$
0.31±0.11±0.11	28	EATON	84 MRK2	$e^+ e^- \rightarrow \Sigma^{*+}$

$\Gamma(\phi f_1(1285))/\Gamma_{\text{total}}$   $\Gamma_{101}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.6±0.5 OUR AVERAGE</b>				
3.4±1.8±1.5	1.1k	<sup>1</sup> ABLIKIM	15H BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \phi \eta \pi^+ \pi^-$
3.2±0.6±0.4		JOUSSET	90 DM2	$J/\psi \rightarrow \phi 2(\pi^+ \pi^-)$
2.1±0.5±0.4	25	<sup>2</sup> JOUSSET	90 DM2	$J/\psi \rightarrow \phi \eta \pi^+ \pi^-$
0.6±0.2±0.1	16	BECKER	87 MRK3	$J/\psi \rightarrow \phi K \bar{K} \pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup> ABLIKIM 15H reports  $[\Gamma(J/\psi(1S) \rightarrow \phi f_1(1285))/\Gamma_{\text{total}}] \times [B(f_1(1285) \rightarrow \eta \pi^+ \pi^-)] = (1.20 \pm 0.6 \pm 0.14) \times 10^{-4}$  which we divide by our best value  $B(f_1(1285) \rightarrow \eta \pi^+ \pi^-) = (35 \pm 15) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> We attribute to the  $f_1(1285)$  the signal observed in the  $\pi^+ \pi^- \eta$  invariant mass distribution at 1297 MeV.

$\Gamma(\phi f_1(1285) \rightarrow \phi \pi^0 f_0(980) \rightarrow \phi \pi^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$   $\Gamma_{102}/\Gamma$

VALUE (units $10^{-7}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>9.36±2.31±1.54</b>	78	ABLIKIM	15P BES3	$J/\psi \rightarrow K^+ K^- 3\pi$

$\Gamma(\phi f_1(1285) \rightarrow \phi \pi^0 f_0(980) \rightarrow \phi \pi^0 \pi^0 \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{103}/\Gamma$

VALUE (units $10^{-7}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.08±1.63±1.47</b>	9	ABLIKIM	15P BES3	$J/\psi \rightarrow K^+ K^- 3\pi$

$\Gamma(\eta \rho)/\Gamma_{\text{total}}$   $\Gamma_{105}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.193±0.023 OUR AVERAGE</b>				
0.194±0.017±0.029	299	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
0.193±0.013±0.029		COFFMAN	88 MRK3	$e^+ e^- \rightarrow \pi^+ \pi^- \eta$

$\Gamma(\omega\eta'(958))/\Gamma_{\text{total}}$   $\Gamma_{106}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.89±0.18 OUR AVERAGE</b>				
2.08±0.30±0.14	137	<sup>1</sup> ABLIKIM	17AK BES3	$J/\psi \rightarrow \pi^+\pi^-\eta'$
2.26±0.43	218	<sup>2</sup> ABLIKIM	06F BES2	$J/\psi \rightarrow \omega\eta'$
1.8 $^{+1.0}_{-0.8}$ ±0.3	6	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
1.66±0.17±0.19		COFFMAN	88 MRK3	$e^+e^- \rightarrow 3\pi\eta'$

<sup>1</sup> From a partial wave analysis of the decay  $J/\psi \rightarrow \pi^+\pi^-\eta'$ .

<sup>2</sup> Using  $B(\eta' \rightarrow \pi^+\pi^-\eta) = (44.3 \pm 1.5)\%$ ,  $B(\eta' \rightarrow \pi^+\pi^-\gamma) = 29.5 \pm 1.0\%$ ,  $B(\eta \rightarrow 2\gamma) = 39.43 \pm 0.26\%$ , and  $B(\omega \rightarrow \pi^+\pi^-\pi^0) = (89.1 \pm 0.7)\%$ .

$\Gamma(\omega f_0(980))/\Gamma_{\text{total}}$   $\Gamma_{107}/\Gamma$

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
<b>1.41±0.27±0.47</b>	<sup>1</sup> AUGUSTIN	89 DM2	$J/\psi \rightarrow 2(\pi^+\pi^-)\pi^0$

<sup>1</sup> Assuming  $B(f_0(980) \rightarrow \pi\pi) = 0.78$ .

$\Gamma(\rho\eta'(958))/\Gamma_{\text{total}}$   $\Gamma_{108}/\Gamma$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>8.1 ±0.8 OUR AVERAGE</b>				Error includes scale factor of 1.6.
7.90±0.19±0.49	3476	<sup>1</sup> ABLIKIM	17AK BES3	$J/\psi \rightarrow \pi^+\pi^-\eta'$
8.3 ±3.0 ±1.2	19	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
11.4 ±1.4 ±1.6		COFFMAN	88 MRK3	$J/\psi \rightarrow \pi^+\pi^-\eta'$

<sup>1</sup> From a partial wave analysis of the decay  $J/\psi \rightarrow \pi^+\pi^-\eta'$ .

$\Gamma(a_2(1320)^\pm \pi^\mp)/\Gamma_{\text{total}}$   $\Gamma_{109}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;43 × 10<sup>-4</sup></b>	90	BRAUNSCH...	76 DASP	$e^+e^-$

$\Gamma(K\bar{K}_2^*(1430) + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{110}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;40 × 10<sup>-4</sup></b>	90	VANNUCCI	77 MRK1	$e^+e^- \rightarrow K^0\bar{K}_2^{*0}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<66 × 10 <sup>-4</sup>	90	BRAUNSCH...	76 DASP	$e^+e^- \rightarrow K^\pm\bar{K}_2^{*\mp}$
------------------------	----	-------------	---------	--

$\Gamma(K_1(1270)^\pm K^\mp)/\Gamma_{\text{total}}$   $\Gamma_{111}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;3.0 × 10<sup>-3</sup></b>	90	<sup>1</sup> BAI	99C BES	$e^+e^-$

<sup>1</sup> Assuming  $B(K_1(1270) \rightarrow K\rho) = 0.42 \pm 0.06$

$\Gamma(K_1(1270)K_S^0 \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$   $\Gamma_{112}/\Gamma$

VALUE (units $10^{-7}$ )	DOCUMENT ID	TECN	COMMENT
<b>8.54 <math>^{+1.07+2.35}_{-1.20-2.13}</math></b>	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

$\Gamma(K_2^*(1430)^0 \bar{K}_2^*(1430)^0)/\Gamma_{\text{total}}$   $\Gamma_{114}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<29 \times 10^{-4}$	90	VANNUCCI 77	MRK1	$e^+e^- \rightarrow \pi^+\pi^-K^+K^-$

$\Gamma(\phi\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{115}/\Gamma$

The two different fit values of ABLIKIM 15K below have the same statistical significance of  $6.4\sigma$  and cannot be distinguished at this moment.

VALUE (units $10^{-6}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$2.94 \pm 0.16 \pm 0.16$		0.8k	<sup>1</sup> ABLIKIM	15K BES3	$e^+e^- \rightarrow J/\psi \rightarrow K^+K^-\gamma\gamma$
$0.124 \pm 0.033 \pm 0.030$		$35 \pm 9$	<sup>2</sup> ABLIKIM	15K BES3	$e^+e^- \rightarrow J/\psi \rightarrow K^+K^-\gamma\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<6.4$	90	<sup>3</sup> ABLIKIM	05B BES2	$e^+e^- \rightarrow J/\psi \rightarrow \phi\gamma\gamma$
$<6.8$	90	COFFMAN	88 MRK3	$e^+e^- \rightarrow K^+K^-\pi^0$

<sup>1</sup> Corresponding to one of the two fit solutions with  $\delta = (-95.9 \pm 1.5)^\circ$  for the phase angle between the resonant  $J/\psi \rightarrow \phi\pi^0$  and non-phi  $J/\psi \rightarrow K^+K^-\pi^0$  contributions.

<sup>2</sup> Corresponding to one of the two fit solutions with  $\delta = (-152.1 \pm 7.7)^\circ$  for the phase angle between the resonant  $J/\psi \rightarrow \phi\pi^0$  and non-phi  $J/\psi \rightarrow K^+K^-\pi^0$  contributions.

<sup>3</sup> Superseded by ABLIKIM 15K.

$\Gamma(\phi\eta(1405) \rightarrow \phi\eta\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{116}/\Gamma$

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$2.01 \pm 0.58 \pm 0.82$		172	<sup>1</sup> ABLIKIM	15H BES3	$e^+e^- \rightarrow J/\psi \rightarrow \phi\eta\pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<17$	90	<sup>2</sup> FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
-------	----	----------------------	--------	-------------------------------------

<sup>1</sup> With  $3.6\sigma$  significance.

<sup>2</sup> Includes unknown branching fraction  $\eta(1405) \rightarrow \eta\pi\pi$ .

$\Gamma(\omega f_2'(1525))/\Gamma_{\text{total}}$   $\Gamma_{117}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.2 \times 10^{-4}$	90	<sup>1</sup> VANNUCCI 77	MRK1	$e^+e^- \rightarrow \pi^+\pi^-\pi^0K^+K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<2.8 \times 10^{-4}$	90	<sup>1</sup> FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
-----------------------	----	----------------------	--------	-------------------------------------

<sup>1</sup> Re-evaluated assuming  $B(f_2'(1525) \rightarrow K\bar{K}) = 0.713$ .

$\Gamma(\omega X(1835) \rightarrow \omega p\bar{p})/\Gamma_{\text{total}}$   $\Gamma_{118}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.9 \times 10^{-6}$	95	ABLIKIM	13P BES3	$J/\psi \rightarrow \gamma\pi^0 p\bar{p}$

$\Gamma(\phi X(1835) \rightarrow \phi p\bar{p})/\Gamma_{\text{total}}$   $\Gamma_{119}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.1 \times 10^{-7}$	90	<sup>1</sup> ABLIKIM	16K BES3	$J/\psi \rightarrow p\bar{p}K_S^0K_L^0, p\bar{p}K^+K^-$

<sup>1</sup> Upper limit applies to any  $p\bar{p}$  mass enhancement near threshold.



$\Gamma(\phi X(1835) \rightarrow \phi \eta \pi^+ \pi^-) / \Gamma_{\text{total}}$					$\Gamma_{120} / \Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<2.8 \times 10^{-4}$	90	ABLIKIM	15H BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \phi \eta \pi^+ \pi^-$	
$\Gamma(\phi X(1870) \rightarrow \phi \eta \pi^+ \pi^-) / \Gamma_{\text{total}}$					$\Gamma_{121} / \Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<6.13 \times 10^{-5}$	90	ABLIKIM	15H BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \phi \eta \pi^+ \pi^-$	
$\Gamma(\eta \phi(2170) \rightarrow \eta \phi f_0(980) \rightarrow \eta \phi \pi^+ \pi^-) / \Gamma_{\text{total}}$					$\Gamma_{122} / \Gamma$
VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
$1.20 \pm 0.14 \pm 0.37$	471	ABLIKIM	15H BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \phi \eta \pi^+ \pi^-$	
$\Gamma(\eta \phi(2170) \rightarrow \eta K^*(892)^0 \bar{K}^*(892)^0) / \Gamma_{\text{total}}$					$\Gamma_{123} / \Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<2.52 \times 10^{-4}$	90	ABLIKIM	10C BES2	$J/\psi \rightarrow \eta K^+ \pi^- K^- \pi^+$	
$\Gamma(\Sigma(1385)^0 \bar{\Lambda} + \text{c.c.}) / \Gamma_{\text{total}}$					$\Gamma_{124} / \Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<0.82 \times 10^{-5}$	90	ABLIKIM	13F BES3	$J/\psi \rightarrow p \bar{p} \pi^+ \pi^- \gamma \gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<0.2 \times 10^{-3}$	90	HENRARD	87 DM2	$e^+ e^-$	
$\Gamma(\Delta(1232)^+ \bar{p}) / \Gamma_{\text{total}}$					$\Gamma_{125} / \Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<0.1 \times 10^{-3}$	90	HENRARD	87 DM2	$e^+ e^-$	
$\Gamma(\Lambda(1520) \bar{\Lambda} + \text{c.c.} \rightarrow \gamma \Lambda \bar{\Lambda}) / \Gamma_{\text{total}}$					$\Gamma_{126} / \Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<4.1 \times 10^{-6}$	90	ABLIKIM	12B BES3	$J/\psi \rightarrow \Lambda \bar{\Lambda} \gamma$	
$\Gamma(\bar{\Lambda}(1520) \Lambda + \text{c.c.}) / \Gamma_{\text{total}}$					$\Gamma_{127} / \Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.80 \times 10^{-3}$	90	LU	19 BELL	$B^+ \rightarrow \bar{p} \Lambda K^+ K^+$	
$\Gamma(\Theta(1540) \bar{\Theta}(1540) \rightarrow K_S^0 p K^- \bar{n} + \text{c.c.}) / \Gamma_{\text{total}}$					$\Gamma_{128} / \Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.1 \times 10^{-5}$	90	BAI	04G BES2	$e^+ e^-$	
$\Gamma(\Theta(1540) K^- \bar{n} \rightarrow K_S^0 p K^- \bar{n}) / \Gamma_{\text{total}}$					$\Gamma_{129} / \Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<2.1 \times 10^{-5}$	90	BAI	04G BES2	$e^+ e^-$	
$\Gamma(\Theta(1540) K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n) / \Gamma_{\text{total}}$					$\Gamma_{130} / \Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.6 \times 10^{-5}$	90	BAI	04G BES2	$e^+ e^-$	

$\Gamma(\bar{\Theta}(1540)K^+n \rightarrow K_S^0\bar{p}K^+n)/\Gamma_{\text{total}}$   $\Gamma_{131}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.6 \times 10^{-5}$	90	BAI	04G	BES2 $e^+e^-$

$\Gamma(\bar{\Theta}(1540)K_S^0p \rightarrow K_S^0pK^-\bar{n})/\Gamma_{\text{total}}$   $\Gamma_{132}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.1 \times 10^{-5}$	90	BAI	04G	BES2 $e^+e^-$

————— STABLE HADRONS —————

$\Gamma(2(\pi^+\pi^-\pi^0))/\Gamma_{\text{total}}$   $\Gamma_{133}/\Gamma$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.37 ± 0.26 OUR AVERAGE</b>				
3.25 ± 0.49	46055	AUGUSTIN	89 DM2	$J/\psi \rightarrow 2(\pi^+\pi^-\pi^0)$
3.17 ± 0.42	147	FRANKLIN	83 MRK2	$e^+e^- \rightarrow \text{hadrons}$
3.64 ± 0.52	1500	BURMESTER	77D PLUT	$e^+e^-$
4 ± 1	675	JEAN-MARIE	76 MRK1	$e^+e^-$

$\Gamma(3(\pi^+\pi^-\pi^0))/\Gamma_{\text{total}}$   $\Gamma_{134}/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.029 ± 0.006 OUR AVERAGE</b>				
0.028 ± 0.009	11	FRANKLIN	83 MRK2	$e^+e^- \rightarrow \text{hadrons}$
0.029 ± 0.007	181	JEAN-MARIE	76 MRK1	$e^+e^-$

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{135}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>21.0 ± 0.8 OUR AVERAGE</b> Error includes scale factor of 1.6. See the ideogram below.				
21.37 ± 0.04 <sup>+0.64</sup> <sub>-0.62</sub>	1.8M	<sup>1,2</sup> ABLIKIM	12H BES3	$e^+e^- \rightarrow J/\psi$
23.0 ± 2.0 ± 0.4	256	<sup>3</sup> AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow J/\psi\pi^+\pi^-\gamma$
21.84 ± 0.05 ± 2.01	220k	<sup>1,4</sup> BAI	04H BES	$e^+e^-$
20.91 ± 0.21 ± 1.16		<sup>4,5</sup> BAI	04H BES	$e^+e^-$
15 ± 2	168	FRANKLIN	83 MRK2	$e^+e^-$


<sup>1</sup> From  $J/\psi \rightarrow \pi^+\pi^-\pi^0$  events directly.

<sup>2</sup> The quoted systematic error includes a contribution of 1.23% (added in quadrature) from the uncertainty on the number of  $J/\psi$  events.

<sup>3</sup> AUBERT 07AU reports  $[\Gamma(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] = (18.6 \pm 1.2 \pm 1.1) \times 10^{-3}$  keV which we divide by our best value  $\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}} = 0.808 \pm 0.013$  keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>4</sup> Mostly  $\rho\pi$ , see also  $\rho\pi$  subsection.

<sup>5</sup> Obtained comparing the rates for  $\pi^+\pi^-\pi^0$  and  $\mu^+\mu^-$ , using  $J/\psi$  events produced via  $\psi(2S) \rightarrow \pi^+\pi^-J/\psi$  and with  $B(J/\psi \rightarrow \mu^+\mu^-) = 5.88 \pm 0.10\%$ .



**$\Gamma(\pi^+ \pi^- \pi^0 K^+ K^-) / \Gamma_{\text{total}}$**   **$\Gamma_{139} / \Gamma$**

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.2±0.3</b>	309	VANNUCCI 77	MRK1	$e^+ e^-$

**$\Gamma(4(\pi^+ \pi^-) \pi^0) / \Gamma_{\text{total}}$**   **$\Gamma_{140} / \Gamma$**

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>90±30</b>	13	JEAN-MARIE 76	MRK1	$e^+ e^-$

**$\Gamma(\pi^+ \pi^- K^+ K^-) / \Gamma_{\text{total}}$**   **$\Gamma_{141} / \Gamma$**

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
• • • 7.2±2.3	205	VANNUCCI 77	MRK1	$e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

**$\Gamma(K \bar{K} \pi) / \Gamma_{\text{total}}$**   **$\Gamma_{149} / \Gamma$**

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>61 ±10 OUR AVERAGE</b>				
55.2±12.0	25	FRANKLIN 83	MRK2	$e^+ e^- \rightarrow K^+ K^- \pi^0$
78.0±21.0	126	VANNUCCI 77	MRK1	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp$

**$\Gamma(K^+ K^- \pi^0) / \Gamma(\pi^+ \pi^- \pi^0)$**   **$\Gamma_{150} / \Gamma_{135}$**

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>12.0±0.3±0.9</b>	23K	LEES	17C BABR	$J/\psi \rightarrow h^+ h^- \pi^0$

$\Gamma(K_S^0 K^\pm \pi^\mp)/\Gamma(\pi^+ \pi^- \pi^0)$					$\Gamma_{151}/\Gamma_{135}$
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>26.5±0.5±2.1</b>	24K	LEES	17C	BABR	$J/\psi \rightarrow h^0 h^+ h^-$

$\Gamma(2(\pi^+ \pi^-))/\Gamma_{\text{total}}$					$\Gamma_{156}/\Gamma$
VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>3.57±0.30 OUR AVERAGE</b>					
3.53±0.12±0.29	1107	<sup>1</sup> ABLIKIM	05H	BES2	$e^+ e^- \rightarrow \psi(2S) \rightarrow J/\psi \pi^+ \pi^-, J/\psi \rightarrow 2(\pi^+ \pi^-)$
4.0 ±1.0	76	JEAN-MARIE	76	MRK1	$e^+ e^-$

<sup>1</sup> Computed using  $B(J/\psi \rightarrow \mu^+ \mu^-) = 0.0588 \pm 0.0010$ .

$\Gamma(3(\pi^+ \pi^-))/\Gamma_{\text{total}}$					$\Gamma_{157}/\Gamma$
VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
40±20	32	JEAN-MARIE	76	MRK1	$e^+ e^-$

$\Gamma(2(\pi^+ \pi^-)\eta)/\Gamma_{\text{total}}$					$\Gamma_{159}/\Gamma$
VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>2.26±0.08±0.27</b>	4.8k	ABLIKIM	05C	BES2	$e^+ e^- \rightarrow 2(\pi^+ \pi^-)\eta$

$\Gamma(3(\pi^+ \pi^-)\eta)/\Gamma_{\text{total}}$					$\Gamma_{160}/\Gamma$
VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>7.24±0.96±1.11</b>	616	ABLIKIM	05C	BES2	$e^+ e^- \rightarrow 3(\pi^+ \pi^-)\eta$

$\Gamma(\rho\bar{\rho})/\Gamma_{\text{total}}$					$\Gamma_{163}/\Gamma$
VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>2.121±0.029 OUR AVERAGE</b>					
2.112±0.004±0.031	314k	ABLIKIM	12C	BES3	$e^+ e^-$
2.19 ±0.16 ±0.06	317	<sup>1</sup> WU	06	BELL	$B^+ \rightarrow \rho\bar{\rho}K^+$
2.26 ±0.01 ±0.14	63316	BAI	04E	BES2	$e^+ e^- \rightarrow J/\psi$
1.97 ±0.22	99	BALDINI	98	FENI	$e^+ e^-$
1.91 ±0.04 ±0.30		PALLIN	87	DM2	$e^+ e^-$
2.16 ±0.07 ±0.15	1420	EATON	84	MRK2	$e^+ e^-$
2.5 ±0.4	133	BRANDELIK	79C	DASP	$e^+ e^-$
2.0 ±0.5		BESCH	78	BONA	$e^+ e^-$
2.2 ±0.2	331	<sup>2</sup> PERUZZI	78	MRK1	$e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.0 ±0.3      48      ANTONELLI    93    SPEC     $e^+ e^-$

<sup>1</sup> WU 06 reports  $[\Gamma(J/\psi(1S) \rightarrow \rho\bar{\rho})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow J/\psi(1S)K^+)] = (2.21 \pm 0.13 \pm 0.10) \times 10^{-6}$  which we divide by our best value  $B(B^+ \rightarrow J/\psi(1S)K^+) = (1.010 \pm 0.028) \times 10^{-3}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Assuming angular distribution  $(1+\cos^2\theta)$ .


$\Gamma(\rho\bar{p}\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{164}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.19±0.08 OUR AVERAGE</b>		Error includes scale factor of 1.1.		
1.33±0.02±0.11	11k	ABLIKIM	09B	BES2 $e^+e^-$
1.13±0.09±0.09	685	EATON	84	MRK2 $e^+e^-$
1.4 ±0.4		BRANDELIK	79C	DASP $e^+e^-$
1.00±0.15	109	PERUZZI	78	MRK1 $e^+e^-$

$\Gamma(\rho\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{165}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>6.0 ±0.5 OUR AVERAGE</b>		Error includes scale factor of 1.3. See the ideogram below.		
6.46±0.17±0.43	1435	EATON	84	MRK2 $e^+e^-$
3.8 ±1.6	48	BESCH	81	BONA $e^+e^-$
5.5 ±0.6	533	PERUZZI	78	MRK1 $e^+e^-$

WEIGHTED AVERAGE  
6.0±0.5 (Error scaled by 1.3)



$\Gamma(\rho\bar{p}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{166}/\Gamma$

Including  $\rho\bar{p}\pi^+\pi^-\gamma$  and excluding  $\omega, \eta, \eta'$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.3 ±0.9 OUR AVERAGE</b>		Error includes scale factor of 1.9.		
3.36±0.65±0.28	364	EATON	84	MRK2 $e^+e^-$
1.6 ±0.6	39	PERUZZI	78	MRK1 $e^+e^-$

$\Gamma(p\bar{p}\eta)/\Gamma_{\text{total}}$   $\Gamma_{167}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.00 ± 0.12 OUR AVERAGE</b>				
1.91 ± 0.02 ± 0.17	13k	<sup>1</sup> ABLIKIM	09	BES2 $e^+e^-$
2.03 ± 0.13 ± 0.15	826	EATON	84	MRK2 $e^+e^-$
2.5 ± 1.2		BRANDELIK	79c	DASP $e^+e^-$
2.3 ± 0.4	197	PERUZZI	78	MRK1 $e^+e^-$


<sup>1</sup> From the combination of  $p\bar{p}\eta \rightarrow p\bar{p}\gamma\gamma$  and  $p\bar{p}\eta \rightarrow p\bar{p}\pi^+\pi^-\pi^0$  channels.

$\Gamma(p\bar{p}\rho)/\Gamma_{\text{total}}$   $\Gamma_{168}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 0.31 × 10<sup>-3</sup></b>	90	EATON	84	MRK2 $e^+e^- \rightarrow \text{hadrons}\gamma$

$\Gamma(p\bar{p}\omega)/\Gamma_{\text{total}}$   $\Gamma_{169}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.98 ± 0.10 OUR AVERAGE</b> Error includes scale factor of 1.3. See the ideogram below.				
0.90 ± 0.02 ± 0.09	2670	ABLIKIM	13P	BES3 $e^+e^-$
0.98 ± 0.03 ± 0.14	2449	ABLIKIM	08	BES2 $e^+e^-$
1.10 ± 0.17 ± 0.18	486	EATON	84	MRK2 $e^+e^-$
1.6 ± 0.3	77	PERUZZI	78	MRK1 $e^+e^-$



$\Gamma(p\bar{p}\eta'(958))/\Gamma_{\text{total}}$   $\Gamma_{170}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.129±0.014 OUR AVERAGE</b> Error includes scale factor of 2.0.				
0.126±0.002±0.007	16K	<sup>1</sup> ABLIKIM	19N BES3	$e^+e^-$
0.200±0.023±0.028	265 ± 31	<sup>2</sup> ABLIKIM	09 BES2	$e^+e^-$
0.68 ±0.23 ±0.17	19	EATON	84 MRK2	$e^+e^-$
1.8 ±0.6	19	PERUZZI	78 MRK1	$e^+e^-$

<sup>1</sup> From the combination of  $p\bar{p}\eta' \rightarrow p\bar{p}\pi^+\pi^-\eta$  and  $p\bar{p}\eta' \rightarrow p\bar{p}\pi^+\pi^-\gamma$  channels.  
<sup>2</sup> From the combination of  $p\bar{p}\eta' \rightarrow p\bar{p}\pi^+\pi^-\eta$  and  $p\bar{p}\eta' \rightarrow p\bar{p}\gamma\rho^0$  channels.

$\Gamma(p\bar{p}a_0(980) \rightarrow p\bar{p}\pi^0\eta)/\Gamma_{\text{total}}$   $\Gamma_{171}/\Gamma$

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
<b>6.8±1.2±1.3</b>	ABLIKIM	14N BES3	$e^+e^- \rightarrow J/\psi$

$\Gamma(p\bar{p}\phi)/\Gamma_{\text{total}}$   $\Gamma_{172}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.519±0.033 OUR AVERAGE</b>				
0.523±0.006±0.033	14K	ABLIKIM	16K BES3	$J/\psi \rightarrow p\bar{p}K_S^0 K_L^0,$ $p\bar{p}K^+ K^-$
0.45 ±0.13 ±0.07		FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$

$\Gamma(n\bar{n})/\Gamma_{\text{total}}$   $\Gamma_{173}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.09±0.16 OUR AVERAGE</b>				
2.07±0.01±0.17	36k	ABLIKIM	12C BES3	$e^+e^-$
2.31±0.49	79	BALDINI	98 FENI	$e^+e^-$
1.8 ±0.9		BESCH	78 BONA	$e^+e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.90±0.55	40	ANTONELLI	93 SPEC	$e^+e^-$

$\Gamma(n\bar{n}\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{174}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.8±3.6</b>	5	BESCH	81 BONA	$e^+e^-$

$\Gamma(\Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}$   $\Gamma_{175}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.50±0.10±0.22</b>	399	ABLIKIM	08O BES2	$e^+e^- \rightarrow J/\psi$

$\Gamma(\Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}$   $\Gamma_{176}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.172±0.032 OUR AVERAGE</b> Error includes scale factor of 1.4.				
1.164±0.004±0.023	111k	ABLIKIM	17L BES3	$J/\psi \rightarrow \Sigma^0\bar{\Sigma}^0$
1.33 ±0.04 ±0.11	1.7k	ABLIKIM	06 BES2	$J/\psi \rightarrow \Sigma^0\bar{\Sigma}^0$
1.06 ±0.04 ±0.23	884	PALLIN	87 DM2	$e^+e^- \rightarrow \Sigma^0\bar{\Sigma}^0$
1.58 ±0.16 ±0.25	90	EATON	84 MRK2	$e^+e^- \rightarrow \Sigma^0\bar{\Sigma}^0$
1.3 ±0.4	52	PERUZZI	78 MRK1	$e^+e^- \rightarrow \Sigma^0\bar{\Sigma}^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.4 ±2.6	3	BESCH	81 BONA	$e^+e^- \rightarrow \Sigma^+\bar{\Sigma}^-$

$\Gamma(2(\pi^+\pi^-)K^+K^-)/\Gamma_{\text{total}}$   $\Gamma_{177}/\Gamma$


VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>31±13</b>	30	VANNUCCI 77	MRK1	$e^+e^-$

$\Gamma(\rho\bar{n}\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{178}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.12±0.09 OUR AVERAGE</b>				
2.36±0.02±0.21	59k	ABLIKIM 06K	BES2	$J/\psi \rightarrow \rho\pi^-\bar{n}$
2.47±0.02±0.24	55k	ABLIKIM 06K	BES2	$J/\psi \rightarrow \bar{\rho}\pi^+n$
2.02±0.07±0.16	1288	EATON 84	MRK2	$e^+e^- \rightarrow \rho\pi^-$
1.93±0.07±0.16	1191	EATON 84	MRK2	$e^+e^- \rightarrow \bar{\rho}\pi^+$
1.7 ±0.7	32	BESCH 81	BONA	$e^+e^- \rightarrow \rho\pi^-$
1.6 ±1.2	5	BESCH 81	BONA	$e^+e^- \rightarrow \bar{\rho}\pi^+$
2.16±0.29	194	PERUZZI 78	MRK1	$e^+e^- \rightarrow \rho\pi^-$
2.04±0.27	204	PERUZZI 78	MRK1	$e^+e^- \rightarrow \bar{\rho}\pi^+$

$\Gamma(\Xi^-\Xi^+)/\Gamma_{\text{total}}$   $\Gamma_{182}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.97 ±0.08 OUR AVERAGE</b> Error includes scale factor of 1.4. See the ideogram below.				
1.040±0.006±0.074	43k	ABLIKIM 16L	BES3	$J/\psi \rightarrow \Xi^-\Xi^+$
0.90 ±0.03 ±0.18	961	ABLIKIM 12P	BES2	$J/\psi \rightarrow \Xi^-\Xi^+$
0.70 ±0.06 ±0.12	132	HENRARD 87	DM2	$e^+e^- \rightarrow \Xi^-\Xi^+$
1.14 ±0.08 ±0.20	194	EATON 84	MRK2	$e^+e^- \rightarrow \Xi^-\Xi^+$
1.4 ±0.5	51	PERUZZI 78	MRK1	$e^+e^- \rightarrow \Xi^-\Xi^+$






**$\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$**   **$\Gamma_{183}/\Gamma$**

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.89 ± 0.09 OUR AVERAGE</b>		Error includes scale factor of 2.8.		See the ideogram below.
1.943 ± 0.003 ± 0.033	441k	ABLIKIM	17L BES3	$e^+e^-$
2.03 ± 0.03 ± 0.15	8887	ABLIKIM	06 BES2	$J/\psi \rightarrow \Lambda\bar{\Lambda}$
2.0 $^{+0.5}_{-0.4}$ ± 0.1	46	<sup>1</sup> WU	06 BELL	$B^+ \rightarrow \Lambda\bar{\Lambda}K^+$
1.08 ± 0.06 ± 0.24	631	BAI	98G BES	$e^+e^-$
1.38 ± 0.05 ± 0.20	1847	PALLIN	87 DM2	$e^+e^-$
1.58 ± 0.08 ± 0.19	365	EATON	84 MRK2	$e^+e^-$
2.6 ± 1.6	5	BESCH	81 BONA	$e^+e^-$
1.1 ± 0.2	196	PERUZZI	78 MRK1	$e^+e^-$

<sup>1</sup>WU 06 reports  $[\Gamma(J/\psi(1S) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow J/\psi(1S)K^+)] = (2.00^{+0.34}_{-0.29} \pm 0.34) \times 10^{-6}$  which we divide by our best value  $B(B^+ \rightarrow J/\psi(1S)K^+) = (1.010 \pm 0.028) \times 10^{-3}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

WEIGHTED AVERAGE  
1.89±0.09 (Error scaled by 2.8)



Values above of weighted average, error, and scale factor are based upon the data in this ideogram only. They are not necessarily the same as our 'best' values, obtained from a least-squares constrained fit utilizing measurements of other (related) quantities as additional information.

**$\Gamma(\Lambda\bar{\Sigma}^-\pi^+ \text{ (or c.c.)})/\Gamma_{\text{total}}$**   **$\Gamma_{184}/\Gamma$**

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.83 ± 0.07 OUR AVERAGE</b>		Error includes scale factor of 1.2.		
0.770 ± 0.051 ± 0.083	335	<sup>1</sup> ABLIKIM	07H BES2	$e^+e^- \rightarrow \bar{\Lambda}\Sigma^+\pi^-$
0.747 ± 0.056 ± 0.076	254	<sup>1</sup> ABLIKIM	07H BES2	$e^+e^- \rightarrow \Lambda\bar{\Sigma}^-\pi^+$
0.90 ± 0.06 ± 0.16	225 ± 15	HENRARD	87 DM2	$e^+e^- \rightarrow \bar{\Lambda}\Sigma^+\pi^-$

1.11 ±0.06 ±0.20	342 ± 18	HENRARD	87	DM2	$e^+e^- \rightarrow \Lambda \bar{\Sigma}^- \pi^+$
1.53 ±0.17 ±0.38	135	EATON	84	MRK2	$e^+e^- \rightarrow \bar{\Lambda} \Sigma^+ \pi^-$
1.38 ±0.21 ±0.35	118	EATON	84	MRK2	$e^+e^- \rightarrow \Lambda \bar{\Sigma}^- \pi^+$

<sup>1</sup> Using  $B(\Lambda \rightarrow \pi^- p) = 63.9\%$  and  $B(\Sigma^+ \rightarrow \pi^0 p) = 51.6\%$ .

### $\Gamma(pK^-\bar{\Lambda}+c.c.)/\Gamma_{\text{total}}$ $\Gamma_{185}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.87±0.11 OUR AVERAGE</b>				
0.85 <sup>+0.17</sup> <sub>-0.15</sub> ±0.02	45	<sup>1</sup> LU	19	BELL $B^+ \rightarrow \bar{p} \Lambda K^+ K^+$
0.89±0.07±0.14	307	EATON	84	MRK2 $e^+e^-$

<sup>1</sup> LU 19 reports  $(8.32^{+1.63}_{-1.45} \pm 0.49) \times 10^{-4}$  from a measurement of  $[\Gamma(J/\psi(1S) \rightarrow pK^-\bar{\Lambda}+c.c.)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow J/\psi(1S)K^+)]$  assuming  $B(B^+ \rightarrow J/\psi(1S)K^+) = (1.026 \pm 0.031) \times 10^{-3}$ , which we rescale to our best value  $B(B^+ \rightarrow J/\psi(1S)K^+) = (1.010 \pm 0.028) \times 10^{-3}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

### $\Gamma(2(K^+K^-))/\Gamma_{\text{total}}$ $\Gamma_{186}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.4 <sup>+0.5</sup> <sub>-0.4</sub> ±0.2	11.0 <sup>+4.3</sup> <sub>-3.5</sub>	<sup>1</sup> HUANG	03	BELL $B^+ \rightarrow 2(K^+K^-)K^+$
0.7±0.3		VANNUCCI	77	MRK1 $e^+e^-$

<sup>1</sup> Using  $B(B^+ \rightarrow J/\psi K^+) = (1.01 \pm 0.05) \times 10^{-3}$ .

### $\Gamma(pK^-\bar{\Sigma}^0)/\Gamma_{\text{total}}$ $\Gamma_{187}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.29±0.06±0.05</b>	90	EATON	84	MRK2 $e^+e^-$

### $\Gamma(K^+K^-)/\Gamma_{\text{total}}$ $\Gamma_{188}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.86±0.09±0.19</b>	1k	<sup>1</sup> METREVELI	12	$\psi(2S) \rightarrow \pi^+\pi^-K^+K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.39±0.24±0.22	107	<sup>2</sup> BALTRUSAIT..85D	MRK3	$e^+e^-$
2.2 ±0.9	6	<sup>2</sup> BRANDELIK	79C	DASP $e^+e^-$

<sup>1</sup> Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.


<sup>2</sup> Interference with non-resonant  $K^+K^-$  production not taken into account.

### $\Gamma(K_S^0 K_L^0)/\Gamma_{\text{total}}$ $\Gamma_{189}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.95±0.11 OUR AVERAGE</b> Error includes scale factor of 2.4. See the ideogram below.				
1.93±0.01±0.05	110K	ABLIKIM	17AH	BES3 $J/\psi \rightarrow K_S^0 K_L^0 \rightarrow \pi^+\pi^-X$
2.62±0.15±0.14	0.3k	<sup>1</sup> METREVELI	12	$\psi(2S) \rightarrow \pi^+\pi^-K_S^0 K_L^0$
1.82±0.04±0.13	2.1k	<sup>2</sup> BAI	04A	BES2 $J/\psi \rightarrow K_S^0 K_L^0 \rightarrow \pi^+\pi^-X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.18±0.12±0.18		JOUSSET	90	DM2 $J/\psi \rightarrow$ hadrons
1.01±0.16±0.09	74	BALTRUSAIT..85D	MRK3	$e^+e^-$

<sup>1</sup> Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

<sup>2</sup> Using  $B(K_S^0 \rightarrow \pi^+ \pi^-) = 0.6868 \pm 0.0027$ .



**$\Gamma(\Lambda \bar{\Lambda} \pi^+ \pi^-)/\Gamma_{\text{total}}$   $\Gamma_{190}/\Gamma$**

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>4.30±0.13±0.99</b>	2.4k	ABLIKIM	12P	BES2	$J/\psi$

**$\Gamma(\Lambda \bar{\Lambda} \eta)/\Gamma_{\text{total}}$   $\Gamma_{191}/\Gamma$**

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>16.2±1.7 OUR AVERAGE</b>					
15.7±0.80±1.54	454	<sup>1</sup> ABLIKIM	13F	BES3	$J/\psi \rightarrow p \bar{p} \pi^+ \pi^- \gamma \gamma$
26.2±6.0 ±4.4	44	<sup>2</sup> ABLIKIM	07H	BES2	$e^+ e^- \rightarrow \psi(2S)$

<sup>1</sup> Using  $B(\Lambda \rightarrow \pi^- p) = 63.9\%$  and  $B(\eta \rightarrow \gamma \gamma) = 39.31\%$ .

<sup>2</sup> Using  $B(\Lambda \rightarrow \pi^- p) = 63.9\%$  and  $B(\eta \rightarrow \gamma \gamma) = 39.4\%$ .

**$\Gamma(\Lambda \bar{\Lambda} \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{192}/\Gamma$**

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>3.78±0.27±0.30</b>		323	<sup>1</sup> ABLIKIM	13F	BES3	$J/\psi \rightarrow p \bar{p} \pi^+ \pi^- \gamma \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 6.4	90	<sup>2</sup> ABLIKIM	07H	BES2	$e^+ e^- \rightarrow \psi(2S)$
23 ±7 ±8	11	BAI	98G	BES	$e^+ e^-$
22 ±5 ±5	19	HENRARD	87	DM2	$e^+ e^-$

<sup>1</sup> Using  $B(\Lambda \rightarrow \pi^- p) = 63.9\%$  and  $B(\pi^0 \rightarrow \gamma \gamma) = 98.8\%$ .

<sup>2</sup> Using  $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ .

$\Gamma(\bar{\Lambda}nK_S^0 + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{193}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>6.46 \pm 0.20 \pm 1.07</math></b>	1058	<sup>1</sup> ABLIKIM	08C BES2	$e^+e^- \rightarrow J/\psi$

<sup>1</sup> Using  $B(\bar{\Lambda} \rightarrow \bar{p}\pi^+) = 63.9\%$  and  $B(K_S^0 \rightarrow \pi^+\pi^-) = 69.2\%$ .

$\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{194}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>1.47 \pm 0.14</math> OUR AVERAGE</b>				
$1.47 \pm 0.13 \pm 0.13$	140	<sup>1</sup> METREVELI	12	$\psi(2S) \rightarrow 2(\pi^+\pi^-)$
$1.58 \pm 0.20 \pm 0.15$	84	BALTRUSAIT..85D	MRK3	$e^+e^-$
$1.0 \pm 0.5$	5	BRANDELIK	78B DASP	$e^+e^-$
$1.6 \pm 1.6$	1	VANNUCCI	77 MRK1	$e^+e^-$

<sup>1</sup> Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\Lambda\bar{\Sigma} + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{195}/\Gamma$

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>2.83 \pm 0.23</math> OUR AVERAGE</b>					
$2.74 \pm 0.24 \pm 0.22$		$234 \pm 21$	<sup>1</sup> ABLIKIM	12B BES3	$J/\psi \rightarrow \Lambda\bar{\Sigma}^0$
$2.92 \pm 0.22 \pm 0.24$		$308 \pm 24$	<sup>2</sup> ABLIKIM	12B BES3	$J/\psi \rightarrow \bar{\Lambda}\Sigma^0$
$<18$			<sup>2</sup> HENRARD	87 DM2	$J/\psi \rightarrow \bar{\Lambda}\Sigma^0$
$<15$	90		PERUZZI	78 MRK1	$e^+e^- \rightarrow \Lambda X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup> ABLIKIM 12B quotes  $B(J/\psi \rightarrow \Lambda\bar{\Sigma}^0)$  which we multiply by 2.

<sup>2</sup> ABLIKIM 12B and HENRARD 87 quote results for  $B(J/\psi \rightarrow \bar{\Lambda}\Sigma^0)$  which we multiply by 2.

$\Gamma(K_S^0 K_S^0)/\Gamma_{\text{total}}$   $\Gamma_{196}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>&lt;1.4 \times 10^{-8}</math></b>	95	<sup>1</sup> ABLIKIM	17AH BES3	$J/\psi \rightarrow K_S^0 K_S^0 \rightarrow \pi^+\pi^-\pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1 \times 10^{-6}$	95	<sup>1</sup> BAI	04D BES	$e^+e^-$
$<5.2 \times 10^{-6}$	90	<sup>1</sup> BALTRUSAIT..85C	MRK3	$e^+e^-$

<sup>1</sup> Forbidden by CP.

————— RADIATIVE DECAYS —————

$\Gamma(3\gamma)/\Gamma_{\text{total}}$   $\Gamma_{197}/\Gamma$

VALUE (units $10^{-6}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>11.6 \pm 2.2</math> OUR AVERAGE</b>					
$11.3 \pm 1.8 \pm 2.0$		$113 \pm 18$	ABLIKIM	13I BES3	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$
$12 \pm 3 \pm 2$		$24.2^{+7.2}_{-6.0}$	ADAMS	08 CLEO	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<55$	90	PARTRIDGE	80 CBAL	$e^+e^-$
-------	----	-----------	---------	----------

$\Gamma(4\gamma)/\Gamma_{\text{total}}$   $\Gamma_{198}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>&lt;9 \times 10^{-6}</math></b>	90	ADAMS	08 CLEO	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$

$\Gamma(5\gamma)/\Gamma_{\text{total}}$					$\Gamma_{199}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<15 \times 10^{-6}$	90	ADAMS	08	CLEO	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$

$\Gamma(\gamma\pi^0\pi^0)/\Gamma_{\text{total}}$					$\Gamma_{200}/\Gamma$
VALUE (units $10^{-3}$ )		DOCUMENT ID	TECN	COMMENT	
$1.15 \pm 0.05$		<sup>1</sup> ABLIKIM	15AE	BES3	$J/\psi \rightarrow \gamma\pi^0\pi^0$

<sup>1</sup> The uncertainty is systematic as statistical is negligible.

$\Gamma(\gamma\eta\pi^0)/\Gamma_{\text{total}}$					$\Gamma_{201}/\Gamma$
VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
$21.4 \pm 1.8 \pm 2.5$	596	ABLIKIM	16P	BES3	$J/\psi \rightarrow 5\gamma$

$\Gamma(\gamma a_0(980)^0 \rightarrow \gamma\eta\pi^0)/\Gamma_{\text{total}}$					$\Gamma_{202}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<2.5 \times 10^{-6}$	95	ABLIKIM	16P	BES3	$J/\psi \rightarrow 5\gamma$

$\Gamma(\gamma a_2(1320)^0 \rightarrow \gamma\eta\pi^0)/\Gamma_{\text{total}}$					$\Gamma_{203}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<6.6 \times 10^{-6}$	95	ABLIKIM	16P	BES3	$J/\psi \rightarrow 5\gamma$

$\Gamma(\gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$					$\Gamma_{204}/\Gamma$
VALUE (units $10^{-4}$ )		DOCUMENT ID	TECN	COMMENT	
$8.1 \pm 0.4$		ABLIKIM	18AA	BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

$\Gamma(\gamma\eta_c(1S))/\Gamma_{\text{total}}$					$\Gamma_{205}/\Gamma$
VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
$1.7 \pm 0.4$		<b>OUR AVERAGE</b>			Error includes scale factor of 1.5.

$2.00 \pm 0.31 \pm 0.02$		<sup>1</sup> MITCHELL	09	CLEO	$e^+ e^- \rightarrow \gamma X$
$1.27 \pm 0.36$		GAISER	86	CBAL	$J/\psi \rightarrow \gamma X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen		ANASHIN	14	KEDR	$J/\psi \rightarrow \gamma\eta_c$
$0.79 \pm 0.20$	$273 \pm 43$	<sup>2</sup> AUBERT	06E	BABR	$B^\pm \rightarrow K^\pm X_{c\bar{c}}$
seen	16	BALTRUSAITIS	84	MRK3	$J/\psi \rightarrow 2\phi\gamma$

<sup>1</sup> MITCHELL 09 reports  $(1.98 \pm 0.09 \pm 0.30) \times 10^{-2}$  from a measurement of  $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)]$  assuming  $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (35.04 \pm 0.07 \pm 0.77) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.68 \pm 0.30) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Calculated by the authors using an average of  $B(J/\psi \rightarrow \gamma\eta_c) \times B(\eta_c \rightarrow K\bar{K}\pi)$  from BALTRUSAITIS 86, BISELLO 91, BAI 04 and  $B(\eta_c \rightarrow K\bar{K}\pi) = (8.5 \pm 1.8)\%$  from AUBERT 06E.

$\Gamma(\gamma\eta_c(1S) \rightarrow 3\gamma)/\Gamma_{\text{total}}$   $\Gamma_{206}/\Gamma$

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**$3.8^{+1.3}_{-1.0}$  OUR AVERAGE** Error includes scale factor of 1.1.

$4.5 \pm 1.2 \pm 0.6$	$33 \pm 9$	ABLIKIM	13I	BES3 $\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$
$1.2^{+2.7}_{-1.1} \pm 0.3$	$1.2^{+2.8}_{-1.1}$	ADAMS	08	CLEO $\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$

$\Gamma(\gamma\pi^+ \pi^- 2\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{207}/\Gamma$

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

**$8.3 \pm 0.2 \pm 3.1$**  <sup>1</sup> BALTRUSAIT ..86B MRK3  $J/\psi \rightarrow 4\pi\gamma$

<sup>1</sup>  $4\pi$  mass less than 2.0 GeV.

$\Gamma(\gamma\eta\pi\pi)/\Gamma_{\text{total}}$   $\Gamma_{208}/\Gamma$

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

**$6.1 \pm 1.0$  OUR AVERAGE**

$5.85 \pm 0.3 \pm 1.05$	<sup>1</sup> EDWARDS	83B	CBAL $J/\psi \rightarrow \eta\pi^+\pi^-$
$7.8 \pm 1.2 \pm 2.4$	<sup>1</sup> EDWARDS	83B	CBAL $J/\psi \rightarrow \eta 2\pi^0$

<sup>1</sup> Broad enhancement at 1700 MeV.

$\Gamma(\gamma\eta_2(1870) \rightarrow \gamma\eta\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{209}/\Gamma$

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

**$6.2 \pm 2.2 \pm 0.9$**  BAI 99 BES  $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$

$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma K \bar{K} \pi)/\Gamma_{\text{total}}$   $\Gamma_{210}/\Gamma$

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

**$2.8 \pm 0.6$  OUR AVERAGE** Error includes scale factor of 1.6. See the ideogram below.

$1.66 \pm 0.1 \pm 0.58$	<sup>1,2</sup> BAI	00D	BES $J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$
$3.8 \pm 0.3 \pm 0.6$	<sup>3</sup> AUGUSTIN	90	DM2 $J/\psi \rightarrow \gamma K \bar{K} \pi$
$4.0 \pm 0.7 \pm 1.0$	<sup>3</sup> EDWARDS	82E	CBAL $J/\psi \rightarrow K^+ K^- \pi^0 \gamma$
$4.3 \pm 1.7$	<sup>3,4</sup> SCHARRE	80	MRK2 $e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.78 \pm 0.21 \pm 0.33$	<sup>3,5,6</sup> AUGUSTIN	92	DM2 $J/\psi \rightarrow \gamma K \bar{K} \pi$
$0.83 \pm 0.13 \pm 0.18$	<sup>3,7,8</sup> AUGUSTIN	92	DM2 $J/\psi \rightarrow \gamma K \bar{K} \pi$
$0.66^{+0.17+0.24}_{-0.16-0.15}$	<sup>3,6,9</sup> BAI	90C	MRK3 $J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
$1.03^{+0.21+0.26}_{-0.18-0.19}$	<sup>3,8,10</sup> BAI	90C	MRK3 $J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$

<sup>1</sup> Interference with the  $J/\psi(1S)$  radiative transition to the broad  $K \bar{K} \pi$  pseudoscalar state around 1800 is  $(0.15 \pm 0.01 \pm 0.05) \times 10^{-3}$ .

<sup>2</sup> Interference with  $J/\psi \rightarrow \gamma f_1(1420)$  is  $(-0.03 \pm 0.01 \pm 0.01) \times 10^{-3}$ .

<sup>3</sup> Includes unknown branching fraction  $\eta(1405) \rightarrow K \bar{K} \pi$ .

<sup>4</sup> Corrected for spin-zero hypothesis for  $\eta(1405)$ .

<sup>5</sup> From fit to the  $a_0(980)\pi 0^-+$  partial wave.


<sup>6</sup>  $a_0(980)\pi$  mode.

<sup>7</sup> From fit to the  $K^*(892)K 0^-+$  partial wave.

<sup>8</sup>  $K^* K$  mode.

<sup>9</sup> From  $a_0(980)\pi$  final state.

<sup>10</sup> From  $K^*(890)K$  final state.



**$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\gamma\rho^0) / \Gamma_{\text{total}}$   $\Gamma_{211}/\Gamma$**

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
<b><math>0.78 \pm 0.20</math> OUR AVERAGE</b>	Error includes scale factor of 1.8.		
$1.07 \pm 0.17 \pm 0.11$	<sup>1</sup> BAI	04J BES2	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
$0.64 \pm 0.12 \pm 0.07$	<sup>1</sup> COFFMAN	90 MRK3	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$

<sup>1</sup>Includes unknown branching fraction  $\eta(1405) \rightarrow \gamma\rho^0$ .

**$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\eta\pi^+\pi^-) / \Gamma_{\text{total}}$   $\Gamma_{212}/\Gamma$**

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>3.0 \pm 0.5</math> OUR AVERAGE</b>				
$2.6 \pm 0.7 \pm 0.4$		BAI	99 BES	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
$3.38 \pm 0.33 \pm 0.64$		<sup>1</sup> BOLTON	92B MRK3	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$7.0 \pm 0.6 \pm 1.1$	261	<sup>2</sup> AUGUSTIN	90 DM2	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$

<sup>1</sup> Via  $a_0(980)\pi$ .  
<sup>2</sup> Includes unknown branching fraction to  $\eta\pi^+\pi^-$ .

**$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\gamma\phi) / \Gamma_{\text{total}}$   $\Gamma_{213}/\Gamma$**

VALUE (units $10^{-6}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>&lt;82</b>	95		BAI	04J BES2	$J/\psi \rightarrow \gamma\gamma K^+ K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$7.03 \pm 0.92 \pm 0.91$		1.3k	<sup>1</sup> ABLIKIM	18i BES3	$J/\psi \rightarrow \gamma\gamma\phi(1020)$
$10.36 \pm 1.51 \pm 1.54$		1.9k	<sup>2</sup> ABLIKIM	18i BES3	$J/\psi \rightarrow \gamma\gamma\phi(1020)$

<sup>1</sup> Constructive interference between the  $X(1835)$  and  $\eta(1405)/\eta(1475)$  is assumed in a fit to the  $\gamma\phi$  invariant mass.

<sup>2</sup> Destructive interference between the  $X(1835)$  and  $\eta(1405)/\eta(1475)$  is assumed in a fit to the  $\gamma\phi$  invariant mass.

**$\Gamma(\gamma\eta(1405) \rightarrow \gamma\gamma\gamma)/\Gamma_{\text{total}}$   $\Gamma_{214}/\Gamma$**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.63 \times 10^{-6}$	90	ABLIKIM	180	BES3 $\psi(2S) \rightarrow \pi^+\pi^-\gamma\gamma\gamma$

**$\Gamma(\gamma\eta(1475) \rightarrow \gamma\gamma\gamma)/\Gamma_{\text{total}}$   $\Gamma_{215}/\Gamma$**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.86 \times 10^{-6}$	90	ABLIKIM	180	BES3 $\psi(2S) \rightarrow \pi^+\pi^-\gamma\gamma\gamma$

**$\Gamma(\gamma\rho\rho)/\Gamma_{\text{total}}$   $\Gamma_{216}/\Gamma$**

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>4.5 \pm 0.8</math> OUR AVERAGE</b>				
$4.7 \pm 0.3 \pm 0.9$		<sup>1</sup> BALTRUSAIT..86B	MRK3	$J/\psi \rightarrow 4\pi\gamma$
$3.75 \pm 1.05 \pm 1.20$		<sup>2</sup> BURKE	82	MRK2 $J/\psi \rightarrow 4\pi\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<0.09$	90	<sup>3</sup> BISELLO	89B	$J/\psi \rightarrow 4\pi\gamma$
---------	----	----------------------	-----	---------------------------------

<sup>1</sup>  $4\pi$  mass less than 2.0 GeV.  
<sup>2</sup>  $4\pi$  mass less than 2.0 GeV. We have multiplied  $2\rho^0$  measurement by 3 to obtain  $2\rho$ .  
<sup>3</sup>  $4\pi$  mass in the range 2.0–25 GeV.

**$\Gamma(\gamma\rho\omega)/\Gamma_{\text{total}}$   $\Gamma_{217}/\Gamma$**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.4 \times 10^{-4}$	90	ABLIKIM	08A	BES2 $e^+e^- \rightarrow J/\psi$

**$\Gamma(\gamma\rho\phi)/\Gamma_{\text{total}}$   $\Gamma_{218}/\Gamma$**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<8.8 \times 10^{-5}$	90	ABLIKIM	08A	BES2 $e^+e^- \rightarrow J/\psi$

**$\Gamma(\gamma\eta'(958))/\Gamma_{\text{total}}$   $\Gamma_{219}/\Gamma$**

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>5.21 \pm 0.17</math> OUR AVERAGE</b>		Error includes scale factor of 1.4. See the ideogram below.		

$5.64 \pm 0.24 \pm 0.20$	5.0k	<sup>1</sup> ABLIKIM	180	BES3 $\psi(2S) \rightarrow \pi^+\pi^-\gamma\gamma\gamma$
$4.78 \pm 0.22 \pm 0.08$		<sup>2</sup> ABLIKIM	11	BES3 $J/\psi \rightarrow \eta'\gamma$
$5.24 \pm 0.12 \pm 0.11$		PEDLAR	09	CLE3 $J/\psi \rightarrow \eta'\gamma$
$5.55 \pm 0.44$	35k	ABLIKIM	06E	BES2 $J/\psi \rightarrow \eta'\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •


$4.50 \pm 0.14 \pm 0.53$		BOLTON	92B	MRK3 $J/\psi \rightarrow \gamma\pi^+\pi^-\eta, \eta \rightarrow \gamma\gamma$
$4.30 \pm 0.31 \pm 0.71$		BOLTON	92B	MRK3 $J/\psi \rightarrow \gamma\pi^+\pi^-\eta, \eta \rightarrow \pi^+\pi^-\pi^0$
$4.04 \pm 0.16 \pm 0.85$	622	AUGUSTIN	90	DM2 $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
$4.39 \pm 0.09 \pm 0.66$	2420	AUGUSTIN	90	DM2 $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
$4.1 \pm 0.3 \pm 0.6$		BLOOM	83	CBAL $e^+e^- \rightarrow 3\gamma + \text{hadrons}$
$2.9 \pm 1.1$	6	BRANDELIK	79C	DASP $e^+e^- \rightarrow 3\gamma$
$2.4 \pm 0.7$	57	BARTEL	76	CNTR $e^+e^- \rightarrow 2\gamma\rho$



<sup>1</sup> ABLIKIM 180 reports  $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta'(958))/\Gamma_{\text{total}}] \times [B(\eta'(958) \rightarrow \gamma\gamma)] = (1.26 \pm 0.02 \pm 0.05) \times 10^{-4}$  from a measurement of  $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta'(958))/\Gamma_{\text{total}}] \times [B(\eta'(958) \rightarrow \gamma\gamma)] \times [B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)]$  assuming  $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.49 \pm 0.30) \times 10^{-2}$ , which we rescale to our best values  $B(\eta'(958) \rightarrow \gamma\gamma) = (2.22 \pm 0.08) \times 10^{-2}$ ,  $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.68 \pm 0.30) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.

<sup>2</sup> ABLIKIM 11 reports  $(4.84 \pm 0.03 \pm 0.24) \times 10^{-3}$  from a measurement of  $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta'(958))/\Gamma_{\text{total}}] / [B(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)]$  assuming  $B(\eta'(958) \rightarrow \pi^+\pi^-\eta) = (43.2 \pm 0.7) \times 10^{-2}$ ,  $B(\eta \rightarrow 2\gamma) = (39.31 \pm 0.20) \times 10^{-2}$ , which we rescale to our best values  $B(\eta'(958) \rightarrow \pi^+\pi^-\eta) = (42.6 \pm 0.7) \times 10^{-2}$ ,  $B(\eta \rightarrow 2\gamma) = (39.41 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.

WEIGHTED AVERAGE  
5.21±0.17 (Error scaled by 1.4)




$\Gamma(\gamma 2\pi^+ 2\pi^-)/\Gamma_{\text{total}}$

$\Gamma_{220}/\Gamma$

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>2.8 ± 0.5 OUR AVERAGE</b>	Error includes scale factor of 1.9. See the ideogram below.		
4.32 ± 0.14 ± 0.73	<sup>1</sup> BISELLO 89B	DM2	$J/\psi \rightarrow 4\pi\gamma$
2.08 ± 0.13 ± 0.35	<sup>2</sup> BISELLO 89B	DM2	$J/\psi \rightarrow 4\pi\gamma$
3.05 ± 0.08 ± 0.45	<sup>2</sup> BALTRUSAIT..86B	MRK3	$J/\psi \rightarrow 4\pi\gamma$
4.85 ± 0.45 ± 1.20	<sup>3</sup> BURKE 82	MRK2	$e^+e^-$

- <sup>1</sup>  $4\pi$  mass less than 3.0 GeV.
- <sup>2</sup>  $4\pi$  mass less than 2.0 GeV.
- <sup>3</sup>  $4\pi$  mass less than 2.5 GeV.



**$\Gamma(\gamma f_2(1270) f_2(1270)) / \Gamma_{\text{total}}$   $\Gamma_{221} / \Gamma$**

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>9.5±0.7±1.6</b>	646 ± 45	ABLIKIM	04M BES	$J/\psi \rightarrow \gamma 2\pi^+ 2\pi^-$

**$\Gamma(\gamma f_2(1270) f_2(1270) (\text{non resonant})) / \Gamma_{\text{total}}$   $\Gamma_{222} / \Gamma$**

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>8.2±0.8±1.7</b>	<sup>1</sup> ABLIKIM	04M BES	$J/\psi \rightarrow \gamma 2\pi^+ 2\pi^-$

<sup>1</sup> Subtracting contribution from intermediate  $\eta_c(1S)$  decays.

**$\Gamma(\gamma K^+ K^- \pi^+ \pi^-) / \Gamma_{\text{total}}$   $\Gamma_{223} / \Gamma$**

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>2.1±0.1±0.6</b>	1516	BAI	00B BES	$J/\psi \rightarrow \gamma K^+ K^0 \pi^+ \pi^-$

**$\Gamma(\gamma f_4(2050)) / \Gamma_{\text{total}}$   $\Gamma_{224} / \Gamma$**

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>2.7±0.5±0.5</b>	<sup>1</sup> BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^-$

<sup>1</sup> Assuming branching fraction  $f_4(2050) \rightarrow \pi\pi / \text{total} = 0.167$ .

$\Gamma(\gamma\omega\omega)/\Gamma_{\text{total}}$   $\Gamma_{225}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.61±0.33 OUR AVERAGE</b>				
6.0 ±4.8 ±1.8		ABLIKIM	08A BES2	$J/\psi \rightarrow \gamma\omega\pi^+\pi^-$
1.41±0.2 ±0.42	120 ± 17	BISELLO	87 SPEC	$e^+e^-$ , hadrons $\gamma$
1.76±0.09±0.45		BALTRUSAIT..85C	MRK3	$e^+e^- \rightarrow$ hadrons $\gamma$

$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\rho^0\rho^0)/\Gamma_{\text{total}}$   $\Gamma_{226}/\Gamma$

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
<b>1.7 ±0.4 OUR AVERAGE</b>	Error includes scale factor of 1.3.		
2.1 ±0.4	BUGG	95 MRK3	$J/\psi \rightarrow \gamma\pi^+\pi^-\pi^+\pi^-$
1.36±0.38	<sup>1,2</sup> BISELLO	89B DM2	$J/\psi \rightarrow 4\pi\gamma$

<sup>1</sup> Estimated by us from various fits.

<sup>2</sup> Includes unknown branching fraction to  $\rho^0\rho^0$ .

$\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$   $\Gamma_{227}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.64±0.12 OUR AVERAGE</b>	Error includes scale factor of 1.3. See the ideogram below.			
2.07±0.16 <sup>+0.02</sup> <sub>-0.07</sub>	2.4k	<sup>1,2</sup> DOBBS	15	$J/\psi \rightarrow \gamma\pi\pi$
1.63±0.26 <sup>+0.02</sup> <sub>-0.06</sub>		<sup>3</sup> ABLIKIM	06V BES2	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$
1.42±0.21 <sup>+0.01</sup> <sub>-0.05</sub>		<sup>4</sup> ABLIKIM	06V BES2	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^0\pi^0$
1.33±0.05±0.20		<sup>5</sup> AUGUSTIN	87 DM2	$J/\psi \rightarrow \gamma\pi^+\pi^-$
1.36±0.09±0.23		<sup>5</sup> BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma\pi^+\pi^-$
1.48±0.25±0.30	178	EDWARDS	82B CBAL	$e^+e^- \rightarrow 2\pi^0\gamma$
2.0 ±0.7	35	ALEXANDER	78 PLUT	$e^+e^-$
1.2 ±0.6	30	<sup>6</sup> BRANDELIK	78B DASP	$e^+e^- \rightarrow \pi^+\pi^-\gamma$

<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.


<sup>2</sup> DOBBS 15 reports  $[\Gamma(J/\psi(1S) \rightarrow \gamma f_2(1270))/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = (1.744 \pm 0.052 \pm 0.122) \times 10^{-3}$  which we divide by our best value  $B(f_2(1270) \rightarrow \pi\pi) = (84.2^{+2.9}_{-0.9}) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>3</sup> ABLIKIM 06V reports  $[\Gamma(J/\psi(1S) \rightarrow \gamma f_2(1270))/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = (1.371 \pm 0.010 \pm 0.222) \times 10^{-3}$  which we divide by our best value  $B(f_2(1270) \rightarrow \pi\pi) = (84.2^{+2.9}_{-0.9}) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>4</sup> ABLIKIM 06V reports  $[\Gamma(J/\psi(1S) \rightarrow \gamma f_2(1270))/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = (1.200 \pm 0.027 \pm 0.174) \times 10^{-3}$  which we divide by our best value  $B(f_2(1270) \rightarrow \pi\pi) = (84.2^{+2.9}_{-0.9}) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>5</sup> Estimated using  $B(f_2(1270) \rightarrow \pi\pi) = 0.843 \pm 0.012$ . The errors do not contain the uncertainty in the  $f_2(1270)$  decay.

<sup>6</sup> Restated by us to take account of spread of E1, M2, E3 transitions.



$\Gamma(\gamma f_2(1270) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$   $\Gamma_{228}/\Gamma$

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
$2.58^{+0.08+0.59}_{-0.09-0.20}$	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

$\Gamma(\gamma f_0(1370) \rightarrow \gamma K \bar{K})/\Gamma_{\text{total}}$   $\Gamma_{229}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	COMMENT
$4.19 \pm 0.73 \pm 1.34$	478	<sup>1</sup> DOBBS 15	$J/\psi \rightarrow \gamma K \bar{K}$

<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\gamma f_0(1370) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$   $\Gamma_{230}/\Gamma$

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
$1.07^{+0.08+0.36}_{-0.07-0.34}$	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

$\Gamma(\gamma f_0(1500) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$   $\Gamma_{231}/\Gamma$

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
$1.59 \pm 0.16^{+0.18}_{-0.56}$	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

$\Gamma(\gamma f_0(1710) \rightarrow \gamma K \bar{K}) / \Gamma_{\text{total}}$   $\Gamma_{232} / \Gamma$

VALUE (units  $10^{-4}$ )    CL%    EVTS    DOCUMENT ID    TECN    COMMENT

**9.5  $\pm$  1.0**    **OUR AVERAGE**    Error includes scale factor of 1.5. See the ideogram below.

8.00 $\pm$ 0.12 $\pm$ 1.24 - 0.08 - 0.40			1	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
11.76 $\pm$ 0.54 $\pm$ 0.94	1.2k		2	DOBBS	15	$J/\psi \rightarrow \gamma K \bar{K}$
9.62 $\pm$ 0.29			3	BAI	03G BES	$J/\psi \rightarrow \gamma K \bar{K}$
5.0 $\pm$ 0.8 $\pm$ 1.8 - 0.4			1,4	BAI	96C BES	$J/\psi \rightarrow \gamma K^+ K^-$
9.2 $\pm$ 1.4 $\pm$ 1.4			1	AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K^+ K^-$
10.4 $\pm$ 1.2 $\pm$ 1.6			1	AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
9.6 $\pm$ 1.2 $\pm$ 1.8			1	BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.6 $\pm$ 0.2 $\pm$ 0.6 - 0.2			1,5	BAI	96C BES	$J/\psi \rightarrow \gamma K^+ K^-$
< 0.8	90		6	BISELLO	89B	$J/\psi \rightarrow 4\pi\gamma$
1.6 $\pm$ 0.4 $\pm$ 0.3			7	BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^-$
3.8 $\pm$ 1.6			8	EDWARDS	82D CBAL	$e^+ e^- \rightarrow \eta\eta\gamma$

<sup>1</sup> Includes unknown branching fraction to  $K^+ K^-$  or  $K_S^0 K_S^0$ . We have multiplied  $K^+ K^-$  measurement by 2, and  $K_S^0 K_S^0$  by 4 to obtain  $K \bar{K}$  result.

<sup>2</sup> Using CLEO-c data but not authored by the CLEO Collaboration.

<sup>3</sup> Includes unknown branching ratio to  $K^+ K^-$  or  $K_S^0 K_S^0$ .


<sup>4</sup> Assuming  $J^P = 2^+$  for  $f_0(1710)$ .

<sup>5</sup> Assuming  $J^P = 0^+$  for  $f_0(1710)$ .

<sup>6</sup> Includes unknown branching fraction to  $\rho^0 \rho^0$ .

<sup>7</sup> Includes unknown branching fraction to  $\pi^+ \pi^-$ .

<sup>8</sup> Includes unknown branching fraction to  $\eta\eta$ .



$\Gamma(\gamma f_0(1710) \rightarrow \gamma \pi \pi) / \Gamma_{\text{total}}$   $\Gamma_{233} / \Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.8 ± 0.5 OUR AVERAGE</b>				
3.72 ± 0.30 ± 0.43	483	<sup>1</sup> DOBBS	15	$J/\psi \rightarrow \gamma \pi \pi$
3.96 ± 0.06 ± 1.12		<sup>2</sup> ABLIKIM	06V BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^+ \pi^-$
3.99 ± 0.15 ± 2.64		<sup>2</sup> ABLIKIM	06V BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^0 \pi^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
2.5 ± 1.6 ± 0.8		BAI	98H BES	$J/\psi \rightarrow \gamma \pi^0 \pi^0$

<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.

<sup>2</sup> Including unknown branching fraction to  $\pi \pi$ .

$\Gamma(\gamma f_0(1710) \rightarrow \gamma \omega \omega) / \Gamma_{\text{total}}$   $\Gamma_{234} / \Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.31 ± 0.06 ± 0.08</b>	180	ABLIKIM	06H BES	$J/\psi \rightarrow \gamma \omega \omega$

$\Gamma(\gamma f_0(1710) \rightarrow \gamma \eta \eta) / \Gamma_{\text{total}}$   $\Gamma_{235} / \Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.35<sup>+0.13+1.24</sup><sub>-0.11-0.74</sub></b>	5.5k	<sup>1</sup> ABLIKIM	13N BES3	$J/\psi \rightarrow \gamma \eta \eta$

<sup>1</sup> From partial wave analysis including all possible combinations of  $0^{++}$ ,  $2^{++}$ , and  $4^{++}$  resonances.

$\Gamma(\gamma\eta)/\Gamma_{\text{total}}$   $\Gamma_{236}/\Gamma$

VALUE (units  $10^{-3}$ )      EVTS      DOCUMENT ID      TECN      COMMENT

**1.108 ± 0.027 OUR AVERAGE**

1.12 ± 0.05 ± 0.01	18.6k	<sup>1</sup> ABLIKIM	180	BES3	$\psi(2S) \rightarrow \pi^+ \pi^- \gamma \gamma \gamma$
1.101 ± 0.029 ± 0.022		PEDLAR	09	CLE3	$J/\psi \rightarrow \eta \gamma$
1.123 ± 0.089	11k	ABLIKIM	06E	BES2	$J/\psi \rightarrow \eta \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.88 ± 0.08 ± 0.11		BLOOM	83	CBAL	$e^+ e^-$
0.82 ± 0.10		BRANDELIK	79C	DASP	$e^+ e^-$
1.3 ± 0.4	21	BARTEL	77	CNTR	$e^+ e^-$

<sup>1</sup> ABLIKIM 180 reports  $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] = (4.42 \pm 0.04 \pm 0.18) \times 10^{-4}$  from a measurement of  $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] \times [B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)]$  assuming  $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.49 \pm 0.30) \times 10^{-2}$ , which we rescale to our best values  $B(\eta \rightarrow 2\gamma) = (39.41 \pm 0.20) \times 10^{-2}$ ,  $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.68 \pm 0.30) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.

$\Gamma(\gamma f_1(1420) \rightarrow \gamma K \bar{K} \pi)/\Gamma_{\text{total}}$   $\Gamma_{237}/\Gamma$

VALUE (units  $10^{-3}$ )      DOCUMENT ID      TECN      COMMENT

**0.79 ± 0.13 OUR AVERAGE**

0.68 ± 0.04 ± 0.24		BAI	00D	BES	$J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$
0.76 ± 0.15 ± 0.21		<sup>1,2</sup> AUGUSTIN	92	DM2	$J/\psi \rightarrow \gamma K \bar{K} \pi$
0.87 ± 0.14 <sup>+0.14</sup> <sub>-0.11</sub>		<sup>1</sup> BAI	90C	MRK3	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$

<sup>1</sup> Included unknown branching fraction  $f_1(1420) \rightarrow K \bar{K} \pi$ .

<sup>2</sup> From fit to the  $K^*(892) K 1^{++}$  partial wave.

$\Gamma(\gamma f_1(1285))/\Gamma_{\text{total}}$   $\Gamma_{238}/\Gamma$

VALUE (units  $10^{-3}$ )      DOCUMENT ID      TECN      COMMENT

**0.61 ± 0.08 OUR AVERAGE**

0.69 ± 0.16 ± 0.20		<sup>1</sup> BAI	04J	BES2	$J/\psi \rightarrow \gamma \gamma \rho^0$
0.61 ± 0.04 ± 0.21		<sup>2</sup> BAI	00D	BES	$J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$
0.45 ± 0.09 ± 0.17		<sup>3</sup> BAI	99	BES	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
0.625 ± 0.063 ± 0.103		<sup>4</sup> BOLTON	92	MRK3	$J/\psi \rightarrow \gamma f_1(1285)$
0.70 ± 0.08 ± 0.16		<sup>5</sup> BOLTON	92B	MRK3	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$

<sup>1</sup> Assuming  $B(f_1(1285) \rightarrow \rho^0 \gamma) = 0.055 \pm 0.013$ .

<sup>2</sup> Assuming  $\Gamma(f_1(1285) \rightarrow K \bar{K} \pi)/\Gamma_{\text{total}} = 0.090 \pm 0.004$ .

<sup>3</sup> Assuming  $\Gamma(f_1(1285) \rightarrow \eta \pi \pi)/\Gamma_{\text{total}} = 0.5 \pm 0.18$ .

<sup>4</sup> Obtained summing the sequential decay channels

$B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow \pi \pi \pi \pi) = (1.44 \pm 0.39 \pm 0.27) \times 10^{-4}$ ;  
 $B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow a_0(980) \pi, a_0(980) \rightarrow \eta \pi) = (3.90 \pm 0.42 \pm 0.87) \times 10^{-4}$ ;

$B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow a_0(980) \pi, a_0(980) \rightarrow K \bar{K}) = (0.66 \pm 0.26 \pm 0.29) \times 10^{-4}$ ;

$B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow \gamma \rho^0) = (0.25 \pm 0.07 \pm 0.03) \times 10^{-4}$ .

<sup>5</sup> Using  $B(f_1(1285) \rightarrow a_0(980) \pi) = 0.37$ , and including unknown branching ratio for  $a_0(980) \rightarrow \eta \pi$ .

$\Gamma(\gamma f_1(1510) \rightarrow \gamma \eta \pi^+ \pi^-) / \Gamma_{\text{total}}$   $\Gamma_{239} / \Gamma$

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
<b><math>4.5 \pm 1.0 \pm 0.7</math></b>	BAI	99	BES $J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$

$\Gamma(\gamma f'_2(1525)) / \Gamma_{\text{total}}$   $\Gamma_{240} / \Gamma$

VALUE (units $10^{-4}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

**$5.7^{+0.8}_{-0.5}$  OUR AVERAGE** Error includes scale factor of 1.5. See the ideogram

below.

$8.0 \pm 0.9 \pm 0.2$	750	1,2	DOBBS	15	$J/\psi \rightarrow \gamma K \bar{K}$
$3.85 \pm 0.17^{+1.91}_{-0.73}$		3	BAI	03G	BES $J/\psi \rightarrow \gamma K \bar{K}$
$3.6 \pm 0.4^{+1.4}_{-0.4}$		3	BAI	96C	BES $J/\psi \rightarrow \gamma K^+ K^-$
$5.6 \pm 1.4 \pm 0.9$		3	AUGUSTIN	88	DM2 $J/\psi \rightarrow \gamma K^+ K^-$
$4.5 \pm 0.4 \pm 0.9$		3	AUGUSTIN	88	DM2 $J/\psi \rightarrow \gamma K_S^0 K_S^0$
$6.8 \pm 1.6 \pm 1.4$		3	BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<3.4	90	4	4 BRANDELIK	79C	DASP $e^+ e^- \rightarrow \pi^+ \pi^- \gamma$
<2.3	90	3	ALEXANDER	78	PLUT $e^+ e^- \rightarrow K^+ K^- \gamma$

<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.


<sup>2</sup> DOBBS 15 reports  $[\Gamma(J/\psi(1S) \rightarrow \gamma f'_2(1525)) / \Gamma_{\text{total}}] \times [B(f'_2(1525) \rightarrow K \bar{K})] = (7.09 \pm 0.46 \pm 0.67) \times 10^{-4}$  which we divide by our best value  $B(f'_2(1525) \rightarrow K \bar{K}) = (88.7 \pm 2.2) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>3</sup> Using  $B(f'_2(1525) \rightarrow K \bar{K}) = 0.888$ .

<sup>4</sup> Assuming isotropic production and decay of the  $f'_2(1525)$  and isospin.



WEIGHTED AVERAGE  
 $5.7 \pm 0.8 - 0.5$  (Error scaled by 1.5)



**$\Gamma(\gamma f'_2(1525) \rightarrow \gamma K_S^0 K_S^0) / \Gamma_{\text{total}}$**   **$\Gamma_{241} / \Gamma$**

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
$7.99^{+0.03+0.69}_{-0.04-0.50}$	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

**$\Gamma(\gamma f'_2(1525) \rightarrow \gamma \eta \eta) / \Gamma_{\text{total}}$**   **$\Gamma_{242} / \Gamma$**

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$3.42^{+0.43+1.37}_{-0.51-1.30}$	5.5k	<sup>1</sup> ABLIKIM	13N BES3	$J/\psi \rightarrow \gamma \eta \eta$

<sup>1</sup> From partial wave analysis including all possible combinations of  $0^{++}$ ,  $2^{++}$ , and  $4^{++}$  resonances.

**$\Gamma(\gamma f_2(1640) \rightarrow \gamma \omega \omega) / \Gamma_{\text{total}}$**   **$\Gamma_{243} / \Gamma$**

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$0.28 \pm 0.05 \pm 0.17$	141	ABLIKIM	06H BES	$J/\psi \rightarrow \gamma \omega \omega$

**$\Gamma(\gamma f_2(1910) \rightarrow \gamma \omega \omega) / \Gamma_{\text{total}}$**   **$\Gamma_{244} / \Gamma$**

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$0.20 \pm 0.04 \pm 0.13$	151	ABLIKIM	06H BES	$J/\psi \rightarrow \gamma \omega \omega$

**$\Gamma(\gamma f_0(1750) \rightarrow \gamma K_S^0 K_S^0) / \Gamma_{\text{total}}$**   **$\Gamma_{245} / \Gamma$**

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
$1.11 \pm 0.06^{+0.19}_{-0.32}$	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

$\Gamma(\gamma f_0(1800) \rightarrow \gamma \omega \phi) / \Gamma_{\text{total}}$   $\Gamma_{246} / \Gamma$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>2.5 ± 0.6 OUR AVERAGE</b>				
$2.00 \pm 0.08^{+1.38}_{-1.64}$	1.3k	ABLIKIM	13J BES3	$J/\psi \rightarrow \gamma \omega \phi$
$2.61 \pm 0.27 \pm 0.65$	95	ABLIKIM	06J BES2	$J/\psi \rightarrow \gamma \omega \phi$

$\Gamma(\gamma f_2(1810) \rightarrow \gamma \eta \eta) / \Gamma_{\text{total}}$   $\Gamma_{247} / \Gamma$

<u>VALUE (units <math>10^{-5}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>COMMENT</u>
<b>5.40<sup>+0.60+3.42</sup><sub>-0.67-2.35</sub></b>	5.5k	<sup>1</sup> ABLIKIM	13N $J/\psi \rightarrow \gamma \eta \eta$

<sup>1</sup> From partial wave analysis including all possible combinations of  $0^{++}$ ,  $2^{++}$ , and  $4^{++}$  resonances.

$\Gamma(\gamma f_2(1950) \rightarrow \gamma K^*(892) \bar{K}^*(892)) / \Gamma_{\text{total}}$   $\Gamma_{248} / \Gamma$

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.7 ± 0.1 ± 0.2</b>	BAI	00B BES	$J/\psi \rightarrow \gamma K^+ K^0 \pi^+ \pi^-$

$\Gamma(\gamma K^*(892) \bar{K}^*(892)) / \Gamma_{\text{total}}$   $\Gamma_{249} / \Gamma$


<u>VALUE (units <math>10^{-3}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>4.0 ± 0.3 ± 1.3</b>	320	<sup>1</sup> BAI	00B BES	$J/\psi \rightarrow \gamma K^+ K^0 \pi^+ \pi^-$

<sup>1</sup> Summed over all charges.

$\Gamma(\gamma \phi \phi) / \Gamma_{\text{total}}$   $\Gamma_{250} / \Gamma$

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>4.0 ± 1.2 OUR AVERAGE</b>				Error includes scale factor of 2.1. See the ideogram below.
$7.5 \pm 0.6 \pm 1.2$	168	BAI	90B MRK3	$J/\psi \rightarrow \gamma 4K$
$3.4 \pm 0.8 \pm 0.6$	$33 \pm 7$	<sup>1</sup> BISELLO	90 DM2	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$
$3.1 \pm 0.7 \pm 0.4$		<sup>1</sup> BISELLO	86B DM2	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$

<sup>1</sup>  $\phi \phi$  mass less than 2.9 GeV,  $\eta_c$  excluded.



$\Gamma(\gamma\rho\rho)/\Gamma_{\text{total}}$					$\Gamma_{251}/\Gamma$
VALUE (units $10^{-3}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.38 \pm 0.07 \pm 0.07</math></b>		49	EATON	84	MRK2 $e^+e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.11	90		PERUZZI	78	MRK1 $e^+e^-$

$\Gamma(\gamma\eta(2225))/\Gamma_{\text{total}}$					$\Gamma_{252}/\Gamma$
VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b><math>3.14^{+0.50}_{-0.19}</math></b>					
$2.40 \pm 0.10^{+2.47}_{-0.18}$	1,2	ABLIKIM	16N BES3	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$	
$4.4 \pm 0.4 \pm 0.8$	196	2 ABLIKIM	08I BES	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$	
$3.3 \pm 0.8 \pm 0.5$		2 BAI	90B MRK3	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$	
$2.7 \pm 0.6 \pm 0.6$		2 BAI	90B MRK3	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$	
$2.4^{+1.5}_{-1.0}$	3,4	BISELLO	89B DM2	$J/\psi \rightarrow 4\pi\gamma$	

<sup>1</sup> From a partial wave analysis of  $J/\psi \rightarrow \gamma\phi\phi$  that also finds significant signals for for  $\eta(2100)$ ,  $0^-+$  phase space,  $f_0(2100)$ ,  $f_2(2010)$ ,  $f_2(2300)$ ,  $f_2(2340)$ , and a previously unseen  $0^-+$  state  $X(2500)$  ( $M = 2470^{+15+101}_{-19-23}$  MeV,  $\Gamma = 230^{+64+56}_{-35-33}$  MeV).

<sup>2</sup> Includes unknown branching fraction to  $\phi\phi$ .

<sup>3</sup> Estimated by us from various fits.

<sup>4</sup> Includes unknown branching fraction to  $\rho^0\rho^0$ .

$\Gamma(\gamma\eta(1760) \rightarrow \gamma\rho^0\rho^0)/\Gamma_{\text{total}}$   $\Gamma_{253}/\Gamma$

VALUE (units $10^{-3}$ )		DOCUMENT ID	TECN	COMMENT
<b>0.13±0.09</b>	1,2	BISELLO	89B DM2	$J/\psi \rightarrow 4\pi\gamma$

<sup>1</sup> Estimated by us from various fits.

<sup>2</sup> Includes unknown branching fraction to  $\rho^0\rho^0$ .

$\Gamma(\gamma\eta(1760) \rightarrow \gamma\omega\omega)/\Gamma_{\text{total}}$   $\Gamma_{254}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.98±0.08±0.32</b>	1045	ABLIKIM	06H BES	$J/\psi \rightarrow \gamma\omega\omega$

$\Gamma(\gamma\eta(1760) \rightarrow \gamma\gamma\gamma)/\Gamma_{\text{total}}$   $\Gamma_{255}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;4.80 × 10<sup>-6</sup></b>	90	ABLIKIM	180 BES3	$\psi(2S) \rightarrow \pi^+\pi^-\gamma\gamma\gamma$

$\Gamma(\gamma X(1835) \rightarrow \gamma\pi^+\pi^-\eta')/\Gamma_{\text{total}}$   $\Gamma_{256}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**2.77<sup>+0.34</sup><sub>-0.40</sub> OUR AVERAGE** Error includes scale factor of 1.1.

3.93±0.38 <sup>+0.31</sup> <sub>-0.84</sub>		<sup>1</sup> ABLIKIM	16J BES3	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$
2.87±0.09 <sup>+0.49</sup> <sub>-0.52</sub>	4265	<sup>2</sup> ABLIKIM	11C BES3	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$
2.2 ± 0.4 ± 0.4	264	ABLIKIM	05R BES2	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$

<sup>1</sup> From a fit of the measured  $\pi^+\pi^-\eta'$  lineshape that accounts for the abrupt distortion observed at the  $p\bar{p}$  threshold with a Flatté formula in addition to known backgrounds and contributors, as well as an *ad hoc* Breit-Wigner ( $M \approx 1919$  MeV;  $\Gamma \approx 51$  MeV) that is required for a good fit. Another explanation for the distortion provided by ABLIKIM 16J is that a second resonance near 1870 MeV interferes with the  $X(1835)$ ; fits to this possibility yield product branching fraction values compatible with that shown within the respective systematic uncertainties.

<sup>2</sup> From a fit of the  $\pi^+\pi^-\eta'$  mass distribution to a combination of  $\gamma f_1(1510)$ ,  $\gamma X(1835)$ , and two unconfirmed states  $\gamma X(2120)$ , and  $\gamma X(2370)$ , for  $M(p\bar{p}) < 2.8$  GeV, and accounting for backgrounds from non- $\eta'$  events and  $J/\psi \rightarrow \pi^0\pi^+\pi^-\eta'$ .

$\Gamma(\gamma X(1835) \rightarrow \gamma p\bar{p})/\Gamma_{\text{total}}$   $\Gamma_{257}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**0.77<sup>+0.15</sup><sub>-0.09</sub> OUR AVERAGE**

0.90 <sup>+0.04</sup> <sub>-0.11</sub> <sup>+0.27</sup> <sub>-0.55</sub>		<sup>1</sup> ABLIKIM	12D BES3	$J/\psi \rightarrow \gamma p\bar{p}$
1.14 <sup>+0.43</sup> <sub>-0.30</sub> <sup>+0.42</sup> <sub>-0.26</sub>	231	<sup>2</sup> ALEXANDER	10 CLEO	$J/\psi \rightarrow \gamma p\bar{p}$
0.70±0.04 <sup>+0.19</sup> <sub>-0.08</sub>		BAI	03F BES2	$J/\psi \rightarrow \gamma p\bar{p}$

<sup>1</sup> From the fit including final state interaction effects in isospin 0  $S$ -wave according to SIBIRTSEV 05A.

<sup>2</sup> From a fit of the  $p\bar{p}$  mass distribution to a combination of  $\gamma X(1835)$ ,  $\gamma R$  with  $M(R) = 2100$  MeV and  $\Gamma(R) = 160$  MeV, and  $\gamma p\bar{p}$  phase space, for  $M(p\bar{p}) < 2.85$  GeV.

$\Gamma(\gamma X(1835) \rightarrow \gamma K_S^0 K_S^0 \eta) / \Gamma_{\text{total}}$   $\Gamma_{258} / \Gamma$

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
$3.31^{+0.33+1.96}_{-0.30-1.29}$	ABLIKIM	15T BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta$

$\Gamma(\gamma X(1835) \rightarrow \gamma \gamma \phi(1020)) / \Gamma_{\text{total}}$   $\Gamma_{259} / \Gamma$

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$1.77 \pm 0.35 \pm 0.25$	305	<sup>1</sup> ABLIKIM	18i BES3	$J/\psi \rightarrow \gamma \gamma \phi(1020)$
$8.09 \pm 1.99 \pm 1.36$	1.3k	<sup>2</sup> ABLIKIM	18i BES3	$J/\psi \rightarrow \gamma \gamma \phi(1020)$

<sup>1</sup> Constructive interference between the  $X(1835)$  and  $\eta(1405)/\eta(1475)$  is assumed in a fit to the  $\gamma \phi$  invariant mass.

<sup>2</sup> Destructive interference between the  $X(1835)$  and  $\eta(1405)/\eta(1475)$  is assumed in a fit to the  $\gamma \phi$  invariant mass.

$\Gamma(\gamma X(1835) \rightarrow \gamma \gamma \gamma) / \Gamma_{\text{total}}$   $\Gamma_{260} / \Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 3.56 \times 10^{-6}$	90	ABLIKIM	180 BES3	$\psi(2S) \rightarrow \pi^+ \pi^- \gamma \gamma \gamma$

$\Gamma(\gamma X(1840) \rightarrow \gamma 3(\pi^+ \pi^-)) / \Gamma_{\text{total}}$   $\Gamma_{261} / \Gamma$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$2.44 \pm 0.36^{+0.60}_{-0.74}$	0.6k	ABLIKIM	13U BES3	$J/\psi \rightarrow \gamma 3(\pi^+ \pi^-)$

$\Gamma(\gamma(K \bar{K} \pi) [J^{PC} = 0^{-+}]) / \Gamma_{\text{total}}$   $\Gamma_{262} / \Gamma$

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
<b>0.7 ± 0.4 OUR AVERAGE</b>	Error includes scale factor of 2.1.		
$0.58 \pm 0.03 \pm 0.20$	<sup>1</sup> BAI	00D BES	$J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$
$2.1 \pm 0.1 \pm 0.7$	<sup>2</sup> BAI	00D BES	$J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$

<sup>1</sup> For a broad structure around 1800 MeV.

<sup>2</sup> For a broad structure around 2040 MeV.

$\Gamma(\gamma \pi^0) / \Gamma_{\text{total}}$   $\Gamma_{263} / \Gamma$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.56 ± 0.17 OUR AVERAGE</b>				
$3.59 \pm 0.20 \pm 0.03$	1.6k	<sup>1</sup> ABLIKIM	180 BES3	$\psi(2S) \rightarrow \pi^+ \pi^- \gamma \gamma \gamma$
$3.63 \pm 0.36 \pm 0.13$		PEDLAR	09 CLE3	$J/\psi \rightarrow \pi^0 \gamma$
$3.13^{+0.65}_{-0.47}$	586	ABLIKIM	06E BES2	$J/\psi \rightarrow \pi^0 \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$3.6 \pm 1.1 \pm 0.7$	BLOOM	83 CBAL	$e^+ e^-$
$7.3 \pm 4.7$	10	BRANDELIK 79C	DASP $e^+ e^-$

<sup>1</sup> ABLIKIM 180 reports  $[\Gamma(J/\psi(1S) \rightarrow \gamma \pi^0) / \Gamma_{\text{total}}] \times [B(\pi^0 \rightarrow 2\gamma)] = (3.57 \pm 0.12 \pm 0.16) \times 10^{-5}$  from a measurement of  $[\Gamma(J/\psi(1S) \rightarrow \gamma \pi^0) / \Gamma_{\text{total}}] \times [B(\pi^0 \rightarrow 2\gamma)] \times [B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)]$  assuming  $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (34.49 \pm 0.30) \times 10^{-2}$ , which we rescale to our best values  $B(\pi^0 \rightarrow 2\gamma) = (98.823 \pm 0.034) \times 10^{-2}$ ,  $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (34.68 \pm 0.30) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.

$\Gamma(\gamma\rho\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{264}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<0.79 \times 10^{-3}$	90	EATON	84	MRK2 $e^+e^-$

$\Gamma(\gamma\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$   $\Gamma_{265}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<0.13 \times 10^{-3}$	90	HENRARD	87	DM2 $e^+e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<0.16 \times 10^{-3}$	90	BAI	98G	BES $e^+e^-$
------------------------	----	-----	-----	--------------

$\Gamma(\gamma f_0(2100) \rightarrow \gamma\eta\eta)/\Gamma_{\text{total}}$   $\Gamma_{266}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$1.13^{+0.09+0.64}_{-0.10-0.28}$	5.5k	<sup>1</sup> ABLIKIM	13N	BES3 $J/\psi \rightarrow \gamma\eta\eta$

<sup>1</sup> From partial wave analysis including all possible combinations of  $0^{++}$ ,  $2^{++}$ , and  $4^{++}$  resonances.

$\Gamma(\gamma f_0(2100) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$   $\Gamma_{267}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$6.24 \pm 0.48 \pm 0.87$	744	<sup>1</sup> DOBBS	15	$J/\psi \rightarrow \gamma\pi\pi$

<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\gamma f_0(2200))/\Gamma_{\text{total}}$   $\Gamma_{268}/\Gamma$

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.5	<sup>1</sup> AUGUSTIN	88	DM2 $J/\psi \rightarrow \gamma K_S^0 K_S^0$
-----	-----------------------	----	---

<sup>1</sup> Includes unknown branching fraction to  $K_S^0 K_S^0$ .

$\Gamma(\gamma f_0(2200) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$   $\Gamma_{269}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$5.86 \pm 0.49 \pm 1.20$	490	<sup>1</sup> DOBBS	15	$J/\psi \rightarrow \gamma K\bar{K}$

<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\gamma f_0(2200) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$   $\Gamma_{270}/\Gamma$

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
$2.72^{+0.08+0.17}_{-0.06-0.47}$	ABLIKIM	18AA	BES3 $J/\psi \rightarrow \gamma K_S^0 K_S^0$

$\Gamma(\gamma f_J(2220))/\Gamma_{\text{total}}$   $\Gamma_{271}/\Gamma$

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

>300			<sup>1</sup> BAI	96B	BES $e^+e^- \rightarrow \gamma\bar{p}p, K\bar{K}$
>250	99.9		<sup>2</sup> HASAN	96	SPEC $\bar{p}p \rightarrow \pi^+\pi^-$
< 2.3	95		<sup>3</sup> AUGUSTIN	88	DM2 $J/\psi \rightarrow \gamma K^+K^-$
< 1.6	95		<sup>3</sup> AUGUSTIN	88	DM2 $J/\psi \rightarrow \gamma K_S^0 K_S^0$
$12.4^{+6.4}_{-5.2} \pm 2.8$		23	<sup>3</sup> BALTRUSAIT..86D	MRK3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
$8.4^{+3.4}_{-2.8} \pm 1.6$		93	<sup>3</sup> BALTRUSAIT..86D	MRK3	$J/\psi \rightarrow \gamma K^+K^-$

<sup>1</sup> Using BARNES 93.

<sup>2</sup> Using BAI 96B.

<sup>3</sup> Includes unknown branching fraction to  $K^+ K^-$  or  $K_S^0 K_S^0$ .

$\Gamma(\gamma f_J(2220) \rightarrow \gamma \pi \pi) / \Gamma_{\text{total}}$   $\Gamma_{272} / \Gamma$

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 3.9</b>	90	1,2 DOBBS	15	$J/\psi \rightarrow \gamma \pi \pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

14 ± 8 ± 4		BAI	98H BES	$J/\psi \rightarrow \gamma \pi^0 \pi^0$
8.4 ± 2.6 ± 3.0		BAI	96B BES	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^+ \pi^-$

<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.

<sup>2</sup> For  $\Gamma = 20/50$  MeV, the 90% CL upper limits for  $\pi^+ \pi^-$  and  $\pi^0 \pi^0$  are  $2.6/5.2 \times 10^{-5}$  and  $1.3/1.9 \times 10^{-5}$ , respectively.

$\Gamma(\gamma f_J(2220) \rightarrow \gamma K \bar{K}) / \Gamma_{\text{total}}$   $\Gamma_{273} / \Gamma$

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 4.1</b>	90	1,2 DOBBS	15	$J/\psi \rightarrow \gamma K \bar{K}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 3.6		<sup>3</sup> DEL-AMO-SA..100	BABR	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma K^+ K^-$
< 2.9		<sup>3</sup> DEL-AMO-SA..100	BABR	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma K_S^0 K_S^0$
6.6 ± 2.9 ± 2.4		BAI	96B BES	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma K^+ K^-$
10.8 ± 4.0 ± 3.2		BAI	96B BES	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma K_S^0 K_S^0$

<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.

<sup>2</sup> For  $\Gamma = 20/50$  MeV, the 90% CL upper limits for  $K^+ K^-$  and  $K_S^0 K_S^0$  are  $1.7/3.1 \times 10^{-5}$  and  $1.2/2.0 \times 10^{-5}$ , respectively.

<sup>3</sup> For spin 2 and helicity 0; other combinations lead to more stringent upper limits.

$\Gamma(\gamma f_J(2220) \rightarrow \gamma p \bar{p}) / \Gamma_{\text{total}}$   $\Gamma_{274} / \Gamma$

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
<b>1.5 ± 0.6 ± 0.5</b>	BAI	96B BES	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma p \bar{p}$

$\Gamma(\gamma f_0(2330) \rightarrow \gamma K_S^0 K_S^0) / \Gamma_{\text{total}}$   $\Gamma_{275} / \Gamma$

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
<b>4.95 ± 0.21 <sup>+0.66</sup> <sub>-0.72</sub></b>	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

$\Gamma(\gamma f_2(2340) \rightarrow \gamma \eta \eta) / \Gamma_{\text{total}}$   $\Gamma_{276} / \Gamma$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>5.60 <sup>+0.62 + 2.37</sup> <sub>-0.65 - 2.07</sub></b>	5.5k	<sup>1</sup> ABLIKIM	13N BES3	$J/\psi \rightarrow \gamma \eta \eta$

<sup>1</sup> From partial wave analysis including all possible combinations of  $0^{++}$ ,  $2^{++}$ , and  $4^{++}$  resonances.

$\Gamma(\gamma f_2(2340) \rightarrow \gamma K_S^0 K_S^0) / \Gamma_{\text{total}}$   $\Gamma_{277} / \Gamma$

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
<b>5.54 <sup>+0.34 + 3.82</sup> <sub>-0.40 - 1.49</sub></b>	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

$\Gamma(\gamma f_0(1500) \rightarrow \gamma \pi \pi) / \Gamma_{\text{total}}$   $\Gamma_{278} / \Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.09 ± 0.24 OUR AVERAGE</b>				
1.21 ± 0.29 ± 0.24	174	<sup>1</sup> DOBBS	15	$J/\psi \rightarrow \gamma \pi \pi$
1.00 ± 0.03 ± 0.45		<sup>2</sup> ABLIKIM	06V BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^+ \pi^-$
1.02 ± 0.09 ± 0.45		<sup>2</sup> ABLIKIM	06V BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^0 \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
5.7 ± 0.8		<sup>3,4</sup> BUGG	95 MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$

- <sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.
- <sup>2</sup> Including unknown branching fraction to  $\pi \pi$ .
- <sup>3</sup> Including unknown branching ratio for  $f_0(1500) \rightarrow \pi^+ \pi^- \pi^+ \pi^-$ .
- <sup>4</sup> Assuming that  $f_0(1500)$  decays only to two  $S$ -wave dipions.

$\Gamma(\gamma f_0(1500) \rightarrow \gamma \eta \eta) / \Gamma_{\text{total}}$   $\Gamma_{279} / \Gamma$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.65<sup>+0.26+0.51</sup><sub>-0.31-1.40</sub></b>	5.5k	<sup>1</sup> ABLIKIM	13N BES3	$J/\psi \rightarrow \gamma \eta \eta$

- <sup>1</sup> From partial wave analysis including all possible combinations of  $0^{++}$ ,  $2^{++}$ , and  $4^{++}$  resonances.

$\Gamma(\gamma A \rightarrow \gamma \text{invisible}) / \Gamma_{\text{total}}$   $\Gamma_{280} / \Gamma$   
 (narrow state  $A$  with  $m_A < 960$  MeV)

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 6.3 × 10<sup>-6</sup></b>	90	<sup>1</sup> INSLER	10 CLEO	$e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$

- <sup>1</sup> The limit varies with mass  $m_A$  of a narrow state  $A$  and is  $4.3 \times 10^{-6}$  for  $m_A = 0$  MeV, reaches its largest value of  $6.3 \times 10^{-6}$  at  $m_A = 500$  MeV, and is  $3.6 \times 10^{-6}$  at  $m_A = 960$  MeV.

$\Gamma(\gamma A^0 \rightarrow \gamma \mu^+ \mu^-) / \Gamma_{\text{total}}$   $\Gamma_{281} / \Gamma$   
 (narrow state  $A^0$  with  $0.2 \text{ GeV} < m_{A^0} < 3 \text{ GeV}$ )

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 0.5 × 10<sup>-5</sup></b>	90	<sup>1</sup> ABLIKIM	16E BES3	$J/\psi \rightarrow \gamma \mu^+ \mu^-$
<b>&lt; 2.1 × 10<sup>-5</sup></b>	90	<sup>2</sup> ABLIKIM	12 BES3	$J/\psi \rightarrow \gamma \mu^+ \mu^-$

- • • We do not use the following data for averages, fits, limits, etc. • • •
- <sup>1</sup> For a narrow scalar or pseudoscalar,  $A^0$ , with a mass in the range 0.212–3 GeV. The measured 90% CL limit as a function of  $m_{A^0}$  is in the range  $(2.8\text{--}495.3) \times 10^{-8}$ .
- <sup>2</sup> For a narrow scalar or pseudoscalar,  $A^0$ , with a mass in the range 0.21–3.00 GeV. The measured 90% CL limit as a function of  $m_{A^0}$  ranges from  $4 \times 10^{-7}$  to  $2.1 \times 10^{-5}$ .

————— DALITZ DECAYS —————

$\Gamma(\pi^0 e^+ e^-) / \Gamma_{\text{total}}$   $\Gamma_{282} / \Gamma$

VALUE (units $10^{-7}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>7.56 ± 1.32 ± 0.50</b>	39	ABLIKIM	14I BES3	$J/\psi \rightarrow \pi^0 e^+ e^-$



$\Gamma(\eta e^+ e^-)/\Gamma_{\text{total}}$   $\Gamma_{283}/\Gamma$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

<b><math>1.43 \pm 0.04 \pm 0.06</math></b>	2.47k	<sup>1,2</sup> ABLIKIM	19A	BES3 $J/\psi \rightarrow \eta e^+ e^-$
--	-------	------------------------	-----	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.16 \pm 0.07 \pm 0.06$	320	<sup>1</sup> ABLIKIM	14I	BES3 $J/\psi \rightarrow \eta e^+ e^-$
--------------------------	-----	----------------------	-----	--

<sup>1</sup> Using both  $\eta \rightarrow \gamma\gamma$  and  $\eta \rightarrow \pi^+ \pi^- \pi^0$  decays.

<sup>2</sup> Approximation of the transition form factor squared as an incoherent sum of the  $\rho$ -meson and one-pole non-resonant amplitudes gives the pole mass  $m(\Lambda) = 2.84 \pm 0.11 \pm 0.08$  GeV. Supersedes ABLIKIM 14I.

$\Gamma(\eta'(958) e^+ e^-)/\Gamma_{\text{total}}$   $\Gamma_{284}/\Gamma$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

<b><math>6.59 \pm 0.07 \pm 0.17</math></b>	8.9k	<sup>1</sup> ABLIKIM	19H	BES3 $J/\psi \rightarrow \eta'(958) e^+ e^-$
--	------	----------------------	-----	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

$5.81 \pm 0.16 \pm 0.31$	1.4k	<sup>1,2</sup> ABLIKIM	14I	BES3 $J/\psi \rightarrow \eta'(958) e^+ e^-$
--------------------------	------	------------------------	-----	--

<sup>1</sup> Using both  $\eta' \rightarrow \gamma\pi^+ \pi^-$  and  $\eta' \rightarrow \pi^+ \pi^- \eta$  decays.

<sup>2</sup> Superseded by ABLIKIM 19H.

$\Gamma(\eta U \rightarrow \eta e^+ e^-)/\Gamma_{\text{total}}$   $\Gamma_{285}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<b><math>&lt; 9.11 \times 10^{-7}</math></b>	90	<sup>1</sup> ABLIKIM	19A	BES3 $J/\psi \rightarrow \eta e^+ e^-$
--	----	----------------------	-----	--

<sup>1</sup> For a dark photon  $U$  with mass between 10 and 2400 MeV. Obtained 90% C.L. limits as a function of  $m_U$  range from  $1.9 \times 10^{-8}$  to  $91.1 \times 10^{-8}$ .

$\Gamma(\eta'(958) U \rightarrow \eta'(958) e^+ e^-)/\Gamma_{\text{total}}$   $\Gamma_{286}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<b><math>&lt; 2.0 \times 10^{-7}</math></b>	90	<sup>1</sup> ABLIKIM	19H	BES3 $J/\psi \rightarrow \eta'(958) e^+ e^-$
---	----	----------------------	-----	--

<sup>1</sup> For a dark photon  $U$  with mass between 100 and 2100 MeV. Obtained 90% C.L. limits as a function of  $m_U$  range from  $1.8 \times 10^{-8}$  to  $2.0 \times 10^{-7}$ . The corresponding limits on the branching fraction  $J/\psi \rightarrow \eta' U$  range from  $5.7 \times 10^{-8}$  to  $7.4 \times 10^{-7}$ .

———— WEAK DECAYS ————

$\Gamma(D^- e^+ \nu_e + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{287}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<b><math>&lt; 1.2 \times 10^{-5}</math></b>	90	ABLIKIM	06M	BES2 $e^+ e^- \rightarrow J/\psi$
---	----	---------	-----	-----------------------------------

$\Gamma(\bar{D}^0 e^+ e^- + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{288}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<b><math>&lt; 8.5 \times 10^{-8}</math></b>	90	<sup>1</sup> ABLIKIM	17AF	BES3 $e^+ e^- \rightarrow J/\psi$
---	----	----------------------	------	-----------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 1.1 \times 10^{-5}$	90	ABLIKIM	06M	BES2 $e^+ e^- \rightarrow J/\psi$
------------------------	----	---------	-----	-----------------------------------

<sup>1</sup> Using  $D^0$  decays to  $K^- \pi^+$ ,  $K^- \pi^+ \pi^0$ , and  $K^- \pi^+ \pi^+ \pi^-$ .

$\Gamma(D_s^- e^+ \nu_e + \text{c.c.})/\Gamma_{\text{total}}$					$\Gamma_{289}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.3 \times 10^{-6}$	90	ABLIKIM	14R	BES3 $e^+ e^- \rightarrow J/\psi$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
$<3.6 \times 10^{-5}$	90	<sup>1</sup> ABLIKIM	06M	BES2 $e^+ e^- \rightarrow J/\psi$	
<sup>1</sup> Using $B(D_s^- \rightarrow \phi \pi^-) = 4.4 \pm 0.5\%$ .					

$\Gamma(D_s^{*-} e^+ \nu_e + \text{c.c.})/\Gamma_{\text{total}}$					$\Gamma_{290}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.8 \times 10^{-6}$	90	ABLIKIM	14R	BES3 $e^+ e^- \rightarrow J/\psi$	

$\Gamma(D^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$					$\Gamma_{291}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<7.5 \times 10^{-5}$	90	ABLIKIM	08J	BES2 $e^+ e^- \rightarrow J/\psi$	

$\Gamma(\bar{D}^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$					$\Gamma_{292}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.7 \times 10^{-4}$	90	ABLIKIM	08J	BES2 $e^+ e^- \rightarrow J/\psi$	

$\Gamma(\bar{D}^0 \bar{K}^{*0} + \text{c.c.})/\Gamma_{\text{total}}$					$\Gamma_{293}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<2.5 \times 10^{-6}$	90	ABLIKIM	14K	BES3 $e^+ e^- \rightarrow J/\psi$	

$\Gamma(D_s^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$					$\Gamma_{294}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.3 \times 10^{-4}$	90	ABLIKIM	08J	BES2 $e^+ e^- \rightarrow J/\psi$	

$\Gamma(D_s^- \rho^+ + \text{c.c.})/\Gamma_{\text{total}}$					$\Gamma_{295}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.3 \times 10^{-5}$	90	ABLIKIM	14K	BES3 $e^+ e^- \rightarrow J/\psi$	

———— CHARGE CONJUGATION (C), PARITY (P), ————  
 ———— LEPTON FAMILY NUMBER (LF) VIOLATING MODES ————

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_{296}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$< 2.7 \times 10^{-7}$	90	ABLIKIM	14Q	BES3 $\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
$< 0.5 \times 10^{-5}$	90	ADAMS	08	CLEO $\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$	
$< 1.6 \times 10^{-4}$	90	<sup>1</sup> WICHT	08	BELL $B^\pm \rightarrow K^\pm \gamma\gamma$	
$< 2.2 \times 10^{-5}$	90	ABLIKIM	07J	BES2 $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$	
$< 50 \times 10^{-5}$	90	BARTEL	77	CNTR $e^+ e^-$	

<sup>1</sup> WICHT 08 reports  $[\Gamma(J/\psi(1S) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow J/\psi(1S) K^+)] < 0.16 \times 10^{-6}$  which we divide by our best value  $B(B^+ \rightarrow J/\psi(1S) K^+) = 1.010 \times 10^{-3}$ .

$\Gamma(\gamma\phi)/\Gamma_{\text{total}}$					$\Gamma_{297}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.4 \times 10^{-6}$	90	ABLIKIM	14Q	BES3 $\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$	

$\Gamma(e^\pm \mu^\mp)/\Gamma_{\text{total}}$					$\Gamma_{298}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.6 \times 10^{-7}$	90	ABLIKIM	13L	BES3	$e^+e^- \rightarrow J/\psi$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<1.1 \times 10^{-6}$	90	BAI	03D	BES	$e^+e^- \rightarrow J/\psi$

$\Gamma(e^\pm \tau^\mp)/\Gamma_{\text{total}}$					$\Gamma_{299}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<8.3 \times 10^{-6}$	90	ABLIKIM	04	BES	$e^+e^- \rightarrow J/\psi$

$\Gamma(\mu^\pm \tau^\mp)/\Gamma_{\text{total}}$					$\Gamma_{300}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<2.0 \times 10^{-6}$	90	ABLIKIM	04	BES	$e^+e^- \rightarrow J/\psi$

### OTHER DECAYS

$\Gamma(\text{invisible})/\Gamma(e^+e^-)$					$\Gamma_{301}/\Gamma_5$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<6.6 \times 10^{-2}$	90	LEES	13I	BABR	$B \rightarrow K^{(*)} J/\psi$

$\Gamma(\text{invisible})/\Gamma(\mu^+\mu^-)$					$\Gamma_{301}/\Gamma_7$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.2 \times 10^{-2}$	90	ABLIKIM	08G	BES2	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$

### $J/\psi(1S)$ REFERENCES

ABLIKIM	19A	PR D99 012006	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	19H	PR D99 012013	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	19N	PR D99 032006	M. Ablikim <i>et al.</i>	(BES III Collab.)
LU	19	PR D99 032003	P.-C. Lu <i>et al.</i>	(BELLE Collab.)
ABLIKIM	18AA	PR D98 072003	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	18AB	PR D98 072005	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	18D	PRL 121 022001	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	18I	PR D97 051101	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	18O	PR D97 072014	M. Ablikim <i>et al.</i>	(BES III Collab.)
ANASHIN	18A	JHEP 1805 119	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
LEES	18	PR D97 052007	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	18E	PR D98 112015	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	17AF	PR D96 111101	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	17AH	PR D96 112001	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	17AK	PR D96 112012	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	17E	PL B770 217	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	17L	PR D95 052003	M. Ablikim <i>et al.</i>	(BES III Collab.)
LEES	17A	PR D95 052001	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	17C	PR D95 072007	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	17D	PR D95 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	16E	PR D93 052005	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	16J	PRL 117 042002	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	16K	PR D93 052010	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	16L	PR D93 072003	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	16M	PR D93 072008	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	16N	PR D93 112011	M. Ablikim	(BES III Collab.)
ABLIKIM	16P	PR D94 072005	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	16Q	PL B761 98	M. Ablikim <i>et al.</i>	(BES III Collab.)
PDG	16	CP C40 100001	C. Patrignani <i>et al.</i>	(PDG Collab.)
AAIJ	15BI	EPJ C75 311	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	15AE	PR D92 052003	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	15H	PR D91 052017	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	15K	PR D91 112001	M. Ablikim <i>et al.</i>	(BES III Collab.)

ABLIKIM	15P	PR D92 012007	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	15T	PRL 115 091803	M. Ablikim <i>et al.</i>	(BES III Collab.)
ANASHIN	15	PL B749 50	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)
LEES	15J	PR D92 072008	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	14I	PR D89 092008	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	14K	PR D89 071101	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	14N	PR D90 052009	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	14Q	PR D90 092002	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	14R	PR D90 112014	M. Ablikim <i>et al.</i>	(BES III Collab.)
ANASHIN	14	PL B738 391	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
AULCHENKO	14	PL B731 227	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)
LEES	14H	PR D89 092002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	13F	PR D87 052007	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	13I	PR D87 032003	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	13J	PR D87 032008	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	13L	PR D87 112007	Ablikim M. <i>et al.</i>	(BES III Collab.)
ABLIKIM	13N	PR D87 092009	Ablikim M. <i>et al.</i>	(BES III Collab.)
ABLIKIM	13P	PR D87 112004	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	13R	PR D88 032007	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	13U	PR D88 091502	M. Ablikim <i>et al.</i>	(BES III Collab.)
LEES	13I	PR D87 112005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	13O	PR D87 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	13Q	PR D88 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	13Y	PR D88 072009	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	12	PR D85 092012	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	12B	PR D86 032008	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	12C	PR D86 032014	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	12D	PRL 108 112003	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	12H	PL B710 594	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	12P	CP C36 1031	M. Ablikim <i>et al.</i>	(BES II Collab.)
LEES	12E	PR D85 112009	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	12F	PR D86 012008	J.P. Lees <i>et al.</i>	(BABAR Collab.)
METREVELI	12	PR D85 092007	Z. Metreveli <i>et al.</i>	(NWES, FLOR, WAYN+)
ABLIKIM	11	PR D83 012003	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	11C	PRL 106 072002	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	11D	PR D83 032003	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	10C	PL B685 27	M. Ablikim <i>et al.</i>	(BES II Collab.)
ABLIKIM	10E	PL B693 88	M. Ablikim <i>et al.</i>	(BES II Collab.)
ALEXANDER	10	PR D82 092002	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
ANASHIN	10	PL B685 134	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
DEL-AMO-SA...	10O	PRL 105 172001	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
INSLER	10	PR D81 091101	J. Insler <i>et al.</i>	(CLEO Collab.)
ABLIKIM	09	PL B676 25	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	09B	PR D80 052004	M. Ablikim <i>et al.</i>	(BES II Collab.)
MITCHELL	09	PRL 102 011801	R.E. Mitchell <i>et al.</i>	(CLEO Collab.)
PEDLAR	09	PR D79 111101	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)
SHEN	09	PR D80 031101	C.P. Shen <i>et al.</i>	(BELLE Collab.)
ABLIKIM	08	EPJ C53 15	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08A	PR D77 012001	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08C	PL B659 789	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08E	PR D77 032005	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08F	PRL 100 102003	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08G	PRL 100 192001	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08I	PL B662 330	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08J	PL B663 297	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08O	PR D78 092005	M. Ablikim <i>et al.</i>	(BES Collab.)
ADAMS	08	PRL 101 101801	G.S. Adams <i>et al.</i>	(CLEO Collab.)
AUBERT	08S	PR D77 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)
BESSON	08	PR D78 032012	D. Besson <i>et al.</i>	(CLEO Collab.)
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)
WICHT	08	PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)
ABLIKIM	07H	PR D76 092003	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	07J	PR D76 117101	M. Ablikim <i>et al.</i>	(BES Collab.)
ANDREOTTI	07	PL B654 74	M. Andreotti <i>et al.</i>	(Femilab E835 Collab.)
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)
Also		PR D77 119902E (errat.)	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07BD	PR D76 092006	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABLIKIM	06	PL B632 181	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06C	PL B633 681	M. Ablikim <i>et al.</i>	(BES Collab.)

ABLIKIM	06E	PR D73 052008	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06F	PR D73 052007	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06H	PR D73 112007	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06J	PRL 96 162002	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06K	PRL 97 062001	M. Ablikim <i>et al.</i>	(BES II Collab.)
ABLIKIM	06M	PL B639 418	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06V	PL B642 441	M. Ablikim <i>et al.</i>	(BES Collab.)
ADAMS	06A	PR D73 051103	G.S. Adams <i>et al.</i>	(CLEO Collab.)
AUBERT	06B	PR D73 012005	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	06D	PR D73 052003	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	06E	PRL 96 052002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,BE	06D	PR D74 091103	B. Aubert <i>et al.</i>	(BABAR Collab.)
WU	06	PRL 97 162003	C.-H. Wu <i>et al.</i>	(BELLE Collab.)
ABLIKIM	05	PL B607 243	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05B	PR D71 032003	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05C	PL B610 192	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05H	PR D72 012002	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05R	PRL 95 262001	M. Ablikim <i>et al.</i>	(BES Collab.)
AUBERT	05D	PR D71 052001	B. Aubert <i>et al.</i>	(BABAR Collab.)
LI	05C	PR D71 111103	Z. Li <i>et al.</i>	(CLEO Collab.)
SIBIRTSEV	05A	PR D71 054010	A. Sibirtsev, J. Haidenbauer	
ABLIKIM	04	PL B598 172	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	04M	PR D70 112008	M. Ablikim <i>et al.</i>	(BES Collab.)
AUBERT	04	PR D69 011103	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	04N	PR D70 072004	B. Aubert <i>et al.</i>	(BABAR Collab.)
BAI	04	PL B578 16	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04A	PR D69 012003	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04D	PL B589 7	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04E	PL B591 42	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04G	PR D70 012004	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04H	PR D70 012005	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04J	PL B594 47	J.Z. Bai <i>et al.</i>	(BES Collab.)
SETH	04	PR D69 097503	K.K. Seth	
AULCHENKO	03	PL B573 63	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)
BAI	03D	PL B561 49	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	03F	PRL 91 022001	J.Z. Bai <i>et al.</i>	(BES II Collab.)
BAI	03G	PR D68 052003	J.Z. Bai <i>et al.</i>	(BES Collab.)
HUANG	03	PRL 91 241802	H.-C. Huang <i>et al.</i>	(BELLE Collab.)
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>	
BAI	00	PRL 84 594	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	00B	PL B472 200	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	00D	PL B476 25	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	99	PL B446 356	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	99C	PRL 83 1918	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98D	PR D58 092006	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98G	PL B424 213	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98H	PRL 81 1179	J.Z. Bai <i>et al.</i>	(BES Collab.)
BALDINI	98	PL B444 111	R. Baldini <i>et al.</i>	(FENICE Collab.)
ARMSTRONG	96	PR D54 7067	T.A. Armstrong <i>et al.</i>	(E760 Collab.)
BAI	96B	PRL 76 3502	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	96C	PRL 77 3959	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	96D	PR D54 1221	J.Z. Bai <i>et al.</i>	(BES Collab.)
GRIBUSHIN	96	PR D53 4723	A. Gribushin <i>et al.</i>	(E672 Collab., E706 Collab.)
HASAN	96	PL B388 376	A. Hasan, D.V. Bugg	(BRUN, LOQM)
BAI	95B	PL B355 374	J.Z. Bai <i>et al.</i>	(BES Collab.)
BUGG	95	PL B353 378	D.V. Bugg <i>et al.</i>	(LOQM, PNPI, WASH)
ANTONELLI	93	PL B301 317	A. Antonelli <i>et al.</i>	(FENICE Collab.)
ARMSTRONG	93B	PR D47 772	T.A. Armstrong <i>et al.</i>	(FNAL E760 Collab.)
BARNES	93	PL B309 469	P.D. Barnes <i>et al.</i>	(PS185 Collab.)
AUGUSTIN	92	PR D46 1951	J.E. Augustin, G. Cosme	(DM2 Collab.)
BOLTON	92	PL B278 495	T. Bolton <i>et al.</i>	(Mark III Collab.)
BOLTON	92B	PRL 69 1328	T. Bolton <i>et al.</i>	(Mark III Collab.)
COFFMAN	92	PRL 68 282	D.M. Coffman <i>et al.</i>	(Mark III Collab.)
HSUEH	92	PR D45 2181	S. Hsueh, S. Palestini	(FNAL, TORI)
BISELLO	91	NP B350 1	D. Bisello <i>et al.</i>	(DM2 Collab.)
AUGUSTIN	90	PR D42 10	J.E. Augustin <i>et al.</i>	(DM2 Collab.)
BAI	90B	PRL 65 1309	Z. Bai <i>et al.</i>	(Mark III Collab.)
BAI	90C	PRL 65 2507	Z. Bai <i>et al.</i>	(Mark III Collab.)
BISELLO	90	PL B241 617	D. Bisello <i>et al.</i>	(DM2 Collab.)
COFFMAN	90	PR D41 1410	D.M. Coffman <i>et al.</i>	(Mark III Collab.)

JOUSSET	90	PR D41 1389	J. Jousset <i>et al.</i>	(DM2 Collab.)
ALEXANDER	89	NP B320 45	J.P. Alexander <i>et al.</i>	(LBL, MICH, SLAC)
AUGUSTIN	89	NP B320 1	J.E. Augustin, G. Cosme	(DM2 Collab.)
BISELLO	89B	PR D39 701	G. Busetto <i>et al.</i>	(DM2 Collab.)
AUGUSTIN	88	PRL 60 2238	J.E. Augustin <i>et al.</i>	(DM2 Collab.)
COFFMAN	88	PR D38 2695	D.M. Coffman <i>et al.</i>	(Mark III Collab.)
FALVARD	88	PR D38 2706	A. Falvard <i>et al.</i>	(CLER, FRAS, LALO+)
AUGUSTIN	87	ZPHY C36 369	J.E. Augustin <i>et al.</i>	(LALO, CLER, FRAS+)
BAGLIN	87	NP B286 592	C. Baglin <i>et al.</i>	(LAPP, CERN, GENO, LYON+)
BALTRUSAIT...	87	PR D35 2077	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)
BECKER	87	PRL 59 186	J.J. Becker <i>et al.</i>	(Mark III Collab.)
BISELLO	87	PL B192 239	D. Bisello <i>et al.</i>	(PADO, CLER, FRAS+)
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)
HENRARD	87	NP B292 670	P. Henrard <i>et al.</i>	(CLER, FRAS, LALO+)
PALLIN	87	NP B292 653	D. Pallin <i>et al.</i>	(CLER, FRAS, LALO, PADO)
BALTRUSAIT...	86	PR D33 629	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)
BALTRUSAIT...	86B	PR D33 1222	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)
BALTRUSAIT...	86D	PRL 56 107	R.M. Baltrusaitis	(CIT, UCSC, ILL, SLAC+)
BISELLO	86B	PL B179 294	D. Bisello <i>et al.</i>	(DM2 Collab.)
GAISER	86	PR D34 711	J. Gaiser <i>et al.</i>	(Crystal Ball Collab.)
BALTRUSAIT...	85C	PRL 55 1723	R.M. Baltrusaitis <i>et al.</i>	(CIT, UCSC+)
BALTRUSAIT...	85D	PR D32 566	R.M. Baltrusaitis <i>et al.</i>	(CIT, UCSC+)
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)
BALTRUSAIT...	84	Translated from YAF 41 733	R.M. Baltrusaitis <i>et al.</i>	(CIT, UCSC+)
EATON	84	PR D29 804	M.W. Eaton <i>et al.</i>	(LBL, SLAC)
BLOOM	83	ARNS 33 143	E.D. Bloom, C. Peck	(SLAC, CIT)
EDWARDS	83B	PRL 51 859	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
FRANKLIN	83	PRL 51 963	M.E.B. Franklin <i>et al.</i>	(LBL, SLAC)
BURKE	82	PRL 49 632	D.L. Burke <i>et al.</i>	(LBL, SLAC)
EDWARDS	82B	PR D25 3065	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
EDWARDS	82D	PRL 48 458	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
Also		ARNS 33 143	E.D. Bloom, C. Peck	(SLAC, CIT)
EDWARDS	82E	PRL 49 259	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
LEMOIGNE	82	PL 113B 509	Y. Lemoigne <i>et al.</i>	(SACL, LOIC, SHMP+)
BESCH	81	ZPHY C8 1	H.J. Besch <i>et al.</i>	(BONN, DESY, MANZ)
GIDAL	81	PL 107B 153	G. Gidal <i>et al.</i>	(SLAC, LBL)
PARTRIDGE	80	PRL 44 712	R. Partridge <i>et al.</i>	(CIT, HARV, PRIN+)
SCHARRE	80	PL 97B 329	D.L. Scharre <i>et al.</i>	(SLAC, LBL)
ZHOLENTZ	80	PL 96B 214	A.A. Zholents <i>et al.</i>	(NOVO)
Also		SJNP 34 814	A.A. Zholents <i>et al.</i>	(NOVO)
BRANDELIK	79C	Translated from YAF 34 1471	R. Brandelik <i>et al.</i>	(DASP Collab.)
ALEXANDER	78	ZPHY C1 233	G. Alexander <i>et al.</i>	(DESY, HAMB, SIEG+)
BESCH	78	PL 72B 493	H.J. Besch <i>et al.</i>	(BONN, DESY, MANZ)
BRANDELIK	78B	PL 74B 292	R. Brandelik <i>et al.</i>	(DASP Collab.)
PERUZZI	78	PR D17 2901	I. Peruzzi <i>et al.</i>	(SLAC, LBL)
BARTEL	77	PL 66B 489	W. Bartel <i>et al.</i>	(DESY, HEIDP)
BURMESTER	77D	PL 72B 135	J. Burmester <i>et al.</i>	(DESY, HAMB, SIEG+)
FELDMAN	77	PRPL 33C 285	G.J. Feldman, M.L. Perl	(LBL, SLAC)
VANNUCCI	77	PR D15 1814	F. Vannucci <i>et al.</i>	(SLAC, LBL)
BARTEL	76	PL 64B 483	W. Bartel <i>et al.</i>	(DESY, HEIDP)
BRAUNSCH...	76	PL 63B 487	W. Braunschweig <i>et al.</i>	(DASP Collab.)
JEAN-MARIE	76	PRL 36 291	B. Jean-Marie <i>et al.</i>	(SLAC, LBL) IG
BALDINI...	75	PL 58B 471	R. Baldini-Celio <i>et al.</i>	(FRAS, ROMA)
BOYARSKI	75	PRL 34 1357	A.M. Boyarski <i>et al.</i>	(SLAC, LBL) JPC
DASP	75	PL 56B 491	W. Braunschweig <i>et al.</i>	(DASP Collab.)
ESPOSITO	75B	LNC 14 73	B. Esposito <i>et al.</i>	(FRAS, NAPL, PADO+)
FORD	75	PRL 34 604	R.L. Ford <i>et al.</i>	(SLAC, PENN)