


$f_2(1950)$ 

$$I^G(J^{PC}) = 0^+(2^{++})$$

 **$f_2(1950)$  MASS**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>1936±12 OUR AVERAGE</b>	Error includes scale factor of 1.3. See the ideogram below.		
1940±50	BAI	00A	BES $J/\psi \rightarrow \gamma(\pi^+\pi^-\pi^+\pi^-)$
1980±22	<sup>1</sup> BARBERIS	00C	450 $pp \rightarrow pp4\pi$
1940±22	<sup>2</sup> BARBERIS	00C	450 $pp \rightarrow pp2\pi2\pi^0$
1960±30	BARBERIS	97B	OMEG 450 $pp \rightarrow pp2(\pi^+\pi^-)$
1918±12	ANTINORI	95	OMEG 300,450 $pp \rightarrow pp2(\pi^+\pi^-)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
2038 <sup>+13+12</sup> -11-73	<sup>3</sup> UEHARA	09	BELL 10.6 $e^+e^- \rightarrow e^+e^-\pi^0\pi^0$
1930±25	<sup>4</sup> BINON	05	GAMS 33 $\pi^-p \rightarrow \eta\eta n$
1980± 2±14	ABE	04	BELL 10.6 $e^+e^- \rightarrow e^+e^-K^+K^-$
1867±46	<sup>5</sup> AMSLER	02	CBAR 0.9 $\bar{p}p \rightarrow \pi^0\eta\eta, \pi^0\pi^0\pi^0$
2010±25	ANISOVICH	00J	SPEC
1980±50	ANISOVICH	99B	SPEC 1.35–1.94 $p\bar{p} \rightarrow \eta\eta\pi^0$
~ 1990	<sup>6</sup> OAKDEN	94	RVUE 0.36–1.55 $\bar{p}p \rightarrow \pi\pi$
1950±15	<sup>7</sup> ASTON	91	LASS 11 $K^-p \rightarrow \Lambda K\bar{K}\pi\pi$

<sup>1</sup> Decaying into  $\pi^+\pi^-2\pi^0$ .<sup>2</sup> Decaying into  $2(\pi^+\pi^-)$ .<sup>3</sup> Taking into account  $f_4(2050)$ .<sup>4</sup> First solution, PWA is ambiguous.<sup>5</sup> T-matrix pole.<sup>6</sup> From solution B of amplitude analysis of data on  $\bar{p}p \rightarrow \pi\pi$ . See however KLOET 96 who fit  $\pi^+\pi^-$  only and find waves only up to  $J = 3$  to be important but not significantly resonant.<sup>7</sup> Cannot determine spin to be 2.



### $f_2(1950)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>464 ± 24 OUR AVERAGE</b>			
$380^{+120}_{-90}$	BAI	00A BES	$J/\psi \rightarrow \gamma(\pi^+\pi^-\pi^+\pi^-)$
$520 \pm 50$	<sup>8</sup> BARBERIS	00C	450 $pp \rightarrow pp4\pi$
$485 \pm 55$	<sup>9</sup> BARBERIS	00C	450 $pp \rightarrow pp4\pi$
$460 \pm 40$	BARBERIS	97B OMEG	450 $pp \rightarrow pp2(\pi^+\pi^-)$
$390 \pm 60$	ANTINORI	95 OMEG	300,450 $pp \rightarrow pp2(\pi^+\pi^-)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$441^{+27+28}_{-25-192}$	<sup>10</sup> UEHARA	09 BELL	10.6 $e^+e^- \rightarrow e^+e^-\pi^0\pi^0$
$450 \pm 50$	<sup>11</sup> BINON	05 GAMS	33 $\pi^-p \rightarrow \eta\eta n$
$297 \pm 12 \pm 6$	ABE	04 BELL	10.6 $e^+e^- \rightarrow e^+e^-K^+K^-$
$385 \pm 58$	<sup>12</sup> AMSLER	02 CBAR	0.9 $\bar{p}p \rightarrow \pi^0\eta\eta, \pi^0\pi^0\pi^0$
$495 \pm 35$	ANISOVICH	00J SPEC	
$500 \pm 100$	ANISOVICH	99B SPEC	1.35–1.94 $p\bar{p} \rightarrow \eta\eta\pi^0$
$\sim 100$	<sup>13</sup> OAKDEN	94 RVUE	0.36–1.55 $\bar{p}p \rightarrow \pi\pi$
$250 \pm 50$	<sup>14</sup> ASTON	91 LASS	11 $K^-p \rightarrow \Lambda K\bar{K}\pi\pi$

<sup>8</sup> Decaying into  $\pi^+\pi^-2\pi^0$ .

<sup>9</sup> Decaying into  $2(\pi^+\pi^-)$ .

<sup>10</sup> Taking into account  $f_4(2050)$ .

<sup>11</sup> First solution, PWA is ambiguous.

<sup>12</sup> T-matrix pole.

<sup>13</sup> From solution B of amplitude analysis of data on  $\bar{p}p \rightarrow \pi\pi$ . See however KLOET 96 who fit  $\pi^+\pi^-$  only and find waves only up to  $J = 3$  to be important but not significantly resonant.

<sup>14</sup> Cannot determine spin to be 2.

## $f_2(1950)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $K^*(892)\bar{K}^*(892)$	seen
$\Gamma_2$ $\pi\pi$	
$\Gamma_3$ $\pi^+\pi^-$	seen
$\Gamma_4$ $\pi^0\pi^0$	seen
$\Gamma_5$ $4\pi$	seen
$\Gamma_6$ $\pi^+\pi^-\pi^+\pi^-$	
$\Gamma_7$ $a_2(1320)\pi$	
$\Gamma_8$ $f_2(1270)\pi\pi$	
$\Gamma_9$ $\eta\eta$	seen
$\Gamma_{10}$ $K\bar{K}$	seen
$\Gamma_{11}$ $\gamma\gamma$	seen
$\Gamma_{12}$ $p\bar{p}$	seen

## $f_2(1950)$ $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_{10}\Gamma_{11}/\Gamma$
<u>VALUE (eV)</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>

• • • We do not use the following data for averages, fits, limits, etc. • • •

$122 \pm 4 \pm 26$       <sup>15</sup> ABE      04    BELL     $10.6 e^+e^- \rightarrow e^+e^-K^+K^-$

<sup>15</sup> Assuming spin 2.

$\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_2\Gamma_{11}/\Gamma$
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>

• • • We do not use the following data for averages, fits, limits, etc. • • •

$162^{+69+1137}_{-42-204}$       <sup>16</sup> UEHARA      09    BELL     $10.6 e^+e^- \rightarrow e^+e^-\pi^0\pi^0$

<sup>16</sup> Taking into account  $f_4(2050)$ .

## $f_2(1950)$ BRANCHING RATIOS

$\Gamma(K^*(892)\bar{K}^*(892))/\Gamma_{\text{total}}$	$\Gamma_1/\Gamma$
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>CHG</u> <u>COMMENT</u>
<b>seen</b>	ASTON      91    LASS    0    11 $K^-p \rightarrow \Lambda K\bar{K}\pi\pi$

$\Gamma(a_2(1320)\pi)/\Gamma_{\text{total}}$   $\Gamma_7/\Gamma$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
not seen	BARBERIS	00B	450 $pp \rightarrow p_f \eta \pi^+ \pi^- p_S$
not seen	BARBERIS	00C	450 $pp \rightarrow p_f 4\pi p_S$
possibly seen	BARBERIS	97B OMEG	450 $pp \rightarrow p p 2(\pi^+ \pi^-)$

$\Gamma(\eta\eta)/\Gamma(4\pi)$   $\Gamma_9/\Gamma_5$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$<5.0 \times 10^{-3}$	90	BARBERIS	00E 450 $pp \rightarrow p_f \eta \eta p_S$

$\Gamma(\eta\eta)/\Gamma(\pi^+\pi^-)$   $\Gamma_9/\Gamma_3$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.14±0.05</b>	AMSLER	02	CBAR 0.9 $\bar{p}p \rightarrow \pi^0 \eta \eta, \pi^0 \pi^0 \pi^0$

$\Gamma(p\bar{p})/\Gamma_{\text{total}}$   $\Gamma_{12}/\Gamma$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>seen</b>	111	ALEXANDER	10	CLEO $\psi(2S) \rightarrow \gamma p \bar{p}$

**$f_2(1950)$  REFERENCES**

ALEXANDER	10	PR D82 092002	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
UEHARA	09	PR D79 052009	S. Uehara <i>et al.</i>	(BELLE Collab.)
BINON	05	PAN 68 960	F. Binon <i>et al.</i>	
		Translated from YAF 68 998.		
ABE	04	EPJ C32 323	K. Abe <i>et al.</i>	(BELLE Collab.)
AMSLER	02	EPJ C23 29	C. Amsler <i>et al.</i>	
ANISOVICH	00J	PL B491 47	A.V. Anisovich <i>et al.</i>	(RAL, LOQM, PNPI+)
BAI	00A	PL B472 207	J.Z. Bai <i>et al.</i>	(BES Collab.)
BARBERIS	00B	PL B471 435	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BARBERIS	00C	PL B471 440	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BARBERIS	00E	PL B479 59	D. Barberis <i>et al.</i>	(WA 102 Collab.)
ANISOVICH	99B	PL B449 154	A.V. Anisovich <i>et al.</i>	
BARBERIS	97B	PL B413 217	D. Barberis <i>et al.</i>	(WA 102 Collab.)
KLOET	96	PR D53 6120	W.M. Kloet, F. Myhrer	(RUTG, NORD)
ANTINORI	95	PL B353 589	F. Antinori <i>et al.</i>	(ATHU, BARI, BIRM+) JP
OAKDEN	94	NP A574 731	M.N. Oakden, M.R. Pennington	(DURH)
ASTON	91	NPBPS B21 5	D. Aston <i>et al.</i>	(LASS Collab.)