



$$I(J^P) = 0(\frac{1}{2}^+) \quad \text{Status: } ***$$

In the quark model, a Λ_b^0 is an isospin-0 udb state. The lowest Λ_b^0 ought to have $J^P = 1/2^+$. None of I , J , or P have actually been measured.

Λ_b^0 MASS

$m_{\Lambda_b^0}$

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-----------------------|-----------|---|
| 5619.60 ± 0.17 OUR AVERAGE | | | | |
| 5619.62 ± 0.16 ± 0.13 | | ¹ AAIJ | 17AMLHCB | pp at 7, 8 TeV |
| 5619.30 ± 0.34 | | ² AAIJ | 14AA LHCB | pp at 7 TeV |
| 5620.15 ± 0.31 ± 0.47 | | ³ AALTONEN | 14B CDF | $p\bar{p}$ at 1.96 TeV |
| 5619.7 ± 0.7 ± 1.1 | | ³ AAD | 13U ATLS | pp at 7 TeV |
| 5621 ± 4 ± 3 | | ⁴ ABE | 97B CDF | $p\bar{p}$ at 1.8 TeV |
| 5668 ± 16 ± 8 | 4 | ⁵ ABREU | 96N DLPH | $e^+e^- \rightarrow Z$ |
| 5614 ± 21 ± 4 | 4 | ⁵ BUSKULIC | 96L ALEP | $e^+e^- \rightarrow Z$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| 5619.65 ± 0.17 ± 0.17 | | ⁶ AAIJ | 16Y LHCB | Repl. by AAIJ 17AM |
| 5619.44 ± 0.13 ± 0.38 | | ³ AAIJ | 13AV LHCB | Repl. by AAIJ 17AM |
| 5619.19 ± 0.70 ± 0.30 | | ³ AAIJ | 12E LHCB | Repl. by AAIJ 13AV |
| 5619.7 ± 1.2 ± 1.2 | | ⁷ ACOSTA | 06 CDF | Repl. by AALTONEN 14B |
| not seen | | ⁸ ABE | 93B CDF | Repl. by ABE 97B |
| 5640 ± 50 ± 30 | 16 | ⁹ ALBAJAR | 91E UA1 | $p\bar{p}$ 630 GeV |
| 5640 $^{+100}_{-210}$ | 52 | BARI | 91 SFM | $\Lambda_b^0 \rightarrow p D^0 \pi^-$ |
| 5650 $^{+150}_{-200}$ | 90 | BARI | 91 SFM | $\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^+ \pi^- \pi^-$ |

¹ Uses $\Lambda_b^0 \rightarrow \chi_{c1} p K^-$, $\Lambda_b^0 \rightarrow \chi_{c2} p K^-$, $\Lambda_b^0 \rightarrow J/\psi \Lambda$, $\Lambda_b^0 \rightarrow p \psi(2S) K^-$, $\Lambda_b^0 \rightarrow p J/\psi \pi^+ \pi^- K^-$, and $\Lambda_b^0 \rightarrow p J/\psi K^-$ decays.

² Uses exclusively reconstructed final states $\Lambda_b^0 \rightarrow \Lambda_c^+ D_s^-$, $\Lambda_c^+ D^-$ and $\bar{B}^0 \rightarrow D^+ D_s^-$ decays. The uncertainty includes both statistical and systematic contributions.

³ Uses $\Lambda_b^0 \rightarrow J/\psi \Lambda$ fully reconstructed decays.

⁴ ABE 97B observed 38 events with a background of 18 ± 1.6 events in the mass range 5.60–5.65 GeV/ c^2 , a significance of > 3.4 standard deviations.

⁵ Uses 4 fully reconstructed Λ_b events.

⁶ Uses $\Lambda_b^0 \rightarrow p \psi(2S) K^-$, $\Lambda_b^0 \rightarrow p J/\psi \pi^+ \pi^- K^-$, and $\Lambda_b^0 \rightarrow p J/\psi K^-$ decays.

⁷ Uses exclusively reconstructed final states containing a $J/\psi \rightarrow \mu^+ \mu^-$ decays.

⁸ ABE 93B states that, based on the signal claimed by ALBAJAR 91E, CDF should have found $30 \pm 23 \Lambda_b^0 \rightarrow J/\psi(1S) \Lambda$ events. Instead, CDF found not more than 2 events.

⁹ ALBAJAR 91E claims 16 ± 5 events above a background of 9 ± 1 events, a significance of about 5 standard deviations.

$m_{\Lambda_b^0} - m_{B^0}$

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|----------------------|---------------------|--------|------------------------|
| 339.2±1.4±0.1 | ¹ ACOSTA | 06 CDF | $p\bar{p}$ at 1.96 TeV |

¹ Uses exclusively reconstructed final states containing $J/\psi \rightarrow \mu^+ \mu^-$ decays.

 $m_{\Lambda_b^0} - m_{B^+}$

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|--------------------------------|-------------------|-----------|---------------|
| 339.72±0.28 OUR AVERAGE | | | |
| 339.72±0.24±0.18 | ¹ AAIJ | 14AA LHCb | pp at 7 TeV |
| 339.71±0.71±0.09 | ² AAIJ | 12E LHCb | pp at 7 TeV |

¹ Uses exclusively reconstructed final states $\Lambda_b^0 \rightarrow \Lambda_c^+ D_s^-$, $\Lambda_c^+ D^-$ and $\bar{B}^0 \rightarrow D^+ D_s^-$ decays.

² Uses exclusively reconstructed final states containing $J/\psi \rightarrow \mu^+ \mu^-$ decays.

 Λ_b^0 MEAN LIFE

See b -baryon Admixture section for data on b -baryon mean life average over species of b -baryon particles.

"OUR EVALUATION" is an average using rescaled values of the data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFLAV) and are described at <http://www.slac.stanford.edu/xorg/hflav/>. The averaging/rescaling procedure takes into account correlations between the measurements and asymmetric lifetime errors.

| VALUE (10^{-12} s) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|--------------|-----------------|-------|---------------------------------|
| 1.471±0.009 OUR EVALUATION | | | | |
| 1.477±0.027±0.009 | ¹ | SIRUNYAN 18BY | CMS | pp at 8 TeV |
| 1.415±0.027±0.006 | ² | AAIJ 14E | LHCb | pp at 7 TeV |
| 1.479±0.009±0.010 | ³ | AAIJ 14U | LHCb | pp at 7, 8 TeV |
| 1.565±0.035±0.020 | ² | AALTONEN 14B | CDF | $p\bar{p}$ at 1.96 TeV |
| 1.449±0.036±0.017 | ² | AAD 13U | ATLAS | pp at 7 TeV |
| 1.503±0.052±0.031 | ² | CHATRCHYAN 13AC | CMS | pp at 7 TeV |
| 1.303±0.075±0.035 | ² | ABAZOV 12U | D0 | $p\bar{p}$ at 1.96 TeV |
| 1.401±0.046±0.035 | ⁴ | AALTONEN 10B | CDF | $p\bar{p}$ at 1.96 TeV |
| 1.27 ^{+0.35} _{-0.29} ±0.09 | | ABREU 95S | DLPH | Excess $p\mu^-$, decay lengths |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|---|--------------|---------------|------|-------------------------|
| 1.482±0.018±0.012 | ⁵ | AAIJ 13BB | LHCb | Repl. by AAIJ 14U |
| 1.537±0.045±0.014 | ² | AALTONEN 11 | CDF | Repl. by AALTONEN 14B |
| 1.218 ^{+0.130} _{-0.115} ±0.042 | ² | ABAZOV 07S | D0 | Repl. by ABAZOV 12U |
| 1.290 ^{+0.119} _{-0.110} ^{+0.087} _{-0.091} | ⁶ | ABAZOV 07U | D0 | $p\bar{p}$ at 1.96 TeV |
| 1.593 ^{+0.083} _{-0.078} ±0.033 | ² | ABULENCIA 07A | CDF | Repl. by AALTONEN 11 |
| 1.22 ^{+0.22} _{-0.18} ±0.04 | ² | ABAZOV 05C | D0 | Repl. by ABAZOV 07S |
| 1.11 ^{+0.19} _{-0.18} ±0.05 | ⁷ | ABREU 99W | DLPH | $e^+ e^- \rightarrow Z$ |

| | | | | | | | |
|------|--|--|--------------|------------|-----|------|--------------------------|
| 1.29 | $\begin{smallmatrix} +0.24 \\ -0.22 \end{smallmatrix}$ | ± 0.06 | ⁷ | ACKERSTAFF | 98G | OPAL | $e^+e^- \rightarrow Z$ |
| 1.21 | ± 0.11 | | ⁷ | BARATE | 98D | ALEP | $e^+e^- \rightarrow Z$ |
| 1.32 | ± 0.15 | ± 0.07 | ⁸ | ABE | 96M | CDF | $\rho\bar{p}$ at 1.8 TeV |
| 1.19 | $\begin{smallmatrix} +0.21 \\ -0.18 \end{smallmatrix}$ | $\begin{smallmatrix} +0.07 \\ -0.08 \end{smallmatrix}$ | | ABREU | 96D | DLPH | Repl. by ABREU 99W |
| 1.14 | $\begin{smallmatrix} +0.22 \\ -0.19 \end{smallmatrix}$ | ± 0.07 | 69 | AKERS | 95K | OPAL | Repl. by ACKERSTAFF 98G |
| 1.02 | $\begin{smallmatrix} +0.23 \\ -0.18 \end{smallmatrix}$ | ± 0.06 | 44 | BUSKULIC | 95L | ALEP | Repl. by BARATE 98D |

¹ Measured using $\Lambda_b^0 \rightarrow J/\psi \Lambda$ decays.

² Measured mean life using fully reconstructed $\Lambda_b^0 \rightarrow J/\psi \Lambda$ decays.

³ Used $\Lambda_b^0 \rightarrow J/\psi p K^-$ decays.

⁴ Measured mean life using fully reconstructed $\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-$ decays.

⁵ Measured the lifetime ratio of decays $\Lambda_b^0 \rightarrow J/\psi p K^-$ to $B^0 \rightarrow J/\psi \pi^+ K^-$ to be $0.976 \pm 0.012 \pm 0.006$ with $\tau_{B^0} = 1.519 \pm 0.007$ ps.

⁶ Measured using semileptonic decays $\Lambda_b^0 \rightarrow \Lambda_c^+ \mu \nu X$ and $\Lambda_c^+ \rightarrow K_S^0 p$.

⁷ Measured using $\Lambda_c \ell^-$ and $\Lambda \ell^+ \ell^-$.

⁸ Excess $\Lambda_c \ell^-$, decay lengths.

$\tau_{\Lambda_b^0}/\tau_{\Lambda_b^+}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|------------------------------|-------------------|----------|---------------|
| 0.940 ± 0.035 ± 0.006 | ¹ AAIJ | 14E LHCB | pp at 7 TeV |

¹ Measured using $\Lambda_b^0 \rightarrow J/\psi \Lambda$ decays.

$\tau_{\Lambda_b^0}/\tau_{B^0}$ MEAN LIFE RATIO

$\tau_{\Lambda_b^0}/\tau_{B^0}$ (direct measurements)

“OUR EVALUATION” has been obtained by the Heavy Flavor Averaging Group (HFLAV) by including both B^0 and B^+ decays.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------------------------------------|-------------|------|---------|
| 0.964 ± 0.007 OUR EVALUATION | | | |


0.970 ± 0.009 OUR AVERAGE Error includes scale factor of 1.4. See the ideogram below.

| | | | |
|-----------------------|-----------------------|----------|---------------------------|
| 0.978 ± 0.018 ± 0.006 | ¹ SIRUNYAN | 18BY CMS | pp at 8 TeV |
| 0.929 ± 0.018 ± 0.004 | ¹ AAIJ | 14E LHCB | pp at 7 TeV |
| 0.974 ± 0.006 ± 0.004 | ² AAIJ | 14U LHCB | pp at 7, 8 TeV |
| 0.960 ± 0.025 ± 0.016 | ³ AAD | 13U ATLS | pp at 7 TeV |
| 0.864 ± 0.052 ± 0.033 | ^{4,5} ABAZOV | 12U D0 | $\rho\bar{p}$ at 1.96 TeV |
| 1.020 ± 0.030 ± 0.008 | ⁴ AALTONEN | 11 CDF | $\rho\bar{p}$ at 1.96 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|--|------------------------|-----------|----------------------|
| 0.976 ± 0.012 ± 0.006 | ⁶ AAIJ | 13BB LHCB | Repl. by AAIJ 14U |
| 0.811 $\begin{smallmatrix} +0.096 \\ -0.087 \end{smallmatrix}$ ± 0.034 | ^{4,5} ABAZOV | 07S D0 | Repl. by ABAZOV 12U |
| 1.041 ± 0.057 | ⁷ ABULENCIA | 07A CDF | Repl. by AALTONEN 11 |
| 0.87 $\begin{smallmatrix} +0.17 \\ -0.14 \end{smallmatrix}$ ± 0.03 | ⁷ ABAZOV | 05C D0 | Repl. by ABAZOV 07S |

- 1 Measured using $\Lambda_b^0 \rightarrow J/\psi \Lambda$ and $B^0 \rightarrow J/\psi K^*(892)^0$ decays.
- 2 Used $\Lambda_b^0 \rightarrow J/\psi p K^-$ and $B^0 \rightarrow J/\psi K^*(892)^0$ decays.
- 3 Measured with $\Lambda_b^0 \rightarrow J/\psi(\mu^+ \mu^-) \Lambda^0(p\pi^-)$ decays.
- 4 Uses fully reconstructed $\Lambda_b \rightarrow J/\psi \Lambda$ decays.
- 5 Uses $B^0 \rightarrow J/\psi K_S^0$ decays for denominator.
- 6 Measures $1/\tau_{\Lambda_b^0} - 1/\tau_{B^0}$ and uses $\tau_{B^0} = 1.519 \pm 0.007$ ps to extract lifetime ratio.
- 7 Measured mean life ratio using fully reconstructed decays.



Λ_b^0 DECAY MODES

The branching fractions $B(b\text{-baryon} \rightarrow \Lambda \ell^- \bar{\nu}_\ell \text{ anything})$ and $B(\Lambda_b^0 \rightarrow \Lambda_c^+ \ell^- \bar{\nu}_\ell \text{ anything})$ are not pure measurements because the underlying measured products of these with $B(b \rightarrow b\text{-baryon})$ were used to determine $B(b \rightarrow b\text{-baryon})$, as described in the note "Production and Decay of b -Flavored Hadrons."

For inclusive branching fractions, e.g., $\Lambda_b \rightarrow \bar{\Lambda}_c \text{ anything}$, the values usually are multiplicities, not branching fractions. They can be greater than one.

| Mode | Fraction (Γ_i/Γ) | Scale factor/ Confidence level |
|---|--------------------------------------|-----------------------------------|
| Γ_1 $J/\psi(1S)\Lambda \times B(b \rightarrow \Lambda_b^0)$ | $(5.8 \pm 0.8) \times 10^{-5}$ | |
| Γ_2 $J/\psi(1S)\Lambda$ | | |
| Γ_3 $\psi(2S)\Lambda$ | | |
| Γ_4 $pD^0\pi^-$ | $(6.3 \pm 0.7) \times 10^{-4}$ | |
| Γ_5 $\Lambda_c(2860)^+\pi^-, \Lambda_c^+ \rightarrow D^0 p$ | | |
| Γ_6 $\Lambda_c(2880)^+\pi^-, \Lambda_c^+ \rightarrow D^0 p$ | | |
| Γ_7 $\Lambda_c(2940)^+\pi^-, \Lambda_c^+ \rightarrow D^0 p$ | | |
| Γ_8 pD^0K^- | $(4.6 \pm 0.8) \times 10^{-5}$ | |
| Γ_9 $pJ/\psi\pi^-$ | $(2.6^{+0.5}_{-0.4}) \times 10^{-5}$ | |
| Γ_{10} $p\pi^- J/\psi, J/\psi \rightarrow \mu^+\mu^-$ | $(1.6 \pm 0.8) \times 10^{-6}$ | |
| Γ_{11} $pJ/\psi K^-$ | $(3.2^{+0.6}_{-0.5}) \times 10^{-4}$ | |
| Γ_{12} $P_c(4380)^+K^-, P_c \rightarrow pJ/\psi$ [a] | $(2.7 \pm 1.4) \times 10^{-5}$ | |
| Γ_{13} $P_c(4450)^+K^-, P_c \rightarrow pJ/\psi$ [a] | $(1.3 \pm 0.4) \times 10^{-5}$ | |
| Γ_{14} $\chi_{c1}(1P)pK^-$ | $(7.6^{+1.5}_{-1.3}) \times 10^{-5}$ | |
| Γ_{15} $\chi_{c2}(1P)pK^-$ | $(7.9^{+1.6}_{-1.4}) \times 10^{-5}$ | |
| Γ_{16} $pJ/\psi(1S)\pi^+\pi^-K^-$ | $(6.6^{+1.3}_{-1.1}) \times 10^{-5}$ | |
| Γ_{17} $p\psi(2S)K^-$ | $(6.6^{+1.2}_{-1.0}) \times 10^{-5}$ | |
| Γ_{18} $\psi(2S)p\pi^-$ | $(7.5^{+1.6}_{-1.4}) \times 10^{-6}$ | |
| Γ_{19} $p\bar{K}^0\pi^-$ | $(1.3 \pm 0.4) \times 10^{-5}$ | |
| Γ_{20} pK^0K^- | $< 3.5 \times 10^{-6}$ | CL=90% |
| Γ_{21} $\Lambda_c^+\pi^-$ | $(4.9 \pm 0.4) \times 10^{-3}$ | S=1.2 |
| Γ_{22} $\Lambda_c^+K^-$ | $(3.59 \pm 0.30) \times 10^{-4}$ | S=1.2 |
| Γ_{23} $\Lambda_c^+ a_1(1260)^-$ | seen | |
| Γ_{24} $\Lambda_c^+ D^-$ | $(4.6 \pm 0.6) \times 10^{-4}$ | |
| Γ_{25} $\Lambda_c^+ D_s^-$ | $(1.10 \pm 0.10) \%$ | |
| Γ_{26} $\Lambda_c^+\pi^+\pi^-\pi^-$ | $(7.7 \pm 1.1) \times 10^{-3}$ | S=1.1 |
| Γ_{27} $\Lambda_c(2595)^+\pi^-, \Lambda_c(2595)^+ \rightarrow \Lambda_c^+\pi^+\pi^-$ | $(3.4 \pm 1.5) \times 10^{-4}$ | |
| Γ_{28} $\Lambda_c(2625)^+\pi^-, \Lambda_c(2625)^+ \rightarrow \Lambda_c^+\pi^+\pi^-$ | $(3.3 \pm 1.3) \times 10^{-4}$ | |
| Γ_{29} $\Sigma_c(2455)^0\pi^+\pi^-, \Sigma_c^0 \rightarrow \Lambda_c^+\pi^-$ | $(5.7 \pm 2.2) \times 10^{-4}$ | |
| Γ_{30} $\Sigma_c(2455)^{++}\pi^-\pi^-, \Sigma_c^{++} \rightarrow \Lambda_c^+\pi^+$ | $(3.2 \pm 1.6) \times 10^{-4}$ | |

| | | | | |
|---------------|---|-----|--|--------|
| Γ_{31} | $\Lambda_c^+ p \bar{p} \pi^-$ | | $(2.65 \pm 0.29) \times 10^{-4}$ | |
| Γ_{32} | $\Sigma_c(2455)^0 p \bar{p}, \Sigma_c(2455)^0 \rightarrow$ $\Lambda_c^+ \pi^-$ | | $(2.4 \pm 0.5) \times 10^{-5}$ | |
| Γ_{33} | $\Sigma_c(2520)^0 p \bar{p}, \Sigma_c(2520)^0 \rightarrow$ $\Lambda_c^+ \pi^-$ | | $(3.2 \pm 0.7) \times 10^{-5}$ | |
| Γ_{34} | $\Lambda K^0 2\pi^+ 2\pi^-$ | | | |
| Γ_{35} | $\Lambda_c^+ \ell^- \bar{\nu}_\ell$ anything | [b] | $(10.4 \pm 2.2) \%$ | |
| Γ_{36} | $\Lambda_c^+ \ell^- \bar{\nu}_\ell$ | | $(6.2^{+1.4}_{-1.3}) \%$ | |
| Γ_{37} | $\Lambda_c^+ \pi^+ \pi^- \ell^- \bar{\nu}_\ell$ | | $(5.6 \pm 3.1) \%$ | |
| Γ_{38} | $\Lambda_c(2595)^+ \ell^- \bar{\nu}_\ell$ | | $(7.9^{+4.0}_{-3.5}) \times 10^{-3}$ | |
| Γ_{39} | $\Lambda_c(2625)^+ \ell^- \bar{\nu}_\ell$ | | $(1.3^{+0.6}_{-0.5}) \%$ | |
| Γ_{40} | $\Sigma_c(2455)^0 \pi^+ \ell^- \bar{\nu}_\ell$ | | | |
| Γ_{41} | $\Sigma_c(2455)^{++} \pi^- \ell^- \bar{\nu}_\ell$ | | | |
| Γ_{42} | $p h^-$ | [c] | $< 2.3 \times 10^{-5}$ | CL=90% |
| Γ_{43} | $p \pi^-$ | | $(4.3 \pm 0.8) \times 10^{-6}$ | |
| Γ_{44} | $p K^-$ | | $(5.1 \pm 0.9) \times 10^{-6}$ | |
| Γ_{45} | $p D_s^-$ | | $< 4.8 \times 10^{-4}$ | CL=90% |
| Γ_{46} | $p \mu^- \bar{\nu}_\mu$ | | $(4.1 \pm 1.0) \times 10^{-4}$ | |
| Γ_{47} | $\Lambda \mu^+ \mu^-$ | | $(1.08 \pm 0.28) \times 10^{-6}$ | |
| Γ_{48} | $p \pi^- \mu^+ \mu^-$ | | $(6.9 \pm 2.5) \times 10^{-8}$ | |
| Γ_{49} | $\Lambda \gamma$ | | $< 1.3 \times 10^{-3}$ | CL=90% |
| Γ_{50} | $\Lambda \eta$ | | $(9^{+7}_{-5}) \times 10^{-6}$ | |
| Γ_{51} | $\Lambda \eta'(958)$ | | $< 3.1 \times 10^{-6}$ | CL=90% |
| Γ_{52} | $\Lambda \pi^+ \pi^-$ | | $(4.7 \pm 1.9) \times 10^{-6}$ | |
| Γ_{53} | $\Lambda K^+ \pi^-$ | | $(5.7 \pm 1.3) \times 10^{-6}$ | |
| Γ_{54} | $\Lambda K^+ K^-$ | | $(1.62 \pm 0.23) \times 10^{-5}$ | |
| Γ_{55} | $\Lambda \phi$ | | $(9.3 \pm 2.5) \times 10^{-6}$ | |
| Γ_{56} | $p \pi^- \pi^+ \pi^-$ | | $(2.11 \pm 0.23) \times 10^{-5}$ | |
| Γ_{57} | $p K^- K^+ \pi^-$ | | $(4.1 \pm 0.6) \times 10^{-6}$ | |
| Γ_{58} | $p K^- \pi^+ \pi^-$ | | $(5.1 \pm 0.5) \times 10^{-5}$ | |
| Γ_{59} | $p K^- K^+ K^-$ | | $(1.27 \pm 0.14) \times 10^{-5}$ | |

[a] P_c^+ is a pentaquark-charmonium state.

[b] Not a pure measurement. See note at head of Λ_b^0 Decay Modes.

[c] Here h^- means π^- or K^- .

CONSTRAINED FIT INFORMATION

An overall fit to 10 branching ratios uses 12 measurements and one constraint to determine 7 parameters. The overall fit has a $\chi^2 = 10.7$ for 6 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

| | | | | | |
|----------|----------|----------|----------|----------|----------|
| x_{22} | 94 | | | | |
| x_{26} | 50 | 47 | | | |
| x_{36} | 14 | 14 | 7 | | |
| x_{43} | 0 | 0 | 0 | 0 | |
| x_{44} | 0 | 0 | 0 | 0 | 83 |
| | x_{21} | x_{22} | x_{26} | x_{36} | x_{43} |

Λ_b^0 BRANCHING RATIOS

$\Gamma(J/\psi(1S)\Lambda \times B(b \rightarrow \Lambda_b^0)) / \Gamma_{\text{total}}$ Γ_1/Γ

| VALUE (units 10^{-5}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|

5.8 ± 0.8 OUR AVERAGE

6.01 ± 0.60 ± 0.58 ± 0.28 ¹ ABAZOV 110 DO $p\bar{p}$ at 1.96 TeV

4.7 ± 2.3 ± 0.2 ² ABE 97B CDF $p\bar{p}$ at 1.8 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

180 ± 60 ± 90 16 ALBAJAR 91E UA1 $p\bar{p}$ at 630 GeV

¹ ABAZOV 110 uses $B(B^0 \rightarrow J/\psi K_S^0) \times B(b \rightarrow B^0) = (1.74 \pm 0.08) \times 10^{-4}$ to obtain the result. The $(\pm 0.08) \times 10^{-4}$ uncertainty of this product is listed as the last uncertainty of the measurement, $(\pm 0.28) \times 10^{-5}$.

² ABE 97B reports $[B(\Lambda_b^0 \rightarrow J/\psi \Lambda) \times B(b \rightarrow \Lambda_b^0)] / [B(B^0 \rightarrow J/\psi K_S^0) \times B(b \rightarrow B^0)] = 0.27 \pm 0.12 \pm 0.05$. We multiply by our best value $B(B^0 \rightarrow J/\psi K_S^0) \times B(b \rightarrow B^0) = (1.74 \pm 0.08) \times 10^{-4}$. Our first error is their experiment error and our second error is the systematic error from using our best value.

$\Gamma(\psi(2S)\Lambda) / \Gamma(J/\psi(1S)\Lambda)$ Γ_3/Γ_2

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

0.50 ± 0.03 ± 0.02 ¹ AAD 15CH ATLS pp at 8 TeV

¹ AAD 15CH uses $B(J/\psi \rightarrow \mu^+ \mu^-) = (5.961 \pm 0.033) \times 10^{-2}$ (PDG 14). And $B(\psi(2S) \rightarrow \mu^+ \mu^-) = (7.89 \pm 0.17) \times 10^{-3}$ (PDG 14) is used assuming lepton universality.

$\Gamma(pD^0\pi^-) / \Gamma_{\text{total}}$ Γ_4/Γ

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------|------|-------------|------|---------|
|-------|------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen 52 BARI 91 SFM $D^0 \rightarrow K^- \pi^+$

seen BASILE 81 SFM $D^0 \rightarrow K^- \pi^+$

$\Gamma(\Lambda_c(2860)^+ \pi^-, \Lambda_c^+ \rightarrow D^0 p) / \Gamma(\Lambda_c(2880)^+ \pi^-, \Lambda_c^+ \rightarrow D^0 p) \quad \Gamma_5/\Gamma_6$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|----------------------------------|-------------|------|-----------------------|
| $4.54^{+0.51+0.21}_{-0.39-0.59}$ | AAIJ | 17S | LHCB pp at 7, 8 TeV |

$\Gamma(\Lambda_c(2940)^+ \pi^-, \Lambda_c^+ \rightarrow D^0 p) / \Gamma(\Lambda_c(2880)^+ \pi^-, \Lambda_c^+ \rightarrow D^0 p) \quad \Gamma_7/\Gamma_6$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|----------------------------------|-------------|------|-----------------------|
| $0.83^{+0.31+0.18}_{-0.10-0.43}$ | AAIJ | 17S | LHCB pp at 7, 8 TeV |

$\Gamma(p D^0 K^-) / \Gamma(p D^0 \pi^-) \quad \Gamma_8/\Gamma_4$

| VALUE (units 10^{-2}) | DOCUMENT ID | TECN | COMMENT |
|-----------------------------|-------------|------|--------------------|
| $7.3 \pm 0.8^{+0.5}_{-0.6}$ | AAIJ | 14H | LHCB pp at 7 TeV |

$\Gamma(p J/\psi \pi^-) / \Gamma(p J/\psi K^-) \quad \Gamma_9/\Gamma_{11}$

| VALUE (units 10^{-2}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|------|-----------------------|
| $8.24 \pm 0.25 \pm 0.42$ | AAIJ | 14K | LHCB pp at 7, 8 TeV |

$\Gamma(p J/\psi K^-) / \Gamma_{\text{total}} \quad \Gamma_{11}/\Gamma$

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|---------------------------------|-------------------|------|-----------------------|
| $3.17 \pm 0.04^{+0.57}_{-0.45}$ | ¹ AAIJ | 16A | LHCB pp at 7, 8 TeV |

¹ AAIJ 16A reported the measurement of $(3.17 \pm 0.04 \pm 0.07 \pm 0.34^{+0.45}_{-0.28}) \times 10^{-4}$ where the first uncertainty is statistical, the second is systematic, the third is due to the branching fraction of $B^0 \rightarrow J/\psi K^*(892)^0$, and the fourth is due to the knowledge of f_{Λ_b}/f_d . We combined in quadrature second to fourth uncertainties to a total systematic uncertainty.

$\Gamma(P_c(4380)^+ K^-, P_c \rightarrow p J/\psi) / \Gamma_{\text{total}} \quad \Gamma_{12}/\Gamma$

P_c^+ is a pentaquark-charmonium state.

| VALUE (units 10^{-5}) | DOCUMENT ID | TECN | COMMENT |
|---------------------------------|-------------------|------|-----------------------|
| $2.66 \pm 0.22^{+1.41}_{-1.38}$ | ¹ AAIJ | 16A | LHCB pp at 7, 8 TeV |

¹ AAIJ 16 total systematic includes the uncertainties on $f(P_c^+)$ and $B(\Lambda_b \rightarrow p J/\psi K^-)$.

$\Gamma(P_c(4450)^+ K^-, P_c \rightarrow p J/\psi) / \Gamma_{\text{total}} \quad \Gamma_{13}/\Gamma$

P_c^+ is a pentaquark-charmonium state.

| VALUE (units 10^{-5}) | DOCUMENT ID | TECN | COMMENT |
|---------------------------------|-------------------|------|-----------------------|
| $1.30 \pm 0.16^{+0.42}_{-0.39}$ | ¹ AAIJ | 16A | LHCB pp at 7, 8 TeV |

¹ AAIJ 16 total systematic includes the uncertainties on $f(P_c^+)$ and $B(\Lambda_b \rightarrow p J/\psi K^-)$.

$\Gamma(\chi_{c1}(1P)\rho K^-)/\Gamma(\rho J/\psi K^-)$ Γ_{14}/Γ_{11}

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------------|----------|------------------|
| 0.239±0.019±0.007 | ¹ AAIJ | 17AMLHCB | pp at 7, 8 TeV |

¹ AAIJ 17AM reports $0.242 \pm 0.014 \pm 0.016$ from a measurement of $[\Gamma(\Lambda_b^0 \rightarrow \chi_{c1}(1P)\rho K^-)/\Gamma(\Lambda_b^0 \rightarrow \rho J/\psi K^-)] \times [B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S))]$ assuming $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (33.9 \pm 1.2) \times 10^{-2}$, which we rescale to our best value $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (34.3 \pm 1.0) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\chi_{c2}(1P)\rho K^-)/\Gamma(\rho J/\psi K^-)$ Γ_{15}/Γ_{11}

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------------|----------|------------------|
| 0.250±0.025±0.007 | ¹ AAIJ | 17AMLHCB | pp at 7, 8 TeV |

¹ AAIJ 17AM reports $0.248 \pm 0.02 \pm 0.017$ from a measurement of $[\Gamma(\Lambda_b^0 \rightarrow \chi_{c2}(1P)\rho K^-)/\Gamma(\Lambda_b^0 \rightarrow \rho J/\psi K^-)] \times [B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S))]$ assuming $B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) = (19.2 \pm 0.7) \times 10^{-2}$, which we rescale to our best value $B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) = (19.0 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\rho J/\psi(1S)\pi^+\pi^- K^-)/\Gamma(\rho J/\psi K^-)$ Γ_{16}/Γ_{11}

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-----------------------------|-------------------|----------|------------------|
| 0.2086±0.0096±0.0134 | ¹ AAIJ | 16Y LHCB | pp at 7, 8 TeV |

¹ Excludes $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$.

$\Gamma(\rho\psi(2S)K^-)/\Gamma(\rho J/\psi K^-)$ Γ_{17}/Γ_{11}

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-----------------------------|-------------------|----------|------------------|
| 0.2070±0.0076±0.0059 | ¹ AAIJ | 16Y LHCB | pp at 7, 8 TeV |

¹ AAIJ 16Y reports a measurement of $0.2070 \pm 0.0076 \pm 0.0046 \pm 0.0037$ where the third uncertainty is due to the knowledge of J/ψ and $\psi(2S)$ branching fractions. We have combined both systematic uncertainties in quadrature.

$\Gamma(\psi(2S)\rho\pi^-)/\Gamma(\rho\psi(2S)K^-)$ Γ_{18}/Γ_{17}

| VALUE (%) | DOCUMENT ID | TECN | COMMENT |
|---------------------|-------------|-----------|----------------------|
| 11.4±1.3±0.2 | AAIJ | 18AF LHCB | pp at 7, 8, 13 TeV |

$\Gamma(\rho\bar{K}^0\pi^-)/\Gamma_{\text{total}}$ Γ_{19}/Γ

| VALUE (units 10^{-5}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------------|----------|---------------|
| 1.26±0.19±0.36 | ¹ AAIJ | 14Q LHCB | pp at 7 TeV |

¹ Used the normalizing mode branching fraction value of $B(B^0 \rightarrow K^0 \pi^+ \pi^-) = (4.96 \pm 0.20) \times 10^{-5}$.

$\Gamma(\rho K^0 K^-)/\Gamma_{\text{total}}$ Γ_{20}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|----------------------------------|-----|-------------|----------|---------------|
| <3.5 × 10⁻⁶ | 90 | AAIJ | 14Q LHCB | pp at 7 TeV |

$\Gamma(\Lambda_c^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{21}/Γ

| VALUE (units 10^{-3}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|

4.9 ± 0.4 OUR FIT Error includes scale factor of 1.2.

4.9 ± 0.5 OUR AVERAGE Error includes scale factor of 1.5.

| | | | | |
|---------------------------------|--|------------------------|-----|----------------------------|
| $4.57^{+0.31}_{-0.30} \pm 0.23$ | | ¹ AAIJ | 14I | LHCB pp at 7 TeV |
| $5.97 \pm 0.28 \pm 0.81$ | | ² AAIJ | 14Q | LHCB pp at 7 TeV |
| $8.8 \pm 2.8 \pm 1.5$ | | ³ ABULENCIA | 07B | CDF $p\bar{p}$ at 1.96 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|------|---|----------|-----|---|
| seen | 3 | ABREU | 96N | DLPH $\Lambda_c^+ \rightarrow pK^- \pi^+$ |
| seen | 4 | BUSKULIC | 96L | ALEP $\Lambda_c^+ \rightarrow pK^- \pi^+$, $p\bar{K}^0, \Lambda\pi^+ \pi^+ \pi^-$ |

¹ AAIJ 14I reports $(4.30 \pm 0.03^{+0.12}_{-0.11} \pm 0.26 \pm 0.21) \times 10^{-3}$ from a measurement of $[\Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-)/\Gamma_{\text{total}}] \times [B(B^0 \rightarrow D^- \pi^+)]$ assuming $B(B^0 \rightarrow D^- \pi^+) = (2.68 \pm 0.13) \times 10^{-3}$, which we rescale to our best value $B(B^0 \rightarrow D^- \pi^+) = (2.52 \pm 0.13) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Uses information on f_{baryon}/f_d from measurement in semileptonic decays by the same authors.

² Obtained using the branching fraction of $\Lambda_c^+ \rightarrow pK^- \pi^+$ decay.

³ The result is obtained from $(f_{\text{baryon}}/f_d) (B(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-)/B(\bar{B}^0 \rightarrow D^+ \pi^-)) = 0.82 \pm 0.08 \pm 0.11 \pm 0.22$, assuming $f_{\text{baryon}}/f_d = 0.25 \pm 0.04$ and $B(\bar{B}^0 \rightarrow D^+ \pi^-) = (2.68 \pm 0.13) \times 10^{-3}$.

$\Gamma(pD^0 \pi^-)/\Gamma(\Lambda_c^+ \pi^-)$ Γ_4/Γ_{21}

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

0.128 ± 0.007^{+0.006}_{-0.007} ¹ AAIJ 14H LHCB pp at 7 TeV

¹ AAIJ 14H reports $[\Gamma(\Lambda_b^0 \rightarrow pD^0 \pi^-)/\Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-)] \times [B(D^0 \rightarrow K^- \pi^+)] / [B(\Lambda_c^+ \rightarrow pK^- \pi^+)] = (8.06 \pm 0.23 \pm 0.35) \times 10^{-2}$ which we multiply or divide by our best values $B(D^0 \rightarrow K^- \pi^+) = (3.950 \pm 0.031) \times 10^{-2}$, $B(\Lambda_c^+ \rightarrow pK^- \pi^+) = (6.28 \pm 0.32) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

$\Gamma(\Lambda_c^+ K^-)/\Gamma_{\text{total}}$ Γ_{22}/Γ

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|------|---------|
|--------------------------|-------------|------|---------|

3.59 ± 0.30 OUR FIT Error includes scale factor of 1.2.

3.55 ± 0.44 ± 0.50 ¹ AAIJ 14Q LHCB pp at 7 TeV

¹ Obtained using the branching fraction of $\Lambda_c^+ \rightarrow pK^- \pi^+$ decay.

$\Gamma(\Lambda_c^+ K^-)/\Gamma(\Lambda_c^+ \pi^-)$ Γ_{22}/Γ_{21}

| VALUE (units 10^{-2}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|------|---------|
|--------------------------|-------------|------|---------|

7.31 ± 0.22 OUR FIT

7.31 ± 0.16 ± 0.16 AAIJ 14H LHCB pp at 7 TeV

| $\Gamma(\Lambda_c^+ a_1(1260)^-)/\Gamma_{\text{total}}$ | | | | | Γ_{23}/Γ |
|---|------|-------------|------|---------|---|
| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT | |
| seen | 1 | ABREU | 96N | DLPH | $\Lambda_c^+ \rightarrow p K^- \pi^+, a_1^- \rightarrow \rho^0 \pi^- \rightarrow \pi^+ \pi^- \pi^-$ |

| $\Gamma(\Lambda_c^+ D_s^-)/\Gamma_{\text{total}}$ | | | | | Γ_{25}/Γ |
|--|--|-------------------|------|---------|----------------------|
| VALUE (units 10^{-2}) | | DOCUMENT ID | TECN | COMMENT | |
| 1.1±0.1 | | ¹ AAIJ | 14AA | LHCB | pp at 7 TeV |
| ¹ Uses $B(\bar{B}^0 \rightarrow D^+ D_s^-) = (7.2 \pm 0.8) \times 10^{-3}$ and their measured $B(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-)/B(\bar{B}^0 \rightarrow D^+ \pi^-)$ values. | | | | | |

| $\Gamma(\Lambda_c^+ D^-)/\Gamma(\Lambda_c^+ D_s^-)$ | | | | | Γ_{24}/Γ_{25} |
|---|--|-------------|------|---------|---------------------------|
| VALUE | | DOCUMENT ID | TECN | COMMENT | |
| 0.042±0.003±0.003 | | AAIJ | 14AA | LHCB | pp at 7 TeV |

| $\Gamma(\Lambda_c^+ \pi^+ \pi^- \pi^-)/\Gamma_{\text{total}}$ | | | | | Γ_{26}/Γ |
|---|------|-------------------------------------|------|---------|------------------------|
| VALUE (units 10^{-3}) | EVTS | DOCUMENT ID | TECN | COMMENT | |
| 7.7±1.1 OUR FIT | | Error includes scale factor of 1.1. | | | |
| 14.9^{+3.8}_{-3.2}±1.2 | | ¹ AALTONEN | 12A | CDF | $p\bar{p}$ at 1.96 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|--|----|------|----|-----|---------------------------------------|
| seen | 90 | BARI | 91 | SFM | $\Lambda_c^+ \rightarrow p K^- \pi^+$ |
| ¹ AALTONEN 12A reports $[\Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^+ \pi^- \pi^-)/\Gamma_{\text{total}}] / [B(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-)] = 3.04 \pm 0.33^{+0.70}_{-0.55}$ which we multiply by our best value $B(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-) = (4.9 \pm 0.4) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. | | | | | |

| $\Gamma(\Lambda_c^+ \pi^+ \pi^- \pi^-)/\Gamma(\Lambda_c^+ \pi^-)$ | | | | | Γ_{26}/Γ_{21} |
|---|--|-------------|------|---------|---------------------------|
| VALUE | | DOCUMENT ID | TECN | COMMENT | |
| 1.56±0.21 OUR FIT | | | | | |
| 1.43±0.16±0.13 | | AAIJ | 11E | LHCB | pp at 7 TeV |

| $\Gamma(\Lambda_c(2595)^+ \pi^-, \Lambda_c(2595)^+ \rightarrow \Lambda_c^+ \pi^+ \pi^-)/\Gamma(\Lambda_c^+ \pi^+ \pi^- \pi^-)$ | | | | | Γ_{27}/Γ_{26} |
|--|--|-------------|------|---------|---------------------------|
| VALUE (units 10^{-2}) | | DOCUMENT ID | TECN | COMMENT | |
| 4.4±1.7^{+0.6}_{-0.4} | | AAIJ | 11E | LHCB | pp at 7 TeV |

| $\Gamma(\Lambda_c(2625)^+ \pi^-, \Lambda_c(2625)^+ \rightarrow \Lambda_c^+ \pi^+ \pi^-)/\Gamma(\Lambda_c^+ \pi^+ \pi^- \pi^-)$ | | | | | Γ_{28}/Γ_{26} |
|--|--|-------------|------|---------|---------------------------|
| VALUE (units 10^{-2}) | | DOCUMENT ID | TECN | COMMENT | |
| 4.3±1.5±0.4 | | AAIJ | 11E | LHCB | pp at 7 TeV |

| $\Gamma(\Sigma_c(2455)^0 \pi^+ \pi^-, \Sigma_c^0 \rightarrow \Lambda_c^+ \pi^-)/\Gamma(\Lambda_c^+ \pi^+ \pi^- \pi^-)$ | | | | | Γ_{29}/Γ_{26} |
|--|--|-------------|------|---------|---------------------------|
| VALUE (units 10^{-2}) | | DOCUMENT ID | TECN | COMMENT | |
| 7.4±2.4±1.2 | | AAIJ | 11E | LHCB | pp at 7 TeV |

$\Gamma(\Sigma_c(2455)^{++}\pi^-\pi^-, \Sigma_c^{++} \rightarrow \Lambda_c^+\pi^+)/\Gamma(\Lambda_c^+\pi^+\pi^-\pi^-)$ Γ_{30}/Γ_{26}

| VALUE (units 10^{-2}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|------|--------------------|
| $4.2 \pm 1.8 \pm 0.7$ | AAIJ | 11E | LHCB pp at 7 TeV |

$\Gamma(\Lambda_c^+ p \bar{p} \pi^-)/\Gamma(\Lambda_c^+ \pi^-)$ Γ_{31}/Γ_{21}

| VALUE (units 10^{-2}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|------|--------------------------|
| $5.40 \pm 0.23 \pm 0.32$ | AAIJ | 18AW | LHCB pp at 7 and 8 TeV |

$\Gamma(\Sigma_c(2455)^0 p \bar{p}, \Sigma_c(2455)^0 \rightarrow \Lambda_c^+ \pi^-)/\Gamma(\Lambda_c^+ p \bar{p} \pi^-)$ Γ_{32}/Γ_{31}

| VALUE (units 10^{-2}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|------|--------------------------|
| $8.9 \pm 1.5 \pm 0.6$ | AAIJ | 18AW | LHCB pp at 7 and 8 TeV |

$\Gamma(\Sigma_c(2520)^0 p \bar{p}, \Sigma_c(2520)^0 \rightarrow \Lambda_c^+ \pi^-)/\Gamma(\Lambda_c^+ p \bar{p} \pi^-)$ Γ_{33}/Γ_{31}

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-----------------------------|-------------|------|--------------------------|
| $0.119 \pm 0.020 \pm 0.014$ | AAIJ | 18AW | LHCB pp at 7 and 8 TeV |

$\Gamma(\Lambda K^0 2\pi^+ 2\pi^-)/\Gamma_{\text{total}}$ Γ_{34}/Γ

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------|------|-------------|------|---------|
|-------|------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|------|---|----------------------|----|------|-------------------------------|
| seen | 4 | ¹ ARENTON | 86 | FMPS | $\Lambda K_S^0 2\pi^+ 2\pi^-$ |
|------|---|----------------------|----|------|-------------------------------|

¹ See the footnote to the ARENTON 86 mass value.

$\Gamma(\Lambda_c^+ \ell^- \bar{\nu}_\ell \text{ anything})/\Gamma_{\text{total}}$ Γ_{35}/Γ

The values and averages in this section serve only to show what values result if one assumes our $B(b \rightarrow b\text{-baryon})$. They cannot be thought of as measurements since the underlying product branching fractions were also used to determine $B(b \rightarrow b\text{-baryon})$ as described in the note on "Production and Decay of b -Flavored Hadrons."

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------|------|-------------|------|---------|
|-------|------|-------------|------|---------|

0.104 ± 0.022 OUR AVERAGE

| | | | | | |
|-----------------------------|--|---------------------|-----|------|-------------------------|
| $0.098 \pm 0.018 \pm 0.013$ | | ¹ BARATE | 98D | ALEP | $e^+ e^- \rightarrow Z$ |
|-----------------------------|--|---------------------|-----|------|-------------------------|

| | | | | | |
|---------------------------------|----|--------------------|-----|------|-------------------------|
| $0.13^{+0.05}_{-0.04} \pm 0.02$ | 29 | ² ABREU | 95S | DLPH | $e^+ e^- \rightarrow Z$ |
|---------------------------------|----|--------------------|-----|------|-------------------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|-----------------------------|----|-----------------------|-----|------|---------------------|
| $0.086 \pm 0.021 \pm 0.012$ | 55 | ³ BUSKULIC | 95L | ALEP | Repl. by BARATE 98D |
|-----------------------------|----|-----------------------|-----|------|---------------------|

| | | | | | |
|--------------------------|----|-----------------------|-----|------|---------------------------------------|
| $0.17 \pm 0.06 \pm 0.02$ | 21 | ⁴ BUSKULIC | 92E | ALEP | $\Lambda_c^+ \rightarrow p K^- \pi^+$ |
|--------------------------|----|-----------------------|-----|------|---------------------------------------|

¹ BARATE 98D reports $[\Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+ \ell^- \bar{\nu}_\ell \text{ anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})] = 0.0086 \pm 0.0007 \pm 0.0014$ which we divide by our best value $B(\bar{b} \rightarrow b\text{-baryon}) = (8.8 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Measured using $\Lambda_c \ell^-$ and $\Lambda \ell^+ \ell^-$.

² ABREU 95S reports $[\Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+ \ell^- \bar{\nu}_\ell \text{ anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})] = 0.0118 \pm 0.0026^{+0.0031}_{-0.0021}$ which we divide by our best value $B(\bar{b} \rightarrow b\text{-baryon}) = (8.8 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ BUSKULIC 95L reports $[\Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+ \ell^- \bar{\nu}_\ell \text{ anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})] = 0.00755 \pm 0.0014 \pm 0.0012$ which we divide by our best value $B(\bar{b} \rightarrow b\text{-baryon}) = (8.8 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ BUSKULIC 92E reports $[\Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+ \ell^- \bar{\nu}_\ell \text{ anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})] = 0.015 \pm 0.0035 \pm 0.0045$ which we divide by our best value $B(\bar{b} \rightarrow b\text{-baryon}) = (8.8 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Superseded by BUSKULIC 95L.

| | |
|---|---|
| $\Gamma(\Lambda_c^+ \ell^- \bar{\nu}_\ell)/\Gamma_{\text{total}}$ | Γ_{36}/Γ |
| <u>VALUE</u> | <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u> |

0.062^{+0.014}_{-0.013} OUR FIT

0.050^{+0.011+0.016}_{-0.008-0.012} ¹ ABDALLAH 04A DLPH $e^+e^- \rightarrow Z^0$

¹ Derived from a combined likelihood and event rate fit to the distribution of the Isgur-Wise variable and using HQET. The slope of the form factor is measured to be $\rho^2 = 2.03 \pm 0.46^{+0.72}_{-1.00}$.

| | |
|---|---|
| $\Gamma(\Lambda_c^+ \ell^- \bar{\nu}_\ell)/\Gamma(\Lambda_c^+ \pi^-)$ | Γ_{36}/Γ_{21} |
| <u>VALUE</u> | <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u> |

12.7^{+3.1}_{-2.7} OUR FIT

16.6 \pm 3.0^{+2.8}_{-3.6} AALTONEN 09E CDF $\rho\bar{p}$ at 1.96 TeV

| | |
|---|---|
| $\Gamma(\Lambda_c^+ \pi^+ \pi^- \ell^- \bar{\nu}_\ell)/\Gamma_{\text{total}}$ | Γ_{37}/Γ |
| <u>VALUE</u> | <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u> |

0.056^{+0.031}_{-0.030} ¹ ABDALLAH 04A DLPH $e^+e^- \rightarrow Z^0$

¹ Derived from the fraction of $\Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+ \ell^- \bar{\nu}_\ell) / (\Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+ \ell^- \bar{\nu}_\ell) + \Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^+ \pi^- \ell^- \bar{\nu}_\ell)) = 0.47^{+0.10+0.07}_{-0.08-0.06}$.

| | |
|---|---|
| $\Gamma(\Lambda_c^+ \ell^- \bar{\nu}_\ell)/[\Gamma(\Lambda_c^+ \ell^- \bar{\nu}_\ell) + \Gamma(\Lambda_c^+ \pi^+ \pi^- \ell^- \bar{\nu}_\ell)]$ | $\Gamma_{36}/(\Gamma_{36}+\Gamma_{37})$ |
| <u>VALUE</u> | <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u> |

0.47^{+0.10+0.07}_{-0.08-0.06} ABDALLAH 04A DLPH $e^+e^- \rightarrow Z^0$

| | |
|---|---|
| $\Gamma(\Lambda_c(2595)^+ \ell^- \bar{\nu}_\ell)/\Gamma(\Lambda_c^+ \ell^- \bar{\nu}_\ell)$ | Γ_{38}/Γ_{36} |
| <u>VALUE</u> | <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u> |

0.126 \pm 0.033^{+0.047}_{-0.038} AALTONEN 09E CDF $\rho\bar{p}$ at 1.96 TeV

| | |
|---|---|
| $\Gamma(\Lambda_c(2625)^+ \ell^- \bar{\nu}_\ell)/\Gamma(\Lambda_c^+ \ell^- \bar{\nu}_\ell)$ | Γ_{39}/Γ_{36} |
| <u>VALUE</u> | <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u> |

0.210 \pm 0.042^{+0.071}_{-0.050} AALTONEN 09E CDF $\rho\bar{p}$ at 1.96 TeV

| | |
|--|---|
| $[\frac{1}{2}\Gamma(\Sigma_c(2455)^0 \pi^+ \ell^- \bar{\nu}_\ell) + \frac{1}{2}\Gamma(\Sigma_c(2455)^{++} \pi^- \ell^- \bar{\nu}_\ell)]/\Gamma(\Lambda_c^+ \ell^- \bar{\nu}_\ell)$ | $(\frac{1}{2}\Gamma_{40} + \frac{1}{2}\Gamma_{41})/\Gamma_{36}$ |
| <u>VALUE</u> | <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u> |

0.054 \pm 0.022^{+0.021}_{-0.018} AALTONEN 09E CDF $\rho\bar{p}$ at 1.96 TeV

$\Gamma(\rho h^-)/\Gamma_{\text{total}}$ Γ_{42}/Γ

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------------------------|------------|---------------------|-------------|---------------------------|
| <2.3 × 10⁻⁵ | 90 | ¹ ACOSTA | 050 CDF | $\rho\bar{p}$ at 1.96 TeV |

¹ Assumes $f_{\Lambda} / f_d = 0.25$, and equal momentum distribution for Λ_b and B mesons.

$\Gamma(\rho\pi^-)/\Gamma_{\text{total}}$ Γ_{43}/Γ

| <u>VALUE (units 10⁻⁶)</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------------------------|------------|-----------------------|-------------|---------------------------|
| 4.3 ± 0.8 OUR FIT | | | | |
| 3.8 ± 0.8 ± 0.5 | | ¹ AALTONEN | 09C CDF | $\rho\bar{p}$ at 1.96 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-----|----|-----------------------|----------|------------------------|
| <50 | 90 | ² BUSKULIC | 96V ALEP | $e^+e^- \rightarrow Z$ |
|-----|----|-----------------------|----------|------------------------|

¹ AALTONEN 09C reports $[\Gamma(\Lambda_b^0 \rightarrow \rho\pi^-)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow K^+\pi^-)] \times [B(\bar{b} \rightarrow b\text{-baryon})] / [B(\bar{b} \rightarrow B^0)] = 0.042 \pm 0.007 \pm 0.006$ which we multiply or divide by our best values $B(B^0 \rightarrow K^+\pi^-) = (1.96 \pm 0.05) \times 10^{-5}$, $B(\bar{b} \rightarrow b\text{-baryon}) = (8.8 \pm 1.2) \times 10^{-2}$, $B(\bar{b} \rightarrow B^0) = (40.5 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

² BUSKULIC 96V assumes PDG 96 production fractions for B^0 , B^+ , B_s , b baryons.

$\Gamma(\rho K^-)/\Gamma_{\text{total}}$ Γ_{44}/Γ

| <u>VALUE (units 10⁻⁶)</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------------------------|------------|-----------------------|-------------|---------------------------|
| 5.1 ± 0.9 OUR FIT | | | | |
| 5.9 ± 1.1 ± 0.8 | | ¹ AALTONEN | 09C CDF | $\rho\bar{p}$ at 1.96 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|------|----|-------------------|----------|------------------------|
| <360 | 90 | ² ADAM | 96D DLPH | $e^+e^- \rightarrow Z$ |
|------|----|-------------------|----------|------------------------|

| | | | | |
|------|----|-----------------------|----------|------------------------|
| < 50 | 90 | ³ BUSKULIC | 96V ALEP | $e^+e^- \rightarrow Z$ |
|------|----|-----------------------|----------|------------------------|

¹ AALTONEN 09C reports $[\Gamma(\Lambda_b^0 \rightarrow \rho K^-)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow K^+\pi^-)] \times [B(\bar{b} \rightarrow b\text{-baryon})] / [B(\bar{b} \rightarrow B^0)] = 0.066 \pm 0.009 \pm 0.008$ which we multiply or divide by our best values $B(B^0 \rightarrow K^+\pi^-) = (1.96 \pm 0.05) \times 10^{-5}$, $B(\bar{b} \rightarrow b\text{-baryon}) = (8.8 \pm 1.2) \times 10^{-2}$, $B(\bar{b} \rightarrow B^0) = (40.5 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

² ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$.

³ BUSKULIC 96V assumes PDG 96 production fractions for B^0 , B^+ , B_s , b baryons.

$\Gamma(\rho\pi^-)/\Gamma(\rho K^-)$ Γ_{43}/Γ_{44}

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------------------|--------------------|-------------|----------------|
| 0.84 ± 0.09 OUR FIT | | | |
| 0.86 ± 0.08 ± 0.05 | AAIJ | 12AR LHCb | pp at 7 TeV |

$\Gamma(\rho D_s^-)/\Gamma_{\text{total}}$ Γ_{45}/Γ

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------------------------|------------|--------------------|-------------|----------------|
| <4.8 × 10⁻⁴ | 90 | AAIJ | 14Q LHCb | pp at 7 TeV |

$\Gamma(\rho\mu^-\bar{\nu}_\mu)/\Gamma_{\text{total}}$ Γ_{46}/Γ

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------------|------|--------------------|
| 4.1±1.0 | ¹ AAIJ | 15BG | LHCB pp at 8 TeV |

¹ The ratio of $B(\Lambda_b^0 \rightarrow \rho\mu^-\bar{\nu}_\mu)$ to $B(\Lambda_b^0 \rightarrow \Lambda_c^+\mu^-\bar{\nu}_\mu)$ is measured within a restricted q^2 region. Combined with theoretical calculations of the form factors and the previously measured value of $|V_{cb}|$, the first $|V_{ub}| = (3.27 \pm 0.15 \pm 0.16 \pm 0.06) \times 10^{-3}$ measurement from the Λ_b decay is obtained, consistent with the exclusively measured world averages.

$\Gamma(\rho\mu^-\bar{\nu}_\mu)/\Gamma(\Lambda_c^+\ell^-\bar{\nu}_\ell)$ Γ_{46}/Γ_{36}

| VALUE (units 10^{-2}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|------|---------|
|--------------------------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|---------------|-------------------|------|--------------------|
| 1.0±0.04±0.08 | ¹ AAIJ | 15BG | LHCB pp at 8 TeV |
|---------------|-------------------|------|--------------------|

¹ This measurement is a ratio of $\Gamma(\Lambda_b^0 \rightarrow \rho\mu^-\bar{\nu}_\mu)[q^2 > 15 \text{ GeV}/c^2]$ to $\Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+\mu^-\bar{\nu}_\mu)[q^2 > 7 \text{ GeV}/c^2]$ within a restricted q^2 region. Combined with theoretical calculations of the form factors and the previously measured value of $|V_{cb}|$, the first $|V_{ub}| = (3.27 \pm 0.15 \pm 0.16 \pm 0.06) \times 10^{-3}$ measurement from the Λ_b decay is obtained, consistent with the exclusively measured world averages.

$\Gamma(\Lambda\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_{47}/Γ

| VALUE (units 10^{-7}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|------|---------|
|--------------------------|-------------|------|---------|

10.8±2.8 OUR AVERAGE

| | | | |
|--------------|-------------------|------|----------------------------|
| 9.6±1.6±2.5 | ¹ AAIJ | 13AJ | LHCB pp at 7 TeV |
| 17.3±4.2±5.5 | AALTONEN | 11AI | CDF $p\bar{p}$ at 1.96 TeV |

¹ Uses $B(\Lambda_b^0 \rightarrow J/\psi\Lambda) = (6.2 \pm 1.4) \times 10^{-4}$. This measurement comes from the sum of the differential rates in q^2 regions excluding those corresponding to J/ψ and $\psi(2S)$ ([8.68,10.09] and [12.86, 14.18] GeV^2/c^4).

$\Gamma(\rho\pi^-\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_{48}/Γ

| VALUE (units 10^{-8}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|------|---------|
|--------------------------|-------------|------|---------|

| | | | |
|--|-------------------|-----|-----------------------|
| 6.9±1.9^{+1.7}_{-1.5} | ¹ AAIJ | 17P | LHCB pp at 7, 8 TeV |
|--|-------------------|-----|-----------------------|

¹ Excludes J/ψ and $\psi(2S)$ decays to $\mu^+\mu^-$.

$\Gamma(\rho\pi^-\mu^+\mu^-)/\Gamma(\rho\pi^-J/\psi, J/\psi \rightarrow \mu^+\mu^-)$ Γ_{48}/Γ_{10}

| VALUE (units 10^{-2}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|------|---------|
|--------------------------|-------------|------|---------|

| | | | |
|--------------------|-------------------|-----|-----------------------|
| 4.4±1.2±0.7 | ¹ AAIJ | 17P | LHCB pp at 7, 8 TeV |
|--------------------|-------------------|-----|-----------------------|

¹ The $\rho\pi^-\mu^+\mu^-$ mode excludes J/ψ and $\psi(2S)$ decays to $\mu^+\mu^-$.

$\Gamma(\Lambda\gamma)/\Gamma_{\text{total}}$ Γ_{49}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

| | | | | |
|----------------------------------|----|--------|-----|---------------------------|
| <1.3 × 10⁻³ | 90 | ACOSTA | 02G | CDF $p\bar{p}$ at 1.8 TeV |
|----------------------------------|----|--------|-----|---------------------------|

$\Gamma(\Lambda\eta)/\Gamma_{\text{total}}$ Γ_{50}/Γ

| VALUE (units 10^{-6}) | DOCUMENT ID | TECN | COMMENT |
|---------------------------------------|-------------------|-----------|------------------|
| $9^{+7}_{-5} \pm 1$ | ¹ AAIJ | 15AH LHCB | pp at 7, 8 TeV |

¹ AAIJ 15AH reports $[\Gamma(\Lambda_b^0 \rightarrow \Lambda\eta)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow \eta' K^0)] = 0.142^{+0.11}_{-0.08}$ which we multiply by our best value $B(B^0 \rightarrow \eta' K^0) = (6.6 \pm 0.4) \times 10^{-5}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. The single uncertainty quoted with the original measurement combines in quadrature statistical and systematic uncertainties.

$\Gamma(\Lambda\eta'(958))/\Gamma_{\text{total}}$ Γ_{51}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------------|-----------|------------------|
| $< 3.1 \times 10^{-6}$ | 90 | ¹ AAIJ | 15AH LHCB | pp at 7, 8 TeV |

¹ AAIJ 15AH reports $[\Gamma(\Lambda_b^0 \rightarrow \Lambda\eta'(958))/\Gamma_{\text{total}}] / [B(B^0 \rightarrow \eta' K^0)] < 0.047$ which we multiply by our best value $B(B^0 \rightarrow \eta' K^0) = 6.6 \times 10^{-5}$.

$\Gamma(\Lambda\pi^+\pi^-)/\Gamma(\Lambda_c^+\pi^-)$ Γ_{52}/Γ_{21}

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|---|-------------------|----------|------------------|
| $9.5 \pm 3.8 \pm 0.5$ | ¹ AAIJ | 16W LHCB | pp at 7, 8 TeV |

¹ AAIJ 16W reports $[\Gamma(\Lambda_b^0 \rightarrow \Lambda\pi^+\pi^-)/\Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+\pi^-)] / [B(\Lambda_c^+ \rightarrow \Lambda\pi^+)] = (7.3 \pm 1.9 \pm 2.2) \times 10^{-2}$ which we multiply by our best value $B(\Lambda_c^+ \rightarrow \Lambda\pi^+) = (1.30 \pm 0.07) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\Lambda K^+\pi^-)/\Gamma(\Lambda_c^+\pi^-)$ Γ_{53}/Γ_{21}

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|--|-------------------|----------|------------------|
| $11.6 \pm 2.3 \pm 0.6$ | ¹ AAIJ | 16W LHCB | pp at 7, 8 TeV |

¹ AAIJ 16W reports $[\Gamma(\Lambda_b^0 \rightarrow \Lambda K^+\pi^-)/\Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+\pi^-)] / [B(\Lambda_c^+ \rightarrow \Lambda\pi^+)] = (8.9 \pm 1.2 \pm 1.3) \times 10^{-2}$ which we multiply by our best value $B(\Lambda_c^+ \rightarrow \Lambda\pi^+) = (1.30 \pm 0.07) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\Lambda K^+K^-)/\Gamma(\Lambda_c^+\pi^-)$ Γ_{54}/Γ_{21}

| VALUE (units 10^{-3}) | DOCUMENT ID | TECN | COMMENT |
|--|-------------------|----------|------------------|
| $3.29 \pm 0.35 \pm 0.17$ | ¹ AAIJ | 16W LHCB | pp at 7, 8 TeV |

¹ AAIJ 16W reports $[\Gamma(\Lambda_b^0 \rightarrow \Lambda K^+K^-)/\Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+\pi^-)] / [B(\Lambda_c^+ \rightarrow \Lambda\pi^+)] = (25.3 \pm 1.9 \pm 1.9) \times 10^{-2}$ which we multiply by our best value $B(\Lambda_c^+ \rightarrow \Lambda\pi^+) = (1.30 \pm 0.07) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\Lambda\phi)/\Gamma_{\text{total}}$ Γ_{55}/Γ

| VALUE (units 10^{-6}) | DOCUMENT ID | TECN | COMMENT |
|---|-------------------|----------|------------------|
| $9.3 \pm 2.0 \pm 1.5$ | ¹ AAIJ | 16J LHCB | pp at 7, 8 TeV |

¹ AAIJ 16J reports $[\Gamma(\Lambda_b^0 \rightarrow \Lambda \phi) / \Gamma_{\text{total}}] / [B(B^0 \rightarrow K^0 \phi)] \times [B(\bar{b} \rightarrow b\text{-baryon})] / [B(\bar{b} \rightarrow B^0)] = 0.275 \pm 0.055 \pm 0.020$ which we multiply or divide by our best values $B(B^0 \rightarrow K^0 \phi) = (7.3 \pm 0.7) \times 10^{-6}$, $B(\bar{b} \rightarrow b\text{-baryon}) = (8.8 \pm 1.2) \times 10^{-2}$, $B(\bar{b} \rightarrow B^0) = (40.5 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

$\Gamma(p\pi^-\pi^+\pi^-) / \Gamma(\Lambda_c^+ \pi^-)$ $\Gamma_{56} / \Gamma_{21}$

| <u>VALUE (units 10^{-3})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|-----------------------|
| $4.30 \pm 0.24^{+0.22}_{-0.23}$ | ¹ AAIJ | 18Q | LHCB pp at 7, 8 TeV |

¹ AAIJ 18Q reports $[\Gamma(\Lambda_b^0 \rightarrow p\pi^-\pi^+\pi^-) / \Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-)] / [B(\Lambda_c^+ \rightarrow pK^-\pi^+)] = (6.85 \pm 0.19 \pm 0.08 \pm 0.32) \times 10^{-2}$ which we multiply by our best value $B(\Lambda_c^+ \rightarrow pK^-\pi^+) = (6.28 \pm 0.32) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(pK^-K^+\pi^-) / \Gamma(\Lambda_c^+ \pi^-)$ $\Gamma_{57} / \Gamma_{21}$

| <u>VALUE (units 10^{-3})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|--------------------|-------------|-----------------------|
| $0.83 \pm 0.10 \pm 0.04$ | ¹ AAIJ | 18Q | LHCB pp at 7, 8 TeV |

¹ AAIJ 18Q reports $[\Gamma(\Lambda_b^0 \rightarrow pK^-K^+\pi^-) / \Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-)] / [B(\Lambda_c^+ \rightarrow pK^-\pi^+)] = (1.32 \pm 0.09 \pm 0.09 \pm 0.10) \times 10^{-2}$ which we multiply by our best value $B(\Lambda_c^+ \rightarrow pK^-\pi^+) = (6.28 \pm 0.32) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(pK^-\pi^+\pi^-) / \Gamma(\Lambda_c^+ \pi^-)$ $\Gamma_{58} / \Gamma_{21}$

| <u>VALUE (units 10^{-3})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|-----------------------|
| $10.3 \pm 0.5 \pm 0.5$ | ¹ AAIJ | 18Q | LHCB pp at 7, 8 TeV |

¹ AAIJ 18Q reports $[\Gamma(\Lambda_b^0 \rightarrow pK^-\pi^+\pi^-) / \Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-)] / [B(\Lambda_c^+ \rightarrow pK^-\pi^+)] = (16.4 \pm 0.3 \pm 0.2 \pm 0.7) \times 10^{-2}$ which we multiply by our best value $B(\Lambda_c^+ \rightarrow pK^-\pi^+) = (6.28 \pm 0.32) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(pK^-K^+K^-) / \Gamma(\Lambda_c^+ \pi^-)$ $\Gamma_{59} / \Gamma_{21}$

| <u>VALUE (units 10^{-3})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|-----------------------|
| $2.58 \pm 0.15^{+0.13}_{-0.14}$ | ¹ AAIJ | 18Q | LHCB pp at 7, 8 TeV |

¹ AAIJ 18Q reports $[\Gamma(\Lambda_b^0 \rightarrow pK^-K^+K^-) / \Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-)] / [B(\Lambda_c^+ \rightarrow pK^-\pi^+)] = (4.11 \pm 0.12 \pm 0.06 \pm 0.19) \times 10^{-2}$ which we multiply by our best value $B(\Lambda_c^+ \rightarrow pK^-\pi^+) = (6.28 \pm 0.32) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

PARTIAL BRANCHING FRACTIONS IN $\Lambda_b \rightarrow \Lambda \mu^+ \mu^-$ **$B(\Lambda_b \rightarrow \Lambda \mu^+ \mu^-)$ ($q^2 < 2.0 \text{ GeV}^2/c^4$)**

| <u>VALUE (units 10^{-7})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|--------------------|-------------|------------------------|
| 0.71\pm0.27 OUR AVERAGE | | | |
| 0.72 $^{+0.24}_{-0.22}$ \pm 0.14 | ¹ AAIJ | 15AE LHCB | pp at 7, 8 TeV |
| 0.15 \pm 2.01 \pm 0.05 | AALTONEN | 11AI CDF | $p\bar{p}$ at 1.96 TeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 0.56 \pm 0.76 \pm 0.80 | ² AAIJ | 13AJ LHCB | Repl. by AAIJ 15AE |
| ¹ AAIJ 15AE measurement covers $0.1 < q^2 < 2.0 \text{ GeV}^2/c^4$. | | | |
| ² Uses $B(\Lambda_b^0 \rightarrow J/\psi \Lambda) = (6.2 \pm 1.4) \times 10^{-4}$. | | | |

 $B(\Lambda_b \rightarrow \Lambda \mu^+ \mu^-)$ ($2.0 < q^2 < 4.3 \text{ GeV}^2/c^4$)

| <u>VALUE (units 10^{-7})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|--------------------|-------------|------------------------|
| 0.28$^{+0.28}_{-0.21}$ OUR AVERAGE | | | |
| 0.253 $^{+0.276}_{-0.207}$ \pm 0.046 | ¹ AAIJ | 15AE LHCB | pp at 7, 8 TeV |
| 1.8 \pm 1.7 \pm 0.6 | AALTONEN | 11AI CDF | $p\bar{p}$ at 1.96 TeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 0.71 \pm 0.60 \pm 0.23 | ² AAIJ | 13AJ LHCB | Repl. by AAIJ 15AE |
| ¹ AAIJ 15AE measurement covers $2.0 < q^2 < 4.0 \text{ GeV}^2/c^4$. | | | |
| ² Uses $B(\Lambda_b^0 \rightarrow J/\psi \Lambda) = (6.2 \pm 1.4) \times 10^{-4}$. | | | |

 $B(\Lambda_b \rightarrow \Lambda \mu^+ \mu^-)$ ($q^2 < 4.3 \text{ GeV}^2/c^4$)

| <u>VALUE (units 10^{-7})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|--------------------|-------------|------------------------|
| 2.7\pm2.5\pm0.9 | AALTONEN | 11AI CDF | $p\bar{p}$ at 1.96 TeV |

 $B(\Lambda_b \rightarrow \Lambda \mu^+ \mu^-)$ ($4.0 < q^2 < 6.0 \text{ GeV}^2/c^4$)

| <u>VALUE (units 10^{-7})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|--------------------|-------------|------------------|
| 0.04$^{+0.18}_{-0.00}$$\pm$0.02 | AAIJ | 15AE LHCB | pp at 7, 8 TeV |

 $B(\Lambda_b \rightarrow \Lambda \mu^+ \mu^-)$ ($1.0 < q^2 < 6.0 \text{ GeV}^2/c^4$)

| <u>VALUE (units 10^{-7})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|------------------------|
| 0.47$^{+0.31}_{-0.27}$ OUR AVERAGE | | | |
| 0.45 $^{+0.30}_{-0.25}$ \pm 0.10 | ¹ AAIJ | 15AE LHCB | pp at 7 and 8 TeV |
| 1.3 \pm 2.1 \pm 0.4 | AALTONEN | 11AI CDF | $p\bar{p}$ at 1.96 TeV |
| ¹ AAIJ 15AE measurement covers $1.1 < q^2 < 6.0 \text{ GeV}^2/c^4$. | | | |

 $B(\Lambda_b \rightarrow \Lambda \mu^+ \mu^-)$ ($6.0 < q^2 < 8.0 \text{ GeV}^2/c^4$)

| <u>VALUE (units 10^{-7})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|--------------------|-------------|------------------|
| 0.50$^{+0.24}_{-0.22}$$\pm$0.10 | AAIJ | 15AE LHCB | pp at 7, 8 TeV |

$B(\Lambda_b \rightarrow \Lambda \mu^+ \mu^-) (4.3 < q^2 < 8.68 \text{ GeV}^2/c^4)$

| <u>VALUE (units 10^{-7})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|--------------------|-------------|------------------------|
| 0.5 ± 0.7 OUR AVERAGE | | | |
| $0.66 \pm 0.74 \pm 0.18$ | ¹ AAIJ | 13AJ LHCB | pp at 7 TeV |
| $-0.2 \pm 1.6 \pm 0.1$ | AALTONEN | 11AI CDF | $p\bar{p}$ at 1.96 TeV |
| ¹ Uses $B(\Lambda_b^0 \rightarrow J/\psi \Lambda) = (6.2 \pm 1.4) \times 10^{-4}$. | | | |

 $B(\Lambda_b \rightarrow \Lambda \mu^+ \mu^-) (10.09 < q^2 < 12.86 \text{ GeV}^2/c^4)$

| <u>VALUE (units 10^{-7})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|--------------------|-------------|------------------------|
| 2.2 ± 0.6 OUR AVERAGE | | | |
| $2.08^{+0.42}_{-0.39} \pm 0.42$ | ¹ AAIJ | 15AE LHCB | pp at 7, 8 TeV |
| $3.0 \pm 1.5 \pm 1.0$ | AALTONEN | 11AI CDF | $p\bar{p}$ at 1.96 TeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| $1.55 \pm 0.58 \pm 0.55$ | ² AAIJ | 13AJ LHCB | Repl. by AAIJ 15AE |
| ¹ AAIJ 15AE measurement covers $11.0 < q^2 < 12.5 \text{ GeV}^2/c^4$. | | | |
| ² Uses $B(\Lambda_b^0 \rightarrow J/\psi \Lambda) = (6.2 \pm 1.4) \times 10^{-4}$. | | | |

 $B(\Lambda_b \rightarrow \Lambda \mu^+ \mu^-) (14.18 < q^2 < 16.0 \text{ GeV}^2/c^4)$

| <u>VALUE (units 10^{-7})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|-------------------------------------|-------------|------------------------|
| 1.7 ± 0.5 OUR AVERAGE | Error includes scale factor of 1.1. | | |
| $2.04^{+0.35}_{-0.33} \pm 0.42$ | ¹ AAIJ | 15AE LHCB | pp at 7, 8 TeV |
| $1.0 \pm 0.7 \pm 0.3$ | AALTONEN | 11AI CDF | $p\bar{p}$ at 1.96 TeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| $1.44 \pm 0.44 \pm 0.42$ | ² AAIJ | 13AJ LHCB | Repl. by AAIJ 15AE |
| ¹ AAIJ 15AE measurement covers $15.0 < q^2 < 16.0 \text{ GeV}^2/c^4$. | | | |
| ² Uses $B(\Lambda_b^0 \rightarrow J/\psi \Lambda) = (6.2 \pm 1.4) \times 10^{-4}$. | | | |

 $B(\Lambda_b \rightarrow \Lambda \mu^+ \mu^-) (16.0 < q^2 \text{ GeV}^2/c^4)$

| <u>VALUE (units 10^{-7})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|---------------------|-------------|------------------------|
| $7.0 \pm 1.9 \pm 2.2$ | AALTONEN | 11AI CDF | $p\bar{p}$ at 1.96 TeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| $4.73 \pm 0.77 \pm 1.25$ | ^{1,2} AAIJ | 13AJ LHCB | Repl. by AAIJ 15AE |
| ¹ Uses $B(\Lambda_b^0 \rightarrow J/\psi \Lambda) = (6.2 \pm 1.4) \times 10^{-4}$. | | | |
| ² Requires $16.00 < q^2 < 20.30 \text{ GeV}^2/c^4$. | | | |

 $B(\Lambda_b \rightarrow \Lambda \mu^+ \mu^-) (18.0 < q^2 < 20.0 \text{ GeV}^2/c^4)$

| <u>VALUE (units 10^{-7})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|--------------------|-------------|------------------|
| $2.44 \pm 0.28 \pm 0.50$ | AAIJ | 15AE LHCB | pp at 7, 8 TeV |

 $B(\Lambda_b \rightarrow \Lambda \mu^+ \mu^-) (15.0 < q^2 < 20.0 \text{ GeV}^2/c^4)$

| <u>VALUE (units 10^{-7})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|--------------------|-------------|------------------|
| $6.00 \pm 0.45 \pm 1.25$ | AAIJ | 15AE LHCB | pp at 7, 8 TeV |

CP VIOLATION

A_{CP} is defined as

$$A_{CP} = \frac{B(\Lambda_b^0 \rightarrow f) - B(\bar{\Lambda}_b^0 \rightarrow \bar{f})}{B(\Lambda_b^0 \rightarrow f) + B(\bar{\Lambda}_b^0 \rightarrow \bar{f})},$$

the CP-violation asymmetry of exclusive Λ_b^0 and $\bar{\Lambda}_b^0$ decay.

$A_{CP}(\Lambda_b \rightarrow p\pi^-)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------|------|-------------------------------------|
| -0.025 ± 0.029 OUR AVERAGE | | | Error includes scale factor of 1.2. |
| $-0.035 \pm 0.017 \pm 0.020$ | AAIJ | 18AX | LHCB pp at 7 and 8 TeV |
| $0.06 \pm 0.07 \pm 0.03$ | AALTONEN | 14P | CDF $p\bar{p}$ at 1.96 TeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| $0.03 \pm 0.17 \pm 0.05$ | AALTONEN | 11N | CDF Repl. by AALTONEN 14P |

$A_{CP}(\Lambda_b \rightarrow pK^-)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------|------|----------------------------|
| -0.025 ± 0.022 OUR AVERAGE | | | |
| $-0.020 \pm 0.013 \pm 0.019$ | AAIJ | 18AX | LHCB pp at 7 and 8 TeV |
| $-0.10 \pm 0.08 \pm 0.04$ | AALTONEN | 14P | CDF $p\bar{p}$ at 1.96 TeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| $0.37 \pm 0.17 \pm 0.03$ | AALTONEN | 11N | CDF Repl. by AALTONEN 14P |

$\Delta A_{CP}(pK^-/\pi^-) \equiv A_{CP}(pK^-) - A_{CP}(p\pi^-)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------|------|--------------------------|
| $0.014 \pm 0.022 \pm 0.010$ | AAIJ | 18AX | LHCB pp at 7 and 8 TeV |

$A_{CP}(\Lambda_b \rightarrow p\bar{K}^0\pi^-)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--|-------------|------|--------------------|
| $0.22 \pm 0.13 \pm 0.03$ | AAIJ | 14Q | LHCB pp at 7 TeV |

$\Delta A_{CP}(J/\psi p\pi^-/K^-) \equiv A_{CP}(J/\psi p\pi^-) - A_{CP}(J/\psi pK^-)$

| VALUE (units 10^{-2}) | DOCUMENT ID | TECN | COMMENT |
|---|-------------|------|-----------------------|
| $5.7 \pm 2.4 \pm 1.2$ | AAIJ | 14K | LHCB pp at 7, 8 TeV |

$A_{CP}(\Lambda_b \rightarrow \Lambda K^+\pi^-)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------------|------|-----------------------|
| $-0.53 \pm 0.23 \pm 0.11$ | ¹ AAIJ | 16W | LHCB pp at 7, 8 TeV |

¹ Measured relative to $\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-$ decay.

$A_{CP}(\Lambda_b \rightarrow \Lambda K^+ K^-)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------------|------|-----------------------|
| $-0.28 \pm 0.10 \pm 0.07$ | ¹ AAIJ | 16W | LHCB pp at 7, 8 TeV |

¹ Measured relative to $\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-$ decay.

$$\Delta A_{CP}(\Lambda_b^0 \rightarrow pK^- \mu^+ \mu^-) \equiv A_{CP}(pK^- \mu^+ \mu^-) - A_{CP}(pK^- J/\psi)$$

| VALUE (units 10^{-2}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|------|-----------------------|
| $-3.5 \pm 5.0 \pm 0.2$ | AAIJ | 17T | LHCB pp at 7, 8 TeV |

CP AND T VIOLATION PARAMETERS

Measured values of the triple-product asymmetry parameters, odd under time-reversal, are defined as $A_{c(s)}(\Lambda/\phi) = (N_{c(s)}^+ - N_{c(s)}^-) / (\text{sum})$ where $N_{c(s)}^+$, $N_{c(s)}^-$ are the number of Λ or ϕ candidates for which the $\cos(\phi)$ and $\sin(\phi)$ observables are positive and negative, respectively. Angles $\cos(\phi)$ and $\sin(\phi)$ are defined as in LEITNER 07.

$A_c(\Lambda)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---------------------------|-------------|------|-----------------------|
| $-0.22 \pm 0.12 \pm 0.06$ | AAIJ | 16J | LHCB pp at 7, 8 TeV |

$A_s(\Lambda)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|------|-----------------------|
| $0.13 \pm 0.12 \pm 0.05$ | AAIJ | 16J | LHCB pp at 7, 8 TeV |

$A_c(\phi)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---------------------------|-------------|------|-----------------------|
| $-0.01 \pm 0.12 \pm 0.03$ | AAIJ | 16J | LHCB pp at 7, 8 TeV |

$A_s(\phi)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---------------------------|-------------|------|-----------------------|
| $-0.07 \pm 0.12 \pm 0.01$ | AAIJ | 16J | LHCB pp at 7, 8 TeV |

$a_{CP}(\Lambda_b^0 \rightarrow p\pi^- \pi^+ \pi^-)$

Observable calculated as half of the difference between triple products for Λ_b^0 and $\bar{\Lambda}_b^0$, which is sensitive to CP violation.

| VALUE (%) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------------|------|-----------------------|
| $1.15 \pm 1.45 \pm 0.32$ | ¹ AAIJ | 17H | LHCB pp at 7, 8 TeV |

¹ Measured over full phase space of the decay.

$a_{CP}(\Lambda_b^0 \rightarrow pK^- \pi^+ \pi^-)$

Observable calculated as half of the difference between triple products for Λ_b^0 and $\bar{\Lambda}_b^0$, which is sensitive to CP violation.

| VALUE (%) | DOCUMENT ID | TECN | COMMENT |
|---------------------------|-------------------|------|-----------------------|
| $-0.81 \pm 0.84 \pm 0.31$ | ¹ AAIJ | 18AG | LHCB pp at 7, 8 TeV |

¹ Measured over full phase space of the decay.

$a_{CP}(\Lambda_b^0 \rightarrow pK^- K^+ \pi^-)$

Observable calculated as half of the difference between triple products for Λ_b^0 and $\bar{\Lambda}_b^0$, which is sensitive to CP violation.

| VALUE (%) | DOCUMENT ID | TECN | COMMENT |
|---------------------------|-------------------|------|-----------------------|
| $-0.93 \pm 4.54 \pm 0.42$ | ¹ AAIJ | 17H | LHCB pp at 7, 8 TeV |

¹ Measured over full phase space of the decay.

$a_{CP}(\Lambda_b^0 \rightarrow p K^- K^+ K^-)$

Observable calculated as half of the difference between triple products for Λ_b^0 and $\bar{\Lambda}_b^0$, which is sensitive to CP violation.

| VALUE (%) | DOCUMENT ID | TECN | COMMENT |
|--|-------------------|-----------|------------------|
| $1.12 \pm 1.51 \pm 0.32$ | ¹ AAIJ | 18AG LHCB | pp at 7, 8 TeV |

¹ Measured over full phase space of the decay.

$a_{CP}(\Lambda_b^0 \rightarrow p K^- \mu^+ \mu^-)$

| VALUE (%) | DOCUMENT ID | TECN | COMMENT |
|---|-------------|----------|------------------|
| $1.2 \pm 5.0 \pm 0.7$ | AAIJ | 17T LHCB | pp at 7, 8 TeV |

P VIOLATION PARAMETERS

Observables calculated as average of the triple products for Λ_b^0 and $\bar{\Lambda}_b^0$, which is sensitive to parity violation.

$a_P(\Lambda_b^0 \rightarrow p \pi^- \pi^+ \pi^-)$

| VALUE (%) | DOCUMENT ID | TECN | COMMENT |
|---|-------------------|----------|------------------|
| $-3.71 \pm 1.45 \pm 0.32$ | ¹ AAIJ | 17H LHCB | pp at 7, 8 TeV |

¹ Measured over full phase space of the decay.

$a_P(\Lambda_b^0 \rightarrow p K^- \pi^+ \pi^-)$

| VALUE (%) | DOCUMENT ID | TECN | COMMENT |
|---|-------------------|-----------|------------------|
| $-0.60 \pm 0.84 \pm 0.31$ | ¹ AAIJ | 18AG LHCB | pp at 7, 8 TeV |

¹ Measured over full phase space of the decay.

$a_P(\Lambda_b^0 \rightarrow p K^- K^+ \pi^-)$

| VALUE (%) | DOCUMENT ID | TECN | COMMENT |
|--|-------------------|----------|------------------|
| $3.62 \pm 4.54 \pm 0.42$ | ¹ AAIJ | 17H LHCB | pp at 7, 8 TeV |

¹ Measured over full phase space of the decay.

$a_P(\Lambda_b^0 \rightarrow p K^- K^+ K^-)$

| VALUE (%) | DOCUMENT ID | TECN | COMMENT |
|---|-------------------|-----------|------------------|
| $-1.56 \pm 1.51 \pm 0.32$ | ¹ AAIJ | 18AG LHCB | pp at 7, 8 TeV |

¹ Measured over full phase space of the decay.

$a_P(\Lambda_b^0 \rightarrow p K^- \mu^+ \mu^-)$

| VALUE (%) | DOCUMENT ID | TECN | COMMENT |
|--|-------------|----------|------------------|
| $-4.8 \pm 5.0 \pm 0.7$ | AAIJ | 17T LHCB | pp at 7, 8 TeV |


Λ_b^0 DECAY PARAMETERS

See the note on "Baryon Decay Parameters" in the neutron Listings.

α decay parameter for $\Lambda_b \rightarrow J/\psi \Lambda$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|---|-----------|------------------|
| 0.07 ± 0.13 OUR AVERAGE | Error includes scale factor of 1.3. See the ideogram below. | | |
| $-0.14 \pm 0.14 \pm 0.10$ | ¹ SIRUNYAN | 18R CMS | pp at 7, 8 TeV |
| $0.30 \pm 0.16 \pm 0.06$ | ² AAD | 14L ATLS | pp at 7 TeV |
| $0.05 \pm 0.17 \pm 0.07$ | ³ AAIJ | 13AG LHCB | pp at 7 TeV |

- ¹ An angular analysis of $\Lambda_b \rightarrow J/\psi \Lambda$ decay is performed. Note that the sign of α in CMS definition is the opposite to that used by AAIJ 13AG and AAD 14L. Λ_b transverse production polarization of $0.00 \pm 0.06 \pm 0.06$ is also reported, as well as squares of the helicity amplitudes.
- ² An angular analysis of $\Lambda_b \rightarrow J/\psi \Lambda$ decay is performed and magnitudes of all helicity amplitudes are also reported.
- ³ An angular analysis of $\Lambda_b \rightarrow J/\psi \Lambda$ decay is performed and a Λ_b transverse production polarization of $0.06 \pm 0.07 \pm 0.02$ is also reported.



$f_L(\mu\mu)$ longitudinal polarization fraction in $\Lambda_b \rightarrow \Lambda \mu^+ \mu^-$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---------------------------------|-------------------|-----------|------------------|
| $0.61^{+0.11}_{-0.14} \pm 0.03$ | ¹ AAIJ | 15AE LHCB | pp at 7, 8 TeV |

¹ AAIJ 15AE measurement covers $15.0 < q^2 < 20.0 \text{ GeV}^2/c^4$.

FORWARD-BACKWARD ASYMMETRIES

The forward-backward asymmetry is defined as $A_{FB}(\Lambda_b^0) = [N(F) - N(B)] / [N(F) + N(B)]$, where the forward (F) direction corresponds to a particle (Λ_b^0 or Λ_b^-) sharing valence quark flavors with a beam particle with the same sign of rapidity.

$A_{FB}(\Lambda_b^0 \rightarrow J/\psi \Lambda)$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--------------------------|---------------------|--------|------------------|
| $0.04 \pm 0.07 \pm 0.02$ | ¹ ABAZOV | 15l D0 | pp at 1.96 TeV |

¹ The measured asymmetry integrated over rapidity y in the range of $0.1 < |y| < 2.0$.

$A_{FB}^{\ell}(\mu\mu)$ in $\Lambda_b \rightarrow \Lambda\mu^+\mu^-$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---------------------------|-------------------|-----------|----------------------|
| $-0.39 \pm 0.04 \pm 0.01$ | ¹ AAIJ | 18AP LHCB | pp at 7, 8, 13 TeV |
| $-0.05 \pm 0.09 \pm 0.03$ | ² AAIJ | 15AE LHCB | Repl. by AAIJ 18AP. |

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ The measurement covers $15.0 < q^2 < 20.0 \text{ GeV}^2/c^4$.

² AAIJ 15AE measurement covers $15.0 < q^2 < 20.0 \text{ GeV}^2/c^4$.

$\Delta(A_{FB}^{\ell}(\mu\mu))$ in $\Lambda_b \rightarrow \Lambda\mu^+\mu^-$

Difference of asymmetries $A_{FB}^{\ell}(\mu\mu)$ in $\Lambda_b \rightarrow \Lambda\mu^+\mu^-$ between Λ_b and $\bar{\Lambda}_b$ decays

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---------------------------|-------------|-----------|------------------|
| $-0.05 \pm 0.09 \pm 0.03$ | AAIJ | 18AO LHCB | pp at 7, 8 TeV |

$A_{FB}^h(\rho\pi)$ in $\Lambda_b \rightarrow \Lambda(\rho\pi)\mu^+\mu^-$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---------------------------|-------------------|-----------|----------------------|
| $-0.30 \pm 0.05 \pm 0.02$ | ¹ AAIJ | 18AP LHCB | pp at 7, 8, 13 TeV |
| $-0.29 \pm 0.07 \pm 0.03$ | ² AAIJ | 15AE LHCB | Repl. by AAIJ 18AP. |

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ The measurement covers $15.0 < q^2 < 20.0 \text{ GeV}^2/c^4$.

² AAIJ 15AE measurement covers $15.0 < q^2 < 20.0 \text{ GeV}^2/c^4$.

$A_{FB}^{\ell h}$ in $\Lambda_b \rightarrow \Lambda\mu^+\mu^-$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------------|-----------|----------------------|
| $0.25 \pm 0.04 \pm 0.01$ | ¹ AAIJ | 18AP LHCB | pp at 7, 8, 13 TeV |

¹ The measurement covers $15.0 < q^2 < 20.0 \text{ GeV}^2/c^4$.

$\Lambda_b^0 - \bar{\Lambda}_b^0$ Production Asymmetry

$$A_P(\Lambda_b^0) = [\sigma(\Lambda_b^0) - \sigma(\bar{\Lambda}_b^0)] / [\sigma(\Lambda_b^0) + \sigma(\bar{\Lambda}_b^0)]$$

$A_P(\Lambda_b^0)$

| VALUE (units 10^{-2}) | DOCUMENT ID | TECN | COMMENT |
|---|-------------------------------------|-----------|---------------|
| 2.4 ± 1.6 OUR AVERAGE | Error includes scale factor of 1.1. | | |
| $-0.11 \pm 2.53 \pm 1.08$ | ¹ AAIJ | 17BF LHCB | pp at 7 TeV |
| $3.44 \pm 1.61 \pm 0.76$ | ¹ AAIJ | 17BF LHCB | pp at 8 TeV |

¹ Indirect determination in kinematic range $2 < p_T < 30 \text{ GeV}/c$ and $2.1 < \eta < 4.5$ from production asymmetries of B^+ , B^0 and B_S^0 .


REFERENCES

| | | | | |
|------------|------|------------------------|-----------------------------|--------------------|
| AAIJ | 18AF | JHEP 1808 131 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 18AG | JHEP 1808 039 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 18AO | JHEP 1809 145 (errat.) | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 18AP | JHEP 1809 146 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 18AW | PL B784 101 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 18AX | PL B787 124 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 18Q | JHEP 1802 098 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| SIRUNYAN | 18BY | EPJ C78 457 | A.M. Sirunyan <i>et al.</i> | (CMS Collab.) |
| SIRUNYAN | 18R | PR D97 072010 | A.M. Sirunyan <i>et al.</i> | (CMS Collab.) |
| AAIJ | 17AM | PRL 119 062001 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 17BF | PL B774 139 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 17H | NATP 13 391 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 17P | JHEP 1704 029 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 17S | JHEP 1705 030 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 17T | JHEP 1706 108 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 16 | JHEP 1601 012 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 16A | CP C40 011001 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 16J | PL B759 282 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 16W | JHEP 1605 081 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 16Y | JHEP 1605 132 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAD | 15CH | PL B751 63 | G. Aad <i>et al.</i> | (ATLAS Collab.) |
| AAIJ | 15AE | JHEP 1506 115 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| Also | | JHEP 1809 145 (errat.) | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 15AH | JHEP 1509 006 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 15BG | NATP 11 743 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| ABAZOV | 15I | PR D91 072008 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| AAD | 14L | PR D89 092009 | G. Aad <i>et al.</i> | (ATLAS Collab.) |
| AAIJ | 14AA | PRL 112 202001 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 14E | JHEP 1404 114 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 14H | PR D89 032001 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 14I | JHEP 1408 143 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 14K | JHEP 1407 103 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 14Q | JHEP 1404 087 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 14U | PL B734 122 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AALTONEN | 14B | PR D89 072014 | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| AALTONEN | 14P | PRL 113 242001 | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| PDG | 14 | CP C38 070001 | K. Olive <i>et al.</i> | (PDG Collab.) |
| AAD | 13U | PR D87 032002 | G. Aad <i>et al.</i> | (ATLAS Collab.) |
| AAIJ | 13AG | PL B724 27 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 13AJ | PL B725 25 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 13AV | PRL 110 182001 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 13BB | PRL 111 102003 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| CHATRCHYAN | 13AC | JHEP 1307 163 | S. Chatrchyan <i>et al.</i> | (CMS Collab.) |
| AAIJ | 12AR | JHEP 1210 037 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 12E | PL B708 241 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AALTONEN | 12A | PR D85 032003 | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| ABAZOV | 12U | PR D85 112003 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| AAIJ | 11E | PR D84 092001 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| Also | | PR D85 039904 (errat.) | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AALTONEN | 11 | PRL 106 121804 | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| AALTONEN | 11AI | PRL 107 201802 | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| AALTONEN | 11N | PRL 106 181802 | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| ABAZOV | 11O | PR D84 031102 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| AALTONEN | 10B | PRL 104 102002 | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| AALTONEN | 09C | PRL 103 031801 | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| AALTONEN | 09E | PR D79 032001 | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| ABAZOV | 07S | PRL 99 142001 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| ABAZOV | 07U | PRL 99 182001 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| ABULENCIA | 07A | PRL 98 122001 | A. Abulencia <i>et al.</i> | (FNAL CDF Collab.) |
| ABULENCIA | 07B | PRL 98 122002 | A. Abulencia <i>et al.</i> | (FNAL CDF Collab.) |
| LEITNER | 07 | NPBPS 174 169 | O. Leitner, Z.J. Ajaltouni | |
| ACOSTA | 06 | PRL 96 202001 | D. Acosta <i>et al.</i> | (CDF Collab.) |
| ABAZOV | 05C | PRL 94 102001 | V.M. Abazov <i>et al.</i> | (D0 Collab.) |
| ACOSTA | 05O | PR D72 051104 | D. Acosta <i>et al.</i> | (CDF Collab.) |
| ABDALLAH | 04A | PL B585 63 | J. Abdallah <i>et al.</i> | (DELPHI Collab.) |
| ACOSTA | 02G | PR D66 112002 | D. Acosta <i>et al.</i> | (CDF Collab.) |
| ABREU | 99W | EPJ C10 185 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| ACKERSTAFF | 98G | PL B426 161 | K. Ackerstaff <i>et al.</i> | (OPAL Collab.) |

| | | | | |
|----------|-----|--------------|-----------------------------|---------------------|
| BARATE | 98D | EPJ C2 197 | R. Barate <i>et al.</i> | (ALEPH Collab.) |
| ABE | 97B | PR D55 1142 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ABE | 96M | PRL 77 1439 | F. Abe <i>et al.</i> | (CDF Collab.) |
| ABREU | 96D | ZPHY C71 199 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| ABREU | 96N | PL B374 351 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| ADAM | 96D | ZPHY C72 207 | W. Adam <i>et al.</i> | (DELPHI Collab.) |
| BUSKULIC | 96L | PL B380 442 | D. Buskulic <i>et al.</i> | (ALEPH Collab.) |
| BUSKULIC | 96V | PL B384 471 | D. Buskulic <i>et al.</i> | (ALEPH Collab.) |
| PDG | 96 | PR D54 1 | R. M. Barnett <i>et al.</i> | (PDG Collab.) |
| ABREU | 95S | ZPHY C68 375 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| AKERS | 95K | PL B353 402 | R. Akers <i>et al.</i> | (OPAL Collab.) |
| BUSKULIC | 95L | PL B357 685 | D. Buskulic <i>et al.</i> | (ALEPH Collab.) |
| ABE | 93B | PR D47 2639 | F. Abe <i>et al.</i> | (CDF Collab.) |
| BUSKULIC | 92E | PL B294 145 | D. Buskulic <i>et al.</i> | (ALEPH Collab.) |
| ALBAJAR | 91E | PL B273 540 | C. Albajar <i>et al.</i> | (UA1 Collab.) |
| BARI | 91 | NC 104A 1787 | G. Bari <i>et al.</i> | (CERN R422 Collab.) |
| ARENTON | 86 | NP B274 707 | M.W. Arenton <i>et al.</i> | (ARIZ, NDAM, VAND) |
| BASILE | 81 | LNC 31 97 | M. Basile <i>et al.</i> | (CERN R415 Collab.) |
