



$$I(J^P) = 0(\frac{1}{2}^+) \quad \text{Status: } ****$$

The parity of the Λ_c^+ is defined to be positive (as are the parities of the proton, neutron, and Λ). The quark content is udc . Results of an analysis of $pK^-\pi^+$ decays (JEZABEK 92) are consistent with $J = 1/2$. Nobody doubts that the spin is indeed $1/2$.

We have omitted some results that have been superseded by later experiments. The omitted results may be found in earlier editions.

Λ_c^+ MASS

Our value in 2004, 2284.9 ± 0.6 MeV, was the average of the measurements now filed below as "not used." The BABAR measurement is so much better that we use it alone. Note that it is about 2.6 (old) standard deviations above the 2004 value.

The fit also includes $\Sigma_c - \Lambda_c^+$ and $\Lambda_c^{*+} - \Lambda_c^+$ mass-difference measurements, but this doesn't affect the Λ_c^+ mass. The new (in 2006) Λ_c^+ mass simply pushes all those other masses higher.

| <u>VALUE (MeV)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-------------------------------|-------------|-----------------------|-------------|---|
| 2286.46 ± 0.14 OUR FIT | | | | |
| 2286.46 ± 0.14 | 4891 | ¹ AUBERT,B | 05S BABR | $\Lambda K_S^0 K^+$ and $\Sigma^0 K_S^0 K^+$ |
| • • • | | | | We do not use the following data for averages, fits, limits, etc. • • • |
| 2284.7 ± 0.6 ± 0.7 | 1134 | AVERY | 91 CLEO | Six modes |
| 2281.7 ± 2.7 ± 2.6 | 29 | ALVAREZ | 90B NA14 | $pK^-\pi^+$ |
| 2285.8 ± 0.6 ± 1.2 | 101 | BARLAG | 89 NA32 | $pK^-\pi^+$ |
| 2284.7 ± 2.3 ± 0.5 | 5 | AGUILAR-... | 88B LEBC | $pK^-\pi^+$ |
| 2283.1 ± 1.7 ± 2.0 | 628 | ALBRECHT | 88C ARG | $pK^-\pi^+$, $p\bar{K}^0$, $\Lambda 3\pi$ |
| 2286.2 ± 1.7 ± 0.7 | 97 | ANJOS | 88B E691 | $pK^-\pi^+$ |
| 2281 ± 3 | 2 | JONES | 87 HBC | $pK^-\pi^+$ |
| 2283 ± 3 | 3 | BOSETTI | 82 HBC | $pK^-\pi^+$ |
| 2290 ± 3 | 1 | CALICCHIO | 80 HYBR | $pK^-\pi^+$ |


¹AUBERT,B 05S uses low-Q $\Lambda K_S^0 K^+$ and $\Sigma^0 K_S^0 K^+$ decays to minimize systematic errors. The error above includes systematic as well as statistical errors. Many cross checks and adjustments to properties of the BABAR detector, as well as the large number of clean events, make this by far the best measurement of the Λ_c^+ mass.

Λ_c^+ MEAN LIFE

Measurements with an error $\geq 100 \times 10^{-15}$ s or with fewer than 20 events have been omitted from the Listings.

| <u>VALUE (10^{-15} s)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|-------------|--------------------|-------------|---|
| 200 ± 6 OUR AVERAGE | | Error | | includes scale factor of 1.6. See the ideogram below. |
| 204.6 ± 3.4 ± 2.5 | 8034 | LINK | 02C FOCS | $pK^-\pi^+$ |
| 198.1 ± 7.0 ± 5.6 | 1630 | KUSHNIR... | 01 SELX | $\Lambda_c^+ \rightarrow pK^-\pi^+$ |

| | | | | | |
|---|------|----------|-----|------|--|
| $179.6 \pm 6.9 \pm 4.4$ | 4749 | MAHMOOD | 01 | CLE2 | $e^+e^- \approx \Upsilon(4S)$ |
| $215 \pm 16 \pm 8$ | 1340 | FRABETTI | 93D | E687 | $\gamma \text{Be}, \Lambda_c^+ \rightarrow pK^- \pi^+$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | | |
| $180 \pm 30 \pm 30$ | 29 | ALVAREZ | 90 | NA14 | $\gamma, \Lambda_c^+ \rightarrow pK^- \pi^+$ |
| $200 \pm 30 \pm 30$ | 90 | FRABETTI | 90 | E687 | $\gamma \text{Be}, \Lambda_c^+ \rightarrow pK^- \pi^+$ |
| $196 \begin{smallmatrix} +23 \\ -20 \end{smallmatrix}$ | 101 | BARLAG | 89 | NA32 | $pK^- \pi^+ + \text{c.c.}$ |
| $220 \pm 30 \pm 20$ | 97 | ANJOS | 88B | E691 | $pK^- \pi^+ + \text{c.c.}$ |



Λ_c^+ DECAY MODES

Branching fractions marked with a footnote, e.g. [a], have been corrected for decay modes not observed in the experiments. For example, the sub-mode fraction $\Lambda_c^+ \rightarrow p\bar{K}^*(892)^0$ seen in $\Lambda_c^+ \rightarrow pK^- \pi^+$ has been multiplied up to include $\bar{K}^*(892)^0 \rightarrow \bar{K}^0 \pi^0$ decays.

| Mode | Fraction (Γ_i/Γ) | Scale factor/ Confidence level |
|------|--------------------------------|-----------------------------------|
|------|--------------------------------|-----------------------------------|

Hadronic modes with a p or n : $S = -1$ final states

| | | | |
|---------------|---|----------------------------------|-------|
| Γ_1 | pK_S^0 | (1.59 ± 0.08) % | S=1.1 |
| Γ_2 | $pK^- \pi^+$ | (6.28 ± 0.32) % | S=1.4 |
| Γ_3 | $p\bar{K}^*(892)^0$ | [a] (1.96 ± 0.27) % | |
| Γ_4 | $\Delta(1232)^{++} K^-$ | (1.08 ± 0.25) % | |
| Γ_5 | $\Lambda(1520)\pi^+$ | [a] (2.2 ± 0.5) % | |
| Γ_6 | $pK^- \pi^+$ nonresonant | (3.5 ± 0.4) % | |
| Γ_7 | $pK_S^0 \pi^0$ | (1.97 ± 0.13) % | S=1.1 |
| Γ_8 | $nK_S^0 \pi^+$ | (1.82 ± 0.25) % | |
| Γ_9 | $p\bar{K}^0 \eta$ | (1.6 ± 0.4) % | |
| Γ_{10} | $pK_S^0 \pi^+ \pi^-$ | (1.60 ± 0.12) % | S=1.1 |
| Γ_{11} | $pK^- \pi^+ \pi^0$ | (4.46 ± 0.30) % | S=1.5 |
| Γ_{12} | $pK^*(892)^- \pi^+$ | [a] (1.4 ± 0.5) % | |
| Γ_{13} | $p(K^- \pi^+)_{\text{nonresonant}} \pi^0$ | (4.6 ± 0.8) % | |
| Γ_{14} | $\Delta(1232)\bar{K}^*(892)$ | seen | |
| Γ_{15} | $pK^- 2\pi^+ \pi^-$ | (1.4 ± 0.9) × 10 ⁻³ | |
| Γ_{16} | $pK^- \pi^+ 2\pi^0$ | (1.0 ± 0.5) % | |

Hadronic modes with a p : $S = 0$ final states

| | | | |
|---------------|------------------------------|--|--------|
| Γ_{17} | $p\pi^0$ | < 2.7 × 10 ⁻⁴ | CL=90% |
| Γ_{18} | $p\eta$ | (1.24 ± 0.30) × 10 ⁻³ | |
| Γ_{19} | $p\omega(782)^0$ | (9 ± 4) × 10 ⁻⁴ | |
| Γ_{20} | $p\pi^+ \pi^-$ | (4.61 ± 0.28) × 10 ⁻³ | |
| Γ_{21} | $pf_0(980)$ | [a] (3.5 ± 2.3) × 10 ⁻³ | |
| Γ_{22} | $p2\pi^+ 2\pi^-$ | (2.3 ± 1.4) × 10 ⁻³ | |
| Γ_{23} | $pK^+ K^-$ | (1.06 ± 0.06) × 10 ⁻³ | |
| Γ_{24} | $p\phi$ | [a] (1.06 ± 0.14) × 10 ⁻³ | |
| Γ_{25} | $pK^+ K^-$ non- ϕ | (5.3 ± 1.2) × 10 ⁻⁴ | |
| Γ_{26} | $p\phi\pi^0$ | (10 ± 4) × 10 ⁻⁵ | |
| Γ_{27} | $pK^+ K^- \pi^0$ nonresonant | < 6.3 × 10 ⁻⁵ | CL=90% |

Hadronic modes with a hyperon: $S = -1$ final states

| | | | |
|---------------|--|----------------------------------|--------|
| Γ_{28} | $\Lambda\pi^+$ | (1.30 ± 0.07) % | S=1.1 |
| Γ_{29} | $\Lambda\pi^+ \pi^0$ | (7.1 ± 0.4) % | S=1.1 |
| Γ_{30} | $\Lambda\rho^+$ | < 6 % | CL=95% |
| Γ_{31} | $\Lambda\pi^- 2\pi^+$ | (3.64 ± 0.29) % | S=1.4 |
| Γ_{32} | $\Sigma(1385)^+ \pi^+ \pi^-$, $\Sigma^{*+} \rightarrow$ $\Lambda\pi^+$ | (1.0 ± 0.5) % | |
| Γ_{33} | $\Sigma(1385)^- 2\pi^+$, $\Sigma^{*-} \rightarrow$ $\Lambda\pi^-$ | (7.6 ± 1.4) × 10 ⁻³ | |
| Γ_{34} | $\Lambda\pi^+ \rho^0$ | (1.5 ± 0.6) % | |
| Γ_{35} | $\Sigma(1385)^+ \rho^0$, $\Sigma^{*+} \rightarrow$ $\Lambda\pi^+$ | (5 ± 4) × 10 ⁻³ | |
| Γ_{36} | $\Lambda\pi^- 2\pi^+$ nonresonant | < 1.1 % | CL=90% |
| Γ_{37} | $\Lambda\pi^- \pi^0 2\pi^+$ total | (2.3 ± 0.8) % | |
| Γ_{38} | $\Lambda\pi^+ \eta$ | [a] (2.2 ± 0.5) % | |
| Γ_{39} | $\Sigma(1385)^+ \eta$ | [a] (1.07 ± 0.32) % | |

| | | | | |
|---------------|---|-----|----------------------------------|--------|
| Γ_{40} | $\Lambda\pi^+\omega$ | [a] | $(1.5 \pm 0.5) \%$ | |
| Γ_{41} | $\Lambda\pi^-\pi^0 2\pi^+$, no η or ω | | $< 8 \times 10^{-3}$ | CL=90% |
| Γ_{42} | $\Lambda K^+\bar{K}^0$ | | $(5.7 \pm 1.1) \times 10^{-3}$ | S=1.9 |
| Γ_{43} | $\Xi(1690)^0 K^+$, $\Xi^{*0} \rightarrow \Lambda\bar{K}^0$ | | $(1.6 \pm 0.5) \times 10^{-3}$ | |
| Γ_{44} | $\Sigma^0\pi^+$ | | $(1.29 \pm 0.07) \%$ | S=1.1 |
| Γ_{45} | $\Sigma^+\pi^0$ | | $(1.25 \pm 0.10) \%$ | |
| Γ_{46} | $\Sigma^+\eta$ | | $(6.9 \pm 2.3) \times 10^{-3}$ | |
| Γ_{47} | $\Sigma^+\pi^+\pi^-$ | | $(4.50 \pm 0.25) \%$ | S=1.3 |
| Γ_{48} | $\Sigma^+\rho^0$ | | $< 1.7 \%$ | CL=95% |
| Γ_{49} | $\Sigma^- 2\pi^+$ | | $(1.87 \pm 0.18) \%$ | |
| Γ_{50} | $\Sigma^0\pi^+\pi^0$ | | $(3.5 \pm 0.4) \%$ | |
| Γ_{51} | $\Sigma^+\pi^0\pi^0$ | | $(1.55 \pm 0.15) \%$ | |
| Γ_{52} | $\Sigma^0\pi^- 2\pi^+$ | | $(1.11 \pm 0.30) \%$ | |
| Γ_{53} | $\Sigma^+\pi^+\pi^-\pi^0$ | | — | |
| Γ_{54} | $\Sigma^+\omega$ | [a] | $(1.70 \pm 0.21) \%$ | |
| Γ_{55} | $\Sigma^-\pi^0 2\pi^+$ | | $(2.1 \pm 0.4) \%$ | |
| Γ_{56} | $\Sigma^+ K^+ K^-$ | | $(3.5 \pm 0.4) \times 10^{-3}$ | S=1.1 |
| Γ_{57} | $\Sigma^+\phi$ | [a] | $(3.9 \pm 0.6) \times 10^{-3}$ | S=1.1 |
| Γ_{58} | $\Xi(1690)^0 K^+$, $\Xi^{*0} \rightarrow \Sigma^+ K^-$ | | $(1.02 \pm 0.25) \times 10^{-3}$ | |
| Γ_{59} | $\Sigma^+ K^+ K^-$ nonresonant | | $< 8 \times 10^{-4}$ | CL=90% |
| Γ_{60} | $\Xi^0 K^+$ | | $(5.5 \pm 0.7) \times 10^{-3}$ | |
| Γ_{61} | $\Xi^- K^+ \pi^+$ | | $(6.2 \pm 0.6) \times 10^{-3}$ | S=1.1 |
| Γ_{62} | $\Xi(1530)^0 K^+$ | | $(4.3 \pm 0.9) \times 10^{-3}$ | S=1.1 |

Hadronic modes with a hyperon: $S = 0$ final states

| | | | | |
|---------------|----------------------------|-----|--------------------------------|--------|
| Γ_{63} | ΛK^+ | | $(6.1 \pm 1.2) \times 10^{-4}$ | |
| Γ_{64} | $\Lambda K^+ \pi^+ \pi^-$ | | $< 5 \times 10^{-4}$ | CL=90% |
| Γ_{65} | $\Sigma^0 K^+$ | | $(5.2 \pm 0.8) \times 10^{-4}$ | |
| Γ_{66} | $\Sigma^0 K^+ \pi^+ \pi^-$ | | $< 2.6 \times 10^{-4}$ | CL=90% |
| Γ_{67} | $\Sigma^+ K^+ \pi^-$ | | $(2.1 \pm 0.6) \times 10^{-3}$ | |
| Γ_{68} | $\Sigma^+ K^*(892)^0$ | [a] | $(3.5 \pm 1.0) \times 10^{-3}$ | |
| Γ_{69} | $\Sigma^- K^+ \pi^+$ | | $< 1.2 \times 10^{-3}$ | CL=90% |

Doubly Cabibbo-suppressed modes

| | | | | |
|---------------|------------------|--|----------------------------------|--|
| Γ_{70} | $\rho K^+ \pi^-$ | | $(1.11 \pm 0.18) \times 10^{-4}$ | |
|---------------|------------------|--|----------------------------------|--|

Semileptonic modes

| | | | | |
|---------------|-------------------------|--|--------------------|--|
| Γ_{71} | $\Lambda e^+ \nu_e$ | | $(3.6 \pm 0.4) \%$ | |
| Γ_{72} | $\Lambda \mu^+ \nu_\mu$ | | $(3.5 \pm 0.5) \%$ | |

Inclusive modes

| | | |
|---------------|--------------------|---------------------------|
| Γ_{73} | e^+ anything | $(3.95 \pm 0.35) \%$ |
| Γ_{74} | p anything | $(50 \pm 16) \%$ |
| Γ_{75} | n anything | $(50 \pm 16) \%$ |
| Γ_{76} | Λ anything | $(38.2^{+2.9}_{-2.4}) \%$ |
| Γ_{77} | 3prongs | $(24 \pm 8) \%$ |

**$\Delta C = 1$ weak neutral current ($C1$) modes, or
Lepton Family number (LF), or Lepton number (L), or
Baryon number (B) violating modes**

| | | | | | |
|---------------|------------------------------|--------|---------|------------------|--------|
| Γ_{78} | $p e^+ e^-$ | $C1$ | < 5.5 | $\times 10^{-6}$ | CL=90% |
| Γ_{79} | $p \mu^+ \mu^-$ non-resonant | $C1$ | < 7.7 | $\times 10^{-8}$ | CL=90% |
| Γ_{80} | $p e^+ \mu^-$ | LF | < 9.9 | $\times 10^{-6}$ | CL=90% |
| Γ_{81} | $p e^- \mu^+$ | LF | < 1.9 | $\times 10^{-5}$ | CL=90% |
| Γ_{82} | $\bar{p} 2e^+$ | L, B | < 2.7 | $\times 10^{-6}$ | CL=90% |
| Γ_{83} | $\bar{p} 2\mu^+$ | L, B | < 9.4 | $\times 10^{-6}$ | CL=90% |
| Γ_{84} | $\bar{p} e^+ \mu^+$ | L, B | < 1.6 | $\times 10^{-5}$ | CL=90% |
| Γ_{85} | $\Sigma^- \mu^+ \mu^+$ | L | < 7.0 | $\times 10^{-4}$ | CL=90% |

[a] This branching fraction includes all the decay modes of the final-state resonance.

CONSTRAINED FIT INFORMATION

An overall fit to 41 branching ratios uses 62 measurements and one constraint to determine 21 parameters. The overall fit has a $\chi^2 = 47.4$ for 42 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

| | | | | | | | | | | | |
|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|--|
| x_2 | 54 | | | | | | | | | | |
| x_7 | 46 | 55 | | | | | | | | | |
| x_{10} | 44 | 64 | 39 | | | | | | | | |
| x_{11} | 51 | 61 | 40 | 60 | | | | | | | |
| x_{28} | 54 | 66 | 44 | 42 | 43 | | | | | | |
| x_{29} | 45 | 61 | 41 | 38 | 36 | 65 | | | | | |
| x_{31} | 51 | 37 | 28 | 41 | 60 | 45 | 36 | | | | |
| x_{42} | 16 | 22 | 14 | 14 | 14 | 26 | 19 | 12 | | | |
| x_{44} | 51 | 55 | 38 | 37 | 40 | 74 | 58 | 44 | 20 | | |
| x_{45} | 38 | 39 | 30 | 25 | 29 | 34 | 33 | 23 | 10 | 29 | |
| x_{47} | 51 | 88 | 50 | 60 | 61 | 59 | 55 | 39 | 20 | 50 | |
| x_{49} | 5 | 9 | 5 | 6 | 6 | 6 | 6 | 3 | 2 | 5 | |
| x_{52} | 13 | 14 | 9 | 12 | 15 | 13 | 11 | 20 | 4 | 12 | |
| x_{54} | 19 | 30 | 18 | 23 | 26 | 19 | 18 | 18 | 6 | 16 | |
| x_{56} | 23 | 41 | 23 | 28 | 28 | 27 | 25 | 18 | 9 | 23 | |
| x_{57} | 19 | 32 | 19 | 22 | 23 | 22 | 20 | 14 | 7 | 18 | |
| x_{60} | 8 | 15 | 8 | 10 | 9 | 10 | 9 | 6 | 3 | 8 | |
| x_{61} | 29 | 39 | 25 | 25 | 25 | 51 | 35 | 24 | 14 | 38 | |
| x_{62} | 6 | 11 | 6 | 7 | 7 | 7 | 7 | 4 | 2 | 6 | |
| | x_1 | x_2 | x_7 | x_{10} | x_{11} | x_{28} | x_{29} | x_{31} | x_{42} | x_{44} | |
| x_{47} | 36 | | | | | | | | | | |
| x_{49} | 4 | 8 | | | | | | | | | |
| x_{52} | 7 | 14 | 1 | | | | | | | | |
| x_{54} | 14 | 29 | 3 | 5 | | | | | | | |
| x_{56} | 17 | 45 | 4 | 6 | 13 | | | | | | |
| x_{57} | 13 | 37 | 3 | 5 | 11 | 16 | | | | | |
| x_{60} | 6 | 13 | 1 | 2 | 5 | 6 | 5 | | | | |
| x_{61} | 19 | 34 | 4 | 7 | 11 | 16 | 13 | 6 | | | |
| x_{62} | 4 | 10 | 1 | 2 | 3 | 4 | 4 | 2 | 4 | | |
| | x_{45} | x_{47} | x_{49} | x_{52} | x_{54} | x_{56} | x_{57} | x_{60} | x_{61} | | |

Λ_c^+ BRANCHING RATIOS

A few really obsolete results have been omitted.

————— Hadronic modes with a p : $S = -1$ final states —————

| $\Gamma(pK_S^0)/\Gamma_{\text{total}}$ | | | | | Γ_1/Γ |
|--|-------------|--------------------|-------------|----------------|--|
| <u>VALUE (%)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | |
| 1.59±0.08 OUR FIT Error includes scale factor of 1.1. | | | | | |
| 1.52±0.08±0.03 | 1243 | ABLIKIM | 16 | BES3 | $e^+e^- \rightarrow \Lambda_c \bar{\Lambda}_c$, 4.599 GeV |

| $\Gamma(pK_S^0)/\Gamma(pK^-\pi^+)$ | | | | | Γ_1/Γ_2 |
|---|-------------|--------------------|-------------|----------------|-------------------------------|
| Measurements given as a \bar{K}^0 ratio have been divided by 2 to convert to a K_S^0 ratio. | | | | | |
| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | |
| 0.254±0.012 OUR FIT Error includes scale factor of 1.4. | | | | | |
| 0.234±0.020 OUR AVERAGE | | | | | |
| 0.23 ±0.01 ±0.02 | 1025 | ALAM | 98 | CLE2 | $e^+e^- \approx \Upsilon(4S)$ |
| 0.22 ±0.04 ±0.03 | 133 | AVERY | 91 | CLEO | e^+e^- 10.5 GeV |
| 0.28 ±0.09 ±0.07 | 45 | ANJOS | 90 | E691 | γ Be 70–260 GeV |
| 0.31 ±0.08 ±0.02 | 73 | ALBRECHT | 88C | ARG | e^+e^- 10 GeV |

| $\Gamma(pK^-\pi^+)/\Gamma_{\text{total}}$ | | | | | Γ_2/Γ |
|---|-------------|---------------------|-------------|----------------|--|
| <u>VALUE (%)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | |
| 6.28±0.32 OUR FIT Error includes scale factor of 1.4. | | | | | |
| 6.3 ±0.5 OUR AVERAGE Error includes scale factor of 2.0. | | | | | |
| 5.84±0.27±0.23 | 6.3k | ABLIKIM | 16 | BES3 | $e^+e^- \rightarrow \Lambda_c \bar{\Lambda}_c$, 4.599 GeV |
| 6.84±0.24 ^{+0.21} _{-0.27} | 1.4k | ¹ ZUPANC | 14 | BELL | $e^+e^- \rightarrow D^{(*)-} \bar{p}\pi^+$ recoil |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| 5.0 ±1.3 | | ² PDG | 02 | | See footnote |

¹This ZUPANC 14 value is the FIRST-EVER model-independent measurement of a Λ_c^+ branching fraction.

²See the note by P. Burchat, " Λ_c^+ Branching Fractions," in any edition of the Review from 2002 through 2014 for how this value was obtained. It is now obsolete.

| $\Gamma(p\bar{K}^*(892)^0)/\Gamma(pK^-\pi^+)$ | | | | | Γ_3/Γ_2 |
|---|-------------|---------------------|-------------|----------------|------------------------------------|
| Unseen decay modes of the $\bar{K}^*(892)^0$ are included. | | | | | |
| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | |
| 0.31±0.04 OUR AVERAGE | | | | | |
| 0.29±0.04±0.03 | | ¹ AITALA | 00 | E791 | $\pi^- N$, 500 GeV |
| 0.35 ^{+0.06} _{-0.07} ±0.03 | 39 | BOZEK | 93 | NA32 | π^- Cu 230 GeV |
| 0.42±0.24 | 12 | BASILE | 81B | CNTR | $pp \rightarrow \Lambda_c^+ e^- X$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| 0.35±0.11 | | BARLAG | 90D | NA32 | See BOZEK 93 |

¹AITALA 00 makes a coherent 5-dimensional amplitude analysis of $946 \pm 38 \Lambda_c^+ \rightarrow pK^-\pi^+$ decays.

$\Gamma(\Delta(1232)^{++}K^-)/\Gamma(\rho K^- \pi^+)$ Γ_4/Γ_2

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|---------------------|----------|-------------------------------------|
| 0.17±0.04 OUR AVERAGE | | | | Error includes scale factor of 1.1. |
| 0.18±0.03±0.03 | | ¹ AITALA | 00 E791 | $\pi^- N$, 500 GeV |
| 0.12 ^{+0.04} _{-0.05} ±0.05 | 14 | BOZEK | 93 NA32 | π^- Cu 230 GeV |
| 0.40±0.17 | 17 | BASILE | 81B CNTR | $pp \rightarrow \Lambda_c^+ e^- X$ |

¹ AITALA 00 makes a coherent 5-dimensional amplitude analysis of $946 \pm 38 \Lambda_c^+ \rightarrow \rho K^- \pi^+$ decays.

$\Gamma(\Lambda(1520)\pi^+)/\Gamma(\rho K^- \pi^+)$ Γ_5/Γ_2

Unseen decay modes of the $\Lambda(1520)$ are included.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|---------------------|---------|---------------------|
| 0.35±0.08 OUR AVERAGE | | | | |
| 0.34±0.08±0.05 | | ¹ AITALA | 00 E791 | $\pi^- N$, 500 GeV |
| 0.40 ^{+0.18} _{-0.13} ±0.09 | 12 | BOZEK | 93 NA32 | π^- Cu 230 GeV |

¹ AITALA 00 makes a coherent 5-dimensional amplitude analysis of $946 \pm 38 \Lambda_c^+ \rightarrow \rho K^- \pi^+$ decays.

$\Gamma(\rho K^- \pi^+ \text{ nonresonant})/\Gamma(\rho K^- \pi^+)$ Γ_6/Γ_2

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|---------------------|---------|---------------------|
| 0.55±0.06 OUR AVERAGE | | | | |
| 0.55±0.06±0.04 | | ¹ AITALA | 00 E791 | $\pi^- N$, 500 GeV |
| 0.56 ^{+0.07} _{-0.09} ±0.05 | 71 | BOZEK | 93 NA32 | π^- Cu 230 GeV |

¹ AITALA 00 makes a coherent 5-dimensional amplitude analysis of $946 \pm 38 \Lambda_c^+ \rightarrow \rho K^- \pi^+$ decays.

$\Gamma(\rho K_S^0 \pi^0)/\Gamma_{\text{total}}$ Γ_7/Γ

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|---------|---|
| 1.97±0.13 OUR FIT | | | | Error includes scale factor of 1.1. |
| 1.87±0.13±0.05 | 558 | ABLIKIM | 16 BES3 | $e^+ e^- \rightarrow \Lambda_c \bar{\Lambda}_c$, 4.599 GeV |

$\Gamma(\rho K_S^0 \pi^0)/\Gamma(\rho K^- \pi^+)$ Γ_7/Γ_2

Measurements given as a \bar{K}^0 ratio have been divided by 2 to convert to a K_S^0 ratio.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------------|------|-------------|---------|--------------------------------|
| 0.314±0.018 OUR FIT | | | | |
| 0.33 ±0.03 ±0.04 | 774 | ALAM | 98 CLE2 | $e^+ e^- \approx \Upsilon(4S)$ |

$\Gamma(n K_S^0 \pi^+)/\Gamma_{\text{total}}$ Γ_8/Γ

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------|------|-------------|----------|----------------------|
| 1.82±0.23±0.11 | 83 | ABLIKIM | 17H BES3 | $e^+ e^-$ at 4.6 GeV |

$\Gamma(\rho \bar{K}^0 \eta)/\Gamma(\rho K^- \pi^+)$ Γ_9/Γ_2

Unseen decay modes of the η are included.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------|------|-------------|---------|--------------------------------|
| 0.25±0.04±0.04 | 57 | AMMAR | 95 CLE2 | $e^+ e^- \approx \Upsilon(4S)$ |

$\Gamma(\rho K_S^0 \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{10} / Γ

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------------|------|-------------|------|--|
| 1.60 ± 0.12 OUR FIT | | | | Error includes scale factor of 1.1. |
| 1.53 ± 0.11 ± 0.09 | 485 | ABLIKIM | 16 | BES3 $e^+ e^- \rightarrow \Lambda_c \bar{\Lambda}_c$, 4.599 GeV |

$\Gamma(\rho K_S^0 \pi^+ \pi^-) / \Gamma(\rho K^- \pi^+)$ Γ_{10} / Γ_2
 Measurements given as a \bar{K}^0 ratio have been divided by 2 to convert to a K_S^0 ratio.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------------------|------|-------------|------|-------------------------------------|
| 0.255 ± 0.015 OUR FIT | | | | Error includes scale factor of 1.1. |
| 0.257 ± 0.031 OUR AVERAGE | | | | |
| 0.26 ± 0.02 ± 0.03 | 985 | ALAM | 98 | CLE2 $e^+ e^- \approx \Upsilon(4S)$ |
| 0.22 ± 0.06 ± 0.02 | 83 | AVERY | 91 | CLEO $e^+ e^-$ 10.5 GeV |
| 0.49 ± 0.18 ± 0.04 | 12 | BARLAG | 90D | NA32 π^- 230 GeV |

$\Gamma(\rho K^- \pi^+ \pi^0) / \Gamma_{\text{total}}$ Γ_{11} / Γ

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------------|------|-------------|------|--|
| 4.46 ± 0.30 OUR FIT | | | | Error includes scale factor of 1.5. |
| 4.53 ± 0.23 ± 0.30 | 1849 | ABLIKIM | 16 | BES3 $e^+ e^- \rightarrow \Lambda_c \bar{\Lambda}_c$, 4.599 GeV |

$\Gamma(\rho K^- \pi^+ \pi^0) / \Gamma(\rho K^- \pi^+)$ Γ_{11} / Γ_2

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------------------|------|-------------|------|---|
| 0.71 ± 0.04 OUR FIT | | | | Error includes scale factor of 2.4. |
| 0.685 ± 0.019 OUR AVERAGE | | | | |
| 0.685 ± 0.007 ± 0.018 | 242k | PAL | 17 | BELL $e^+ e^- \approx \Upsilon(4S), \Upsilon(5S)$ |
| 0.67 ± 0.04 ± 0.11 | 2.6k | ALAM | 98 | CLE2 $e^+ e^- \approx \Upsilon(4S)$ |

$\Gamma(\rho K^*(892)^- \pi^+) / \Gamma(\rho K_S^0 \pi^+ \pi^-)$ $\Gamma_{12} / \Gamma_{10}$

Unseen decay modes of the $K^*(892)^-$ are included.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------|------|-------------|------|---------------------|
| 0.88 ± 0.28 | 17 | ALEEV | 94 | BIS2 nN 20–70 GeV |

$\Gamma(\rho(K^- \pi^+)_{\text{nonresonant}} \pi^0) / \Gamma(\rho K^- \pi^+)$ Γ_{13} / Γ_2

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------------|------|-------------|------|-------------------------|
| 0.73 ± 0.12 ± 0.05 | 67 | BOZEK | 93 | NA32 π^- Cu 230 GeV |

$\Gamma(\Delta(1232) \bar{K}^*(892)) / \Gamma_{\text{total}}$ Γ_{14} / Γ

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------------|------|-------------|------|---------------------|
| seen | 35 | AMENDOLIA | 87 | SPEC γ Ge-Si |

$\Gamma(\rho K^- 2\pi^+ \pi^-) / \Gamma(\rho K^- \pi^+)$ Γ_{15} / Γ_2

| VALUE | DOCUMENT ID | TECN | COMMENT |
|----------------------|-------------|------|----------------------|
| 0.022 ± 0.015 | BARLAG | 90D | NA32 π^- 230 GeV |

$\Gamma(\rho K^- \pi^+ 2\pi^0) / \Gamma(\rho K^- \pi^+)$ Γ_{16} / Γ_2

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------------|------|-------------|------|-------------------------|
| 0.16 ± 0.07 ± 0.03 | 15 | BOZEK | 93 | NA32 π^- Cu 230 GeV |

————— Hadronic modes with a p : $S = 0$ final states —————

$\Gamma(p\pi^0)/\Gamma_{\text{total}}$ Γ_{17}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-----|-------------|------|--------------------------|
| $<2.7 \times 10^{-4}$ | 90 | ABLIKIM | 17Q | BES3 e^+e^- at 4.6 GeV |

$\Gamma(p\eta)/\Gamma_{\text{total}}$ Γ_{18}/Γ

Unseen decay modes of the η are included.

| VALUE (units 10^{-3}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|--|
| $1.24 \pm 0.28 \pm 0.10$ | 52 | ABLIKIM | 17Q | BES3 $\eta \rightarrow 2\gamma, \pi^+\pi^0\pi^-$ |

$\Gamma(p\omega(782)^0)/\Gamma_{\text{total}}$ Γ_{19}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|--|
| $9.4 \pm 3.2 \pm 2.2$ | 13 | AAIJ | 18N | LHCB Seen in $\Lambda_c^+ \rightarrow p\mu^+\mu^-$ |

$\Gamma(p\pi^+\pi^-)/\Gamma(pK^-\pi^+)$ Γ_{20}/Γ_2

| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|------|--|
| 7.35 ± 0.24 OUR AVERAGE | | | | Error includes scale factor of 1.3. |
| $7.44 \pm 0.08 \pm 0.18$ | 20k | AAIJ | 18V | LHCB $\Lambda_b^0 \rightarrow \Lambda_c^+\mu^-X$ |
| $6.70 \pm 0.48 \pm 0.25$ | 495 | ABLIKIM | 16U | BES3 e^+e^- at 4.599 GeV |
| 6.9 ± 3.6 | 5 | BARLAG | 90D | NA32 π^- 230 GeV |

$\Gamma(pf_0(980))/\Gamma(pK^-\pi^+)$ Γ_{21}/Γ_2

Unseen decay modes of the $f_0(980)$ are included.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------------------------------------|-------------|------|----------------------|
| 0.055 ± 0.036 | BARLAG | 90D | NA32 π^- 230 GeV |

$\Gamma(p2\pi^+2\pi^-)/\Gamma(pK^-\pi^+)$ Γ_{22}/Γ_2

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------------------------------------|-------------|------|----------------------|
| 0.036 ± 0.023 | BARLAG | 90D | NA32 π^- 230 GeV |

$\Gamma(pK^+K^-)/\Gamma(pK^-\pi^+)$ Γ_{23}/Γ_2

| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|------|--|
| 1.70 ± 0.04 OUR AVERAGE | | | | |
| $1.70 \pm 0.03 \pm 0.03$ | 3.4k | AAIJ | 18V | LHCB $\Lambda_b^0 \rightarrow \Lambda_c^+\mu^-X$ |
| $1.4 \pm 0.2 \pm 0.2$ | 676 | ABE | 02C | BELL $e^+e^- \approx \Upsilon(4S)$ |
| $3.9 \pm 0.9 \pm 0.7$ | 214 | ALEXANDER | 96C | CLE2 $e^+e^- \approx \Upsilon(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| $9.6 \pm 2.9 \pm 1.0$ | 30 | FRABETTI | 93H | E687 $\gamma\text{Be}, \bar{E}_\gamma$ 220 GeV |
| 4.8 ± 2.7 | | BARLAG | 90D | NA32 π^- 230 GeV |

$\Gamma(p\phi)/\Gamma(pK^-\pi^+)$ Γ_{24}/Γ_2

Unseen decay modes of the ϕ are included.

| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|------|------------------------------------|
| 1.70 ± 0.21 OUR AVERAGE | | | | |
| $1.81 \pm 0.33 \pm 0.13$ | 44 | ABLIKIM | 16U | BES3 e^+e^- at 4.599 GeV |
| $1.5 \pm 0.2 \pm 0.2$ | 345 | ABE | 02C | BELL $e^+e^- \approx \Upsilon(4S)$ |
| $2.4 \pm 0.6 \pm 0.3$ | 54 | ALEXANDER | 96C | CLE2 $e^+e^- \approx \Upsilon(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 4.0 ± 2.7 | | BARLAG | 90D | NA32 π^- 230 GeV |

$\Gamma(pK^+K^- \text{ non-}\phi)/\Gamma(pK^-\pi^+)$ Γ_{25}/Γ_2

| <u>VALUE (units 10^{-3})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|-----------------------------|
| 8.4 ± 1.8 OUR AVERAGE | | | | |
| 9.36 ± 2.22 ± 0.71 | 38 | ABLIKIM | 16U BES3 | e^+e^- at 4.599 GeV |
| 7 ± 2 ± 2 | 344 | ABE | 02C BELL | $e^+e^- \approx \gamma(4S)$ |

$\Gamma(p\phi\pi^0)/\Gamma(pK^-\pi^+)$ Γ_{26}/Γ_2

| <u>VALUE (units 10^{-3})</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|--------------------|-------------|---|
| 1.538 ± 0.641 ^{+0.077}_{-0.100} | PAL | 17 BELL | $e^+e^- \approx \gamma(4S), \gamma(5S)$ |

$\Gamma(pK^+K^-\pi^0 \text{ nonresonant})/\Gamma_{\text{total}}$ Γ_{27}/Γ

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------------------|------------|--------------------|-------------|---|
| < 6.3 × 10⁻⁵ | 90 | PAL | 17 BELL | $e^+e^- \approx \gamma(4S), \gamma(5S)$ |

————— Hadronic modes with a hyperon: $S = -1$ final states —————

$\Gamma(\Lambda\pi^+)/\Gamma_{\text{total}}$ Γ_{28}/Γ

| <u>VALUE (%)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------------------|-------------|--------------------|-------------|--|
| 1.30 ± 0.07 OUR FIT | | | | Error includes scale factor of 1.1. |
| 1.24 ± 0.07 ± 0.03 | 706 | ABLIKIM | 16 BES3 | $e^+e^- \rightarrow \Lambda_c \bar{\Lambda}_c$, 4.599 GeV |

$\Gamma(\Lambda\pi^+)/\Gamma(pK^-\pi^+)$ Γ_{28}/Γ_2

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------------------------|-------------|--------------------|-------------|--|
| 0.207 ± 0.009 OUR FIT | | | | Error includes scale factor of 1.2. |
| 0.204 ± 0.019 OUR AVERAGE | | | | |
| 0.217 ± 0.013 ± 0.020 | 750 | LINK | 05F FOCS | γ nucleus, $\bar{E}_\gamma \approx 180$ GeV |
| 0.18 ± 0.03 ± 0.04 | | ALBRECHT | 92 ARG | $e^+e^- \approx 10.4$ GeV |
| 0.18 ± 0.03 ± 0.03 | 87 | AVERY | 91 CLEO | e^+e^- 10.5 GeV |

$\Gamma(\Lambda\pi^+\pi^0)/\Gamma_{\text{total}}$ Γ_{29}/Γ

| <u>VALUE (%)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------|-------------|--------------------|-------------|--|
| 7.1 ± 0.4 OUR FIT | | | | Error includes scale factor of 1.1. |
| 7.01 ± 0.37 ± 0.19 | 1497 | ABLIKIM | 16 BES3 | $e^+e^- \rightarrow \Lambda_c \bar{\Lambda}_c$, 4.599 GeV |

$\Gamma(\Lambda\pi^+\pi^0)/\Gamma(pK^-\pi^+)$ Γ_{29}/Γ_2

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------------------|-------------|--------------------|-------------|---|
| 1.12 ± 0.05 OUR FIT | | | | Error includes scale factor of 1.1. |
| 0.73 ± 0.09 ± 0.16 | 464 | AVERY | 94 CLE2 | $e^+e^- \approx \gamma(3S), \gamma(4S)$ |

$\Gamma(\Lambda\rho^+)/\Gamma(pK^-\pi^+)$ Γ_{30}/Γ_2

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|------------------|------------|--------------------|-------------|---|
| < 0.95 | 95 | AVERY | 94 CLE2 | $e^+e^- \approx \gamma(3S), \gamma(4S)$ |

$\Gamma(\Lambda\pi^-2\pi^+)/\Gamma_{\text{total}}$ Γ_{31}/Γ

| <u>VALUE (%)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------------------|-------------|--------------------|-------------|--|
| 3.64 ± 0.29 OUR FIT | | | | Error includes scale factor of 1.4. |
| 3.81 ± 0.24 ± 0.18 | 609 | ABLIKIM | 16 BES3 | $e^+e^- \rightarrow \Lambda_c \bar{\Lambda}_c$, 4.599 GeV |

$\Gamma(\Lambda\pi^- 2\pi^+)/\Gamma(\rho K^- \pi^+)$ Γ_{31}/Γ_2

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------------------------|-------------|--------------------|-------------|--|
| 0.58 ± 0.04 OUR FIT | | | | Error includes scale factor of 1.9. |
| 0.522 ± 0.032 OUR AVERAGE | | | | |
| 0.508 ± 0.024 ± 0.024 | 1356 | LINK | 05F FOCS | γ nucleus, $\bar{E}_\gamma \approx 180$ GeV |
| 0.65 ± 0.11 ± 0.12 | 289 | AVERY | 91 CLEO | $e^+ e^-$ 10.5 GeV |
| 0.82 ± 0.29 ± 0.27 | 44 | ANJOS | 90 E691 | γ Be 70–260 GeV |
| 0.94 ± 0.41 ± 0.13 | 10 | BARLAG | 90D NA32 | π^- 230 GeV |
| 0.61 ± 0.16 ± 0.04 | 105 | ALBRECHT | 88C ARG | $e^+ e^-$ 10 GeV |

$\Gamma(\Sigma(1385)^+ \pi^+ \pi^-, \Sigma^{*+} \rightarrow \Lambda\pi^+)/\Gamma(\Lambda\pi^- 2\pi^+)$ Γ_{32}/Γ_{31}

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------|--------------------|-------------|--|
| 0.28 ± 0.10 ± 0.08 | LINK | 05F FOCS | γ nucleus, $\bar{E}_\gamma \approx 180$ GeV |

$\Gamma(\Sigma(1385)^- 2\pi^+, \Sigma^{*-} \rightarrow \Lambda\pi^-)/\Gamma(\Lambda\pi^- 2\pi^+)$ Γ_{33}/Γ_{31}

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------|--------------------|-------------|--|
| 0.21 ± 0.03 ± 0.02 | LINK | 05F FOCS | γ nucleus, $\bar{E}_\gamma \approx 180$ GeV |

$\Gamma(\Lambda\pi^+ \rho^0)/\Gamma(\Lambda\pi^- 2\pi^+)$ Γ_{34}/Γ_{31}

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------|--------------------|-------------|--|
| 0.40 ± 0.12 ± 0.12 | LINK | 05F FOCS | γ nucleus, $\bar{E}_\gamma \approx 180$ GeV |

$\Gamma(\Sigma(1385)^+ \rho^0, \Sigma^{*+} \rightarrow \Lambda\pi^+)/\Gamma(\Lambda\pi^- 2\pi^+)$ Γ_{35}/Γ_{31}

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------|--------------------|-------------|--|
| 0.14 ± 0.09 ± 0.07 | LINK | 05F FOCS | γ nucleus, $\bar{E}_\gamma \approx 180$ GeV |

$\Gamma(\Lambda\pi^- 2\pi^+ \text{ nonresonant})/\Gamma(\Lambda\pi^- 2\pi^+)$ Γ_{36}/Γ_{31}

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------|------------|--------------------|-------------|--|
| < 0.3 | 90 | LINK | 05F FOCS | γ nucleus, $\bar{E}_\gamma \approx 180$ GeV |

$\Gamma(\Lambda\pi^- \pi^0 2\pi^+ \text{ total})/\Gamma(\rho K^- \pi^+)$ Γ_{37}/Γ_2

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------|-------------|-----------------------------|-------------|--------------------------------|
| 0.36 ± 0.09 ± 0.09 | 50 | ¹ CRONIN-HEN..03 | CLE3 | $e^+ e^- \approx \Upsilon(4S)$ |

¹ CRONIN-HENNESSY 03 finds this channel to be dominantly $\Lambda\eta\pi^+$ and $\Lambda\omega\pi^+$; see below.

$\Gamma(\Lambda\pi^+ \eta)/\Gamma(\rho K^- \pi^+)$ Γ_{38}/Γ_2

Unseen decay modes of the η are included.

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------------------|-------------|--------------------|-------------|--------------------------------|
| 0.36 ± 0.07 OUR AVERAGE | | | | |
| 0.41 ± 0.17 ± 0.10 | 11 | CRONIN-HEN..03 | CLE3 | $e^+ e^- \approx \Upsilon(4S)$ |
| 0.35 ± 0.05 ± 0.06 | 116 | AMMAR | 95 CLE2 | $e^+ e^- \approx \Upsilon(4S)$ |

$\Gamma(\Sigma(1385)^+ \eta)/\Gamma(\rho K^- \pi^+)$ Γ_{39}/Γ_2

Unseen decay modes of the $\Sigma(1385)^+$ and η are included.

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------|-------------|--------------------|-------------|--------------------------------|
| 0.17 ± 0.04 ± 0.03 | 54 | AMMAR | 95 CLE2 | $e^+ e^- \approx \Upsilon(4S)$ |

$\Gamma(\Lambda\pi^+\omega)/\Gamma(pK^-\pi^+)$ Γ_{40}/Γ_2

Unseen decay modes of the ω are included.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------|------|----------------|------|-------------------------------|
| 0.24±0.06±0.06 | 32 | CRONIN-HEN..03 | CLE3 | $e^+e^- \approx \Upsilon(4S)$ |

$\Gamma(\Lambda\pi^-\pi^0 2\pi^+, \text{no } \eta \text{ or } \omega)/\Gamma(pK^-\pi^+)$ Γ_{41}/Γ_2

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------------|-----|----------------|------|-------------------------------|
| <0.13 | 90 | CRONIN-HEN..03 | CLE3 | $e^+e^- \approx \Upsilon(4S)$ |

$\Gamma(\Lambda K^+\bar{K}^0)/\Gamma(pK^-\pi^+)$ Γ_{42}/Γ_2

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------------|-------------------------------------|-------------|----------|--|
| 0.090±0.017 OUR FIT | Error includes scale factor of 1.9. | | | |
| 0.131±0.020 OUR AVERAGE | | | | |
| 0.142±0.018±0.022 | 251 | LINK | 05F FOCS | γ nucleus, $\bar{E}_\gamma \approx 180$ GeV |
| 0.12 ±0.02 ±0.02 | 59 | AMMAR | 95 CLE2 | $e^+e^- \approx \Upsilon(4S)$ |

$\Gamma(\Xi(1690)^0 K^+, \Xi^{*0} \rightarrow \Lambda\bar{K}^0)/\Gamma(\Lambda K^+\bar{K}^0)$ Γ_{43}/Γ_{42}

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------------|-------|-------------|----------|--|
| 0.28±0.07 OUR AVERAGE | | | | |
| 0.32±0.10±0.04 | 84±24 | LINK | 05F FOCS | γ nucleus, $\bar{E}_\gamma \approx 180$ GeV |
| 0.26±0.08±0.03 | 93 | ABE | 02C BELL | $e^+e^- \approx \Upsilon(4S)$ |

$\Gamma(\Lambda K^+\bar{K}^0)/\Gamma(\Lambda\pi^+)$ Γ_{42}/Γ_{28}

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------------|-------------------------------------|-------------|----------|-------------------------------|
| 0.44 ±0.08 OUR FIT | Error includes scale factor of 2.0. | | | |
| 0.395±0.026±0.036 | 460 ± 30 | AUBERT | 07U BABR | $e^+e^- \approx \Upsilon(4S)$ |

$\Gamma(\Sigma^0\pi^+)/\Gamma_{\text{total}}$ Γ_{44}/Γ

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------------------------------|-------------|---------|--|
| 1.29±0.07 OUR FIT | Error includes scale factor of 1.1. | | | |
| 1.27±0.08±0.03 | 522 | ABLIKIM | 16 BES3 | $e^+e^- \rightarrow \Lambda_c\bar{\Lambda}_c, 4.599$ GeV |

$\Gamma(\Sigma^0\pi^+)/\Gamma(pK^-\pi^+)$ Γ_{44}/Γ_2

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------------------------------|-------------------------------------|-------------|---------|---|
| 0.206±0.010 OUR FIT | Error includes scale factor of 1.2. | | | |
| 0.20 ±0.04 OUR AVERAGE | | | | |
| 0.21 ±0.02 ±0.04 | 196 | AVERY | 94 CLE2 | $e^+e^- \approx \Upsilon(3S), \Upsilon(4S)$ |
| 0.17 ±0.06 ±0.04 | | ALBRECHT | 92 ARG | $e^+e^- \approx 10.4$ GeV |

$\Gamma(\Sigma^0\pi^+)/\Gamma(\Lambda\pi^+)$ Γ_{44}/Γ_{28}

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------------------------------|------|-------------|----------|--|
| 0.99 ±0.04 OUR FIT | | | | |
| 0.98 ±0.05 OUR AVERAGE | | | | |
| 0.977±0.015±0.051 | 33k | AUBERT | 07U BABR | $e^+e^- \approx \Upsilon(4S)$ |
| 1.09 ±0.11 ±0.19 | 750 | LINK | 05F FOCS | γ nucleus, $\bar{E}_\gamma \approx 180$ GeV |

$\Gamma(\Sigma^+\pi^0)/\Gamma_{\text{total}}$ Γ_{45}/Γ

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|---------|--|
| 1.25±0.10 OUR FIT | | | | |
| 1.18±0.10±0.03 | 309 | ABLIKIM | 16 BES3 | $e^+e^- \rightarrow \Lambda_c\bar{\Lambda}_c, 4.599$ GeV |

$\Gamma(\Sigma^+ \pi^0)/\Gamma(pK^- \pi^+)$ Γ_{45}/Γ_2

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------------------|-------------|--------------------|-------------|------------------------------------|
| 0.199±0.015 OUR FIT | | | | |
| 0.20 ±0.03 ±0.03 | 93 | KUBOTA | 93 | CLE2 $e^+e^- \approx \Upsilon(4S)$ |

 $\Gamma(\Sigma^+ \eta)/\Gamma(pK^- \pi^+)$ Γ_{46}/Γ_2 Unseen decay modes of the η are included.

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------------|-------------|--------------------|-------------|------------------------------------|
| 0.11±0.03±0.02 | 26 | AMMAR | 95 | CLE2 $e^+e^- \approx \Upsilon(4S)$ |

 $\Gamma(\Sigma^+ \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{47}/Γ

| <u>VALUE (%)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------------|-------------|--------------------|-------------|---|
| 4.50±0.25 OUR FIT | | | | Error includes scale factor of 1.3. |
| 4.25±0.24±0.20 | 1156 | ABLIKIM | 16 | BES3 $e^+e^- \rightarrow \Lambda_c \bar{\Lambda}_c$, 4.599 GeV |

 $\Gamma(\Sigma^+ \pi^+ \pi^-)/\Gamma(pK^- \pi^+)$ Γ_{47}/Γ_2

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|------------------------------------|
| 0.716±0.019 OUR FIT | | | | |
| 0.720±0.024 OUR AVERAGE | | | | |
| 0.719±0.003±0.024 | 2.7M | BERGER | 18 | BELL $e^+e^- \approx \Upsilon(4S)$ |
| 0.74 ±0.07 ±0.09 | 487 | KUBOTA | 93 | CLE2 $e^+e^- \approx \Upsilon(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.72 ±0.14 | 47 ± 9 | VAZQUEZ-JA..08 | SELX | Σ^- nucleus, 600 GeV |
| 0.54 $\begin{smallmatrix} +0.18 \\ -0.15 \end{smallmatrix}$ | 11 | BARLAG | 92 | NA32 π^- Cu 230 GeV |

 $\Gamma(\Sigma^+ \rho^0)/\Gamma(pK^- \pi^+)$ Γ_{48}/Γ_2

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------|------------|--------------------|-------------|------------------------------------|
| <0.27 | 95 | KUBOTA | 93 | CLE2 $e^+e^- \approx \Upsilon(4S)$ |

 $\Gamma(\Sigma^- 2\pi^+)/\Gamma_{\text{total}}$ Γ_{49}/Γ

| <u>VALUE (%)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------------|-------------|--------------------|-------------|--------------------------|
| 1.87±0.18 OUR FIT | | | | |
| 1.81±0.17±0.09 | 161 | ABLIKIM | 17Y | BES3 e^+e^- at 4.6 GeV |

 $\Gamma(\Sigma^- 2\pi^+)/\Gamma(pK^- \pi^+)$ Γ_{49}/Γ_2

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------------------|-------------|--------------------|-------------|-------------------------------------|
| 0.297±0.030 OUR FIT | | | | Error includes scale factor of 1.1. |
| 0.314±0.067 | 30 ± 6 | VAZQUEZ-JA..08 | SELX | Σ^- nucleus, 600 GeV |

 $\Gamma(\Sigma^- 2\pi^+)/\Gamma(\Sigma^+ \pi^+ \pi^-)$ Γ_{49}/Γ_{47}

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------------|-------------|--------------------|-------------|--|
| 0.42±0.04 OUR FIT | | | | Error includes scale factor of 1.1. |
| 0.53±0.15±0.07 | 56 | FRABETTI | 94E | E687 γ Be, \bar{E}_γ 220 GeV |

 $\Gamma(\Sigma^0 \pi^+ \pi^0)/\Gamma(pK^- \pi^+)$ Γ_{50}/Γ_2

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-------------------------------|-------------|--------------------|-------------|--|
| 0.56 ±0.05 OUR AVERAGE | | | | Error includes scale factor of 1.5. |
| 0.575±0.005±0.036 | 2.7M | BERGER | 18 | BELL $e^+e^- \approx \Upsilon(4S)$ |
| 0.36 ±0.09 ±0.10 | 117 | AVERY | 94 | CLE2 $e^+e^- \approx \Upsilon(3S), \Upsilon(4S)$ |

$\Gamma(\Sigma^+ \pi^0 \pi^0)/\Gamma(pK^- \pi^+)$ Γ_{51}/Γ_2

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|-------------------------------------|
| 0.247±0.006±0.019 | 925k | BERGER | 18 | BELL $e^+ e^- \approx \Upsilon(4S)$ |

$\Gamma(\Sigma^0 \pi^- 2\pi^+)/\Gamma(pK^- \pi^+)$ Γ_{52}/Γ_2

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---|
| 0.18±0.05 OUR FIT | | | | |
| 0.21±0.05±0.05 | 90 | AVERY | 94 | CLE2 $e^+ e^- \approx \Upsilon(3S), \Upsilon(4S)$ |

$\Gamma(\Sigma^0 \pi^- 2\pi^+)/\Gamma(\Lambda \pi^- 2\pi^+)$ Γ_{52}/Γ_{31}

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---|
| 0.31±0.08 OUR FIT | | | | |
| 0.26±0.06±0.09 | 480 | LINK | 05F | FOCS γ nucleus, $\bar{E}_\gamma \approx 180$ GeV |

$\Gamma(\Sigma^+ \omega)/\Gamma_{\text{total}}$ Γ_{54}/Γ

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|--|
| 1.70±0.21 OUR FIT | | | | |
| 1.56±0.20±0.07 | 157 | ABLIKIM | 16 | BES3 $e^+ e^- \rightarrow \Lambda_c \bar{\Lambda}_c$, 4.599 GeV |

$\Gamma(\Sigma^+ \omega)/\Gamma(pK^- \pi^+)$ Γ_{54}/Γ_2

Unseen decay modes of the ω are included.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------------|------|-------------|------|-------------------------------------|
| 0.271±0.031 OUR FIT | | | | |
| 0.54 ±0.13 ±0.06 | 107 | KUBOTA | 93 | CLE2 $e^+ e^- \approx \Upsilon(4S)$ |

$\Gamma(\Sigma^- \pi^0 2\pi^+)/\Gamma_{\text{total}}$ Γ_{55}/Γ

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------|------|-------------|------|---------------------------|
| 2.11±0.33±0.14 | 88 | ABLIKIM | 17Y | BES3 $e^+ e^-$ at 4.6 GeV |

$\Gamma(\Sigma^+ K^+ K^-)/\Gamma(pK^- \pi^+)$ Γ_{56}/Γ_2

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------------|------|-------------|------|---------------------------------|
| 0.056±0.006 OUR FIT | | | | |
| 0.070±0.011±0.011 | 59 | AVERY | 93 | CLE2 $e^+ e^- \approx 10.5$ GeV |

$\Gamma(\Sigma^+ K^+ K^-)/\Gamma(\Sigma^+ \pi^+ \pi^-)$ Γ_{56}/Γ_{47}

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------------|------|-------------|------|--|
| 0.078±0.008 OUR FIT | | | | |
| 0.074±0.009 OUR AVERAGE | | | | |
| 0.076±0.007±0.009 | 246 | ABE | 02C | BELL $e^+ e^- \approx \Upsilon(4S)$ |
| 0.071±0.011±0.011 | 103 | LINK | 02G | FOCS γ nucleus, ≈ 180 GeV |

$\Gamma(\Sigma^+ \phi)/\Gamma(pK^- \pi^+)$ Γ_{57}/Γ_2

Unseen decay modes of the ϕ are included.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------------|------|-------------|------|---------------------------------|
| 0.062±0.009 OUR FIT | | | | |
| 0.069±0.023±0.016 | 26 | AVERY | 93 | CLE2 $e^+ e^- \approx 10.5$ GeV |

$\Gamma(\Sigma^+\phi)/\Gamma(\Sigma^+\pi^+\pi^-)$ Γ_{57}/Γ_{47}

Unseen decay modes of the ϕ are included.

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------------------|-------------|--------------------|-------------|--|
| 0.087±0.012 OUR FIT | | | | |
| 0.086±0.012 OUR AVERAGE | | | | |
| 0.085±0.012±0.012 | 129 | ABE | 02C | BELL $e^+e^- \approx \Upsilon(4S)$ |
| 0.087±0.016±0.006 | 57 | LINK | 02G | FOCS γ nucleus, ≈ 180 GeV |

$\Gamma(\Xi(1690)^0 K^+, \Xi^{*0} \rightarrow \Sigma^+ K^-)/\Gamma(\Sigma^+\pi^+\pi^-)$ Γ_{58}/Γ_{47}

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------------------|-------------|--------------------|-------------|--|
| 0.023±0.005 OUR AVERAGE | | | | |
| 0.023±0.005±0.005 | 75 | ABE | 02C | BELL $e^+e^- \approx \Upsilon(4S)$ |
| 0.022±0.006±0.006 | 34 | LINK | 02G | FOCS γ nucleus, ≈ 180 GeV |

$\Gamma(\Sigma^+ K^+ K^- \text{ nonresonant})/\Gamma(\Sigma^+\pi^+\pi^-)$ Γ_{59}/Γ_{47}

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|--|
| <0.018 | | | | |
| | 90 | ABE | 02C | BELL $e^+e^- \approx \Upsilon(4S)$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| <0.028 | 90 | LINK | 02G | FOCS γ nucleus, ≈ 180 GeV |

$\Gamma(\Xi^0 K^+)/\Gamma_{\text{total}}$ Γ_{60}/Γ

| <u>VALUE (units 10^{-3})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|--------------------|-------------|--------------------------|
| 5.5 ±0.7 OUR FIT | | | | |
| 5.90±0.86±0.39 | 68 | ABLIKIM | 18Y | BES3 e^+e^- at 4.6 GeV |


$\Gamma(\Xi^0 K^+)/\Gamma(pK^-\pi^+)$ Γ_{60}/Γ_2

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------------------|-------------|--------------------|-------------|--------------------------------|
| 0.088±0.012 OUR FIT | | | | |
| 0.078±0.013±0.013 | 56 | AVERY | 93 | CLE2 $e^+e^- \approx 10.5$ GeV |

$\Gamma(\Xi^- K^+ \pi^+)/\Gamma(pK^-\pi^+)$ Γ_{61}/Γ_2

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|-------------|--------------------|-------------|--------------------------------|
| 0.099±0.009 OUR FIT | | | | |
| 0.098±0.021 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below. | | | | |
| 0.14 ±0.03 ±0.02 | 34 | ALBRECHT | 95B | ARG $e^+e^- \approx 10.4$ GeV |
| 0.079±0.013±0.014 | 60 | AVERY | 93 | CLE2 $e^+e^- \approx 10.5$ GeV |
| 0.15 ±0.04 ±0.03 | 30 | AVERY | 91 | CLEO $e^+e^- 10.5$ GeV |

WEIGHTED AVERAGE
 0.098 ± 0.021 (Error scaled by 1.3)



Values above of weighted average, error, and scale factor are based upon the data in this ideogram only. They are not necessarily the same as our 'best' values, obtained from a least-squares constrained fit utilizing measurements of other (related) quantities as additional information.

| | | | χ^2 |
|----------------------------|-----|------|----------|
| ALBRECHT | 95B | ARG | 1.3 |
| AVERY | 93 | CLE2 | 1.0 |
| AVERY | 91 | CLEO | 1.1 |
| | | | 3.4 |
| (Confidence Level = 0.180) | | | |

$$\Gamma(\Xi^- K^+ \pi^+) / \Gamma(\rho K^- \pi^+)$$

$$\Gamma(\Xi^- K^+ \pi^+) / \Gamma(\Lambda \pi^+)$$

$\Gamma_{61} / \Gamma_{28}$

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------------|-----------|-------------|----------|------------------------------|
| 0.48 ± 0.04 OUR FIT | | | | |
| 0.480 ± 0.016 ± 0.039 | 2665 ± 84 | AUBERT | 07U BABR | $e^+ e^- \approx \gamma(4S)$ |

$$\Gamma(\Xi(1530)^0 K^+) / \Gamma_{\text{total}}$$

Γ_{62} / Γ

| VALUE (units 10^{-3}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------------|------|-------------|----------|-------------------------------------|
| 4.3 ± 0.9 OUR FIT | | | | Error includes scale factor of 1.1. |
| 5.02 ± 0.99 ± 0.31 | 60 | ABLIKIM | 18Y BES3 | $e^+ e^-$ at 4.6 GeV |

$$\Gamma(\Xi(1530)^0 K^+) / \Gamma(\rho K^- \pi^+)$$

Γ_{62} / Γ_2

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|---------|-------------------------------------|
| 0.068 ± 0.014 OUR FIT | | | | Error includes scale factor of 1.1. |
| 0.053 ± 0.016 ± 0.010 | 24 | AVERY | 93 CLE2 | $e^+ e^- \approx 10.5$ GeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.05 ± 0.02 ± 0.01 | 11 | ALBRECHT | 95B ARG | $e^+ e^- \approx 10.4$ GeV |

———— Hadronic modes with a hyperon: $S = 0$ final states ————

$$\Gamma(\Lambda K^+) / \Gamma(\Lambda \pi^+)$$

$\Gamma_{63} / \Gamma_{28}$

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------------------|------------|-------------|----------|-------------------------------------|
| 0.047 ± 0.009 OUR AVERAGE | | | | Error includes scale factor of 1.8. |
| 0.044 ± 0.004 ± 0.003 | 1162 ± 101 | AUBERT | 07U BABR | $e^+ e^- \approx \gamma(4S)$ |
| 0.074 ± 0.010 ± 0.012 | 265 | ABE | 02C BELL | $e^+ e^- \approx \gamma(4S)$ |

$\Gamma(\Lambda K^+ \pi^+ \pi^-)/\Gamma(\Lambda \pi^+)$ Γ_{64}/Γ_{28}

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-----|-------------|----------|--------------------------------|
| $<4.1 \times 10^{-2}$ | 90 | AUBERT | 07U BABR | $e^+ e^- \approx \Upsilon(4S)$ |

$\Gamma(\Sigma^0 K^+)/\Gamma(\Sigma^0 \pi^+)$ Γ_{65}/Γ_{44}

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------------------|----------|-------------|----------|--------------------------------|
| 0.040 ± 0.006 OUR AVERAGE | | | | |
| 0.038 ± 0.005 ± 0.003 | 366 ± 52 | AUBERT | 07U BABR | $e^+ e^- \approx \Upsilon(4S)$ |
| 0.056 ± 0.014 ± 0.008 | 75 | ABE | 02C BELL | $e^+ e^- \approx \Upsilon(4S)$ |

$\Gamma(\Sigma^0 K^+ \pi^+ \pi^-)/\Gamma(\Sigma^0 \pi^+)$ Γ_{66}/Γ_{44}

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-----|-------------|----------|--------------------------------|
| $<2.0 \times 10^{-2}$ | 90 | AUBERT | 07U BABR | $e^+ e^- \approx \Upsilon(4S)$ |

$\Gamma(\Sigma^+ K^+ \pi^-)/\Gamma(\Sigma^+ \pi^+ \pi^-)$ Γ_{67}/Γ_{47}

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------------|------|-------------|----------|--------------------------------|
| 0.047 ± 0.011 ± 0.008 | 105 | ABE | 02C BELL | $e^+ e^- \approx \Upsilon(4S)$ |

$\Gamma(\Sigma^+ K^*(892)^0)/\Gamma(\Sigma^+ \pi^+ \pi^-)$ Γ_{68}/Γ_{47}

Unseen decay modes of the $K^*(892)^0$ are included.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------------|------|-------------|----------|-------------------------------------|
| 0.078 ± 0.018 ± 0.013 | 49 | LINK | 02G FOCS | γ nucleus, ≈ 180 GeV |

$\Gamma(\Sigma^- K^+ \pi^+)/\Gamma(\Sigma^+ K^*(892)^0)$ Γ_{69}/Γ_{68}

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---------|-----|-------------|----------|-------------------------------------|
| <0.35 | 90 | LINK | 02G FOCS | γ nucleus, ≈ 180 GeV |

———— Doubly Cabibbo-suppressed modes ————

$\Gamma(\rho K^+ \pi^-)/\Gamma(\rho K^- \pi^+)$ Γ_{70}/Γ_2

| VALUE (units 10^{-3}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|-------------|----------|---|
| 1.77 ± 0.27 OUR AVERAGE Error includes scale factor of 1.9. | | | | |
| 1.65 ± 0.15 ± 0.05 | 392 | AAIJ | 18V LHCB | $\Lambda_b^0 \rightarrow \Lambda_c^+ \mu^- X$ |
| 2.35 ± 0.27 ± 0.21 | 3379 | YANG | 16 BELL | At or near Υ s |

———— Semileptonic modes ————

$\Gamma(\Lambda e^+ \nu_e)/\Gamma_{\text{total}}$ Γ_{71}/Γ

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------------|------|-------------|----------|-----------------------------------|
| 3.63 ± 0.38 ± 0.20 | 104 | ABLIKIM | 15Y BES3 | 567 pb^{-1} , 4.599 GeV |

$\Gamma(\Lambda e^+ \nu_e)/\Gamma(e^+ \text{ anything})$ Γ_{71}/Γ_{73}

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|-----------|-------------------|
| 91.9 ± 12.5 ± 5.4 | 214 | ABLIKIM | 18AF BES3 | $e^+ e^-$ 4.6 GeV |

$\Gamma(\Lambda e^+ \nu_e)/\Gamma(\rho K^- \pi^+)$ Γ_{71}/Γ_2

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------|--------------------|-------------|----------------|
|--------------|--------------------|-------------|----------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|-------------|------------------|------|------------------------------------|
| 0.43 ± 0.08 | 1,2 BERGFELD 94 | CLE2 | $e^+ e^- \approx \Upsilon(4S)$ |
| 0.38 ± 0.14 | 2,3 ALBRECHT 91G | ARG | $e^+ e^- \approx 10.4 \text{ GeV}$ |

¹ BERGFELD 94 measures $\sigma(e^+ e^- \rightarrow \Lambda_c^+ X) \cdot B(\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e) = (4.87 \pm 0.28 \pm 0.69) \text{ pb}$.

² To extract $\Gamma(\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e)/\Gamma(\Lambda_c^+ \rightarrow \rho K^- \pi^+)$, we use $\sigma(e^+ e^- \rightarrow \Lambda_c^+ X) \cdot B(\Lambda_c^+ \rightarrow \rho K^- \pi^+) = (11.2 \pm 1.3) \text{ pb}$, which is the weighted average of measurements from ARGUS (ALBRECHT 96E) and CLEO (AVERY 91).

³ ALBRECHT 91G measures $\sigma(e^+ e^- \rightarrow \Lambda_c^+ X) \cdot B(\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e) = (4.20 \pm 1.28 \pm 0.71) \text{ pb}$.

$\Gamma(\Lambda \mu^+ \nu_\mu)/\Gamma_{\text{total}}$ Γ_{72}/Γ

| <u>VALUE (%)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|------------------|-------------|--------------------|-------------|----------------|
|------------------|-------------|--------------------|-------------|----------------|

| | | | | |
|---------------------------|----|-------------|------|----------------------|
| 3.49 ± 0.46 ± 0.27 | 79 | ABLIKIM 17D | BES3 | $e^+ e^-$ at 4.6 GeV |
|---------------------------|----|-------------|------|----------------------|

$\Gamma(\Lambda \mu^+ \nu_\mu)/\Gamma(\rho K^- \pi^+)$ Γ_{72}/Γ_2

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------|--------------------|-------------|----------------|
|--------------|--------------------|-------------|----------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|-------------|------------------|------|------------------------------------|
| 0.40 ± 0.09 | 1,2 BERGFELD 94 | CLE2 | $e^+ e^- \approx \Upsilon(4S)$ |
| 0.35 ± 0.20 | 2,3 ALBRECHT 91G | ARG | $e^+ e^- \approx 10.4 \text{ GeV}$ |

¹ BERGFELD 94 measures $\sigma(e^+ e^- \rightarrow \Lambda_c^+ X) \cdot B(\Lambda_c^+ \rightarrow \Lambda \mu^+ \nu_\mu) = (4.43 \pm 0.51 \pm 0.64) \text{ pb}$.

² To extract $\Gamma(\Lambda_c^+ \rightarrow \Lambda \mu^+ \nu_\mu)/\Gamma(\Lambda_c^+ \rightarrow \rho K^- \pi^+)$, we use $\sigma(e^+ e^- \rightarrow \Lambda_c^+ X) \cdot B(\Lambda_c^+ \rightarrow \rho K^- \pi^+) = (11.2 \pm 1.3) \text{ pb}$, which is the weighted average of measurements from ARGUS (ALBRECHT 96E) and CLEO (AVERY 91).

³ ALBRECHT 91G measures $\sigma(e^+ e^- \rightarrow \Lambda_c^+ X) \cdot B(\Lambda_c^+ \rightarrow \Lambda \mu^+ \nu_\mu) = (3.91 \pm 2.02 \pm 0.90) \text{ pb}$.

$\Gamma(\Lambda \mu^+ \nu_\mu)/\Gamma(\Lambda e^+ \nu_e)$ Γ_{72}/Γ_{71}

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------|--------------------|-------------|----------------|
|--------------|--------------------|-------------|----------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|--------------------|--------------------------|------|----------------------|
| 0.96 ± 0.16 ± 0.04 | ¹ ABLIKIM 17D | BES3 | $e^+ e^-$ at 4.6 GeV |
|--------------------|--------------------------|------|----------------------|

¹ This is the ratio of the ABLIKIM 17D $\Lambda \mu^+ \nu_e$ branching fraction and the ABLIKIM 15Y $\Lambda e^+ \nu_e$ branching fraction (see above), and so is not an independent measurement.

———— Inclusive modes ————

$\Gamma(e^+ \text{ anything})/\Gamma_{\text{total}}$ Γ_{73}/Γ

| <u>VALUE (%)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|------------------|-------------|--------------------|-------------|----------------|
|------------------|-------------|--------------------|-------------|----------------|

| | | | | |
|---------------------------|-----|--------------|------|-------------------|
| 3.95 ± 0.34 ± 0.09 | 214 | ABLIKIM 18AF | BES3 | $e^+ e^-$ 4.6 GeV |
|---------------------------|-----|--------------|------|-------------------|

$\Gamma(\rho \text{ anything})/\Gamma_{\text{total}}$ Γ_{74}/Γ

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------|--------------------|-------------|----------------|
|--------------|--------------------|-------------|----------------|

| | | | |
|---------------------------|--------------------------|------|--------------------|
| 0.50 ± 0.08 ± 0.14 | ¹ CRAWFORD 92 | CLEO | $e^+ e^-$ 10.5 GeV |
|---------------------------|--------------------------|------|--------------------|

¹ This CRAWFORD 92 value includes protons from Λ decay. The value is model dependent, but account is taken of this in the systematic error.

$\Gamma(n \text{ anything})/\Gamma_{\text{total}}$ **Γ_{75}/Γ**

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|--------------------------|-------------|-------------------|
| $0.50 \pm 0.08 \pm 0.14$ | ¹ CRAWFORD 92 | CLEO | e^+e^- 10.5 GeV |

¹This CRAWFORD 92 value includes neutrons from Λ decay. The value is model dependent, but account is taken of this in the systematic error.

$\Gamma(\Lambda \text{ anything})/\Gamma_{\text{total}}$ **Γ_{76}/Γ**

| <u>VALUE (%)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|-------------|--------------------|-------------|---------------------|
| $38.2^{+2.8}_{-2.2} \pm 0.9$ | 700 | ABLIKIM | 18E BES3 | e^+e^- at 4.6 GeV |

$\Gamma(3\text{prongs})/\Gamma_{\text{total}}$ **Γ_{77}/Γ**

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|--------------------|-------------|--------------------------------------|
| $0.24 \pm 0.07 \pm 0.04$ | KAYIS-TOPAK.03 | CHRS | ν_μ emulsion, $\bar{E}=27$ GeV |

————— Rare or forbidden modes —————

$\Gamma(p e^+ e^-)/\Gamma_{\text{total}}$ **Γ_{78}/Γ**

A test for the $\Delta C=1$ weak neutral current. Allowed by higher-order electroweak interactions.

| <u>VALUE</u> | <u>CL%</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|---------------|--------------------|-------------|-------------------------------|
| $< 5.5 \times 10^{-6}$ | 90 | 4.0 ± 7.1 | LEES | 11G BABR | $e^+e^- \approx \Upsilon(4S)$ |

$\Gamma(p \mu^+ \mu^- \text{ non-resonant})/\Gamma_{\text{total}}$ **Γ_{79}/Γ**

A test for the $\Delta C=1$ weak neutral current. Allowed by higher-order electroweak interactions.

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|--|
| $< 7.7 \times 10^{-8}$ | 90 | AAIJ | 18N LHCb | Ratio to $p\phi$, $\phi \rightarrow \mu^+\mu^-$ |
| $< 4.4 \times 10^{-5}$ | 90 | LEES | 11G BABR | $e^+e^- \approx \Upsilon(4S)$ |
| $< 3.4 \times 10^{-4}$ | 90 | KODAMA | 95 E653 | π^- emulsion 600 GeV |

$\Gamma(p e^+ \mu^-)/\Gamma_{\text{total}}$ **Γ_{80}/Γ**

A test of lepton family-number conservation.

| <u>VALUE</u> | <u>CL%</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|----------------|--------------------|-------------|-------------------------------|
| $< 9.9 \times 10^{-6}$ | 90 | -0.7 ± 3.0 | LEES | 11G BABR | $e^+e^- \approx \Upsilon(4S)$ |

$\Gamma(p e^- \mu^+)/\Gamma_{\text{total}}$ **Γ_{81}/Γ**

A test of lepton family-number conservation.

| <u>VALUE</u> | <u>CL%</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|------------|---------------|--------------------|-------------|-------------------------------|
| $< 19 \times 10^{-6}$ | 90 | 6.2 ± 4.9 | LEES | 11G BABR | $e^+e^- \approx \Upsilon(4S)$ |

$\Gamma(\bar{p} 2e^+)/\Gamma_{\text{total}}$ **Γ_{82}/Γ**

A test of lepton- and baryon-number conservation.

| <u>VALUE</u> | <u>CL%</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|----------------|--------------------|-------------|-------------------------------|
| $< 2.7 \times 10^{-6}$ | 90 | -1.5 ± 4.5 | LEES | 11G BABR | $e^+e^- \approx \Upsilon(4S)$ |

$\Gamma(\bar{p} 2\mu^+)/\Gamma_{\text{total}}$ **Γ_{83}/Γ**

A test of lepton- and baryon-number conservation and of lepton family-number conservation.

| <u>VALUE</u> | <u>CL%</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|---------------|--------------------|-------------|-------------------------------|
| $< 9.4 \times 10^{-6}$ | 90 | 0.0 ± 2.2 | LEES | 11G BABR | $e^+e^- \approx \Upsilon(4S)$ |

$\Gamma(\bar{p}e^+\mu^+)/\Gamma_{\text{total}}$ Γ_{84}/Γ
 A test of lepton- and baryon-number conservation and of lepton family-number conservation.

| VALUE | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------|-----|----------------|-------------|----------|-------------------------------|
| $<16 \times 10^{-6}$ | 90 | 10.1 ± 6.8 | LEES | 11G BABR | $e^+e^- \approx \Upsilon(4S)$ |

$\Gamma(\Sigma^-\mu^+\mu^+)/\Gamma_{\text{total}}$ Γ_{85}/Γ
 A test of lepton-number conservation.

| VALUE | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-----|------|-------------|---------|--------------------------|
| $<7.0 \times 10^{-4}$ | 90 | 0 | KODAMA | 95 E653 | π^- emulsion 600 GeV |

Λ_c^+ DECAY PARAMETERS

See the note on “Baryon Decay Parameters” in the neutron Listings.

α FOR $\Lambda_c^+ \rightarrow \Lambda\pi^+$

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|---------------------|----------|--|
| -0.91 ± 0.15 OUR AVERAGE | | | | |
| $-0.78 \pm 0.16 \pm 0.19$ | | LINK | 06A FOCS | γ A, $\bar{E}_\gamma \approx 180$ GeV |
| $-0.94 \pm 0.21 \pm 0.12$ | 414 | ¹ BISHAI | 95 CLE2 | $e^+e^- \approx \Upsilon(4S)$ |
| -0.96 ± 0.42 | | ALBRECHT | 92 ARG | $e^+e^- \approx 10.4$ GeV |
| -1.1 ± 0.4 | 86 | AVERY | 90B CLEO | $e^+e^- \approx 10.6$ GeV |

¹ BISHAI 95 actually gives $\alpha = -0.94^{+0.21+0.12}_{-0.06-0.06}$, chopping the errors at the physical limit -1.0 . However, for $\alpha \approx -1.0$, some experiments should *get* unphysical values ($\alpha < -1.0$), and for averaging with other measurements such values (or errors that extend below -1.0) should *not* be chopped.

α FOR $\Lambda_c^+ \rightarrow \Sigma^+\pi^0$

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|---------|-------------------------------|
| $-0.45 \pm 0.31 \pm 0.06$ | 89 | BISHAI | 95 CLE2 | $e^+e^- \approx \Upsilon(4S)$ |

α FOR $\Lambda_c^+ \rightarrow \Lambda\ell^+\nu_\ell$

The experiments don't cover the complete (or same incomplete) $M(\Lambda\ell^+)$ range, but we average them together anyway.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-----------------------|---------|-------------------------------|
| -0.86 ± 0.04 OUR AVERAGE | | | | |
| $-0.86 \pm 0.03 \pm 0.02$ | 3201 | ¹ HINSON | 05 CLEO | $e^+e^- \approx \Upsilon(4S)$ |
| $-0.91 \pm 0.42 \pm 0.25$ | | ² ALBRECHT | 94B ARG | $e^+e^- \approx 10$ GeV |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| $-0.82^{+0.09+0.06}_{-0.06-0.03}$ | 700 | ³ CRAWFORD | 95 CLE2 | See HINSON 05 |
| $-0.89^{+0.17+0.09}_{-0.11-0.05}$ | 350 | ⁴ BERGFELD | 94 CLE2 | See CRAWFORD 95 |

¹ HINSON 05 measures the form-factor ratio $R \equiv f_2/f_1$ for $\Lambda_c^+ \rightarrow \Lambda e^+\nu_e$ events to be $-0.31 \pm 0.05 \pm 0.04$ and the pole mass to be $2.21 \pm 0.08 \pm 0.14$ GeV/ c^2 , and from these calculates α , averaged over q^2 , where $\langle q^2 \rangle = 0.67$ (GeV/ c)².

² ALBRECHT 94B uses Λe^+ and $\Lambda \mu^+$ events in the mass range $1.85 < M(\Lambda\ell^+) < 2.20$ GeV.

³ CRAWFORD 95 measures the form-factor ratio $R \equiv f_2/f_1$ for $\Lambda_c^+ \rightarrow \Lambda e^+\nu_e$ events to be $-0.25 \pm 0.14 \pm 0.08$ and from this calculates α , averaged over q^2 , to be the above.

⁴ BERGFELD 94 uses Λe^+ events.

Λ_c^+ , $\bar{\Lambda}_c^-$ CP-VIOLATING DECAY ASYMMETRIES

$(\alpha + \bar{\alpha})/(\alpha - \bar{\alpha})$ in $\Lambda_c^+ \rightarrow \Lambda\pi^+$, $\bar{\Lambda}_c^- \rightarrow \bar{\Lambda}\pi^-$

This is zero if CP is conserved.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---------------------------|-------------|------|---|
| $-0.07 \pm 0.19 \pm 0.24$ | LINK | 06A | FOCS γ A, $\bar{E}_\gamma \approx 180$ GeV |

$(\alpha + \bar{\alpha})/(\alpha - \bar{\alpha})$ in $\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e$, $\bar{\Lambda}_c^- \rightarrow \bar{\Lambda} e^- \bar{\nu}_e$

This is zero if CP is conserved.

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|------|-------------------------------------|
| $0.00 \pm 0.03 \pm 0.02$ | HINSON | 05 | CLEO $e^+ e^- \approx \Upsilon(4S)$ |

$A_{CP}(\Lambda X)$ in $\Lambda_c \rightarrow \Lambda X$, $\bar{\Lambda}_c \rightarrow \bar{\Lambda} X$

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------------|------|-------------|------|---------------------------|
| $2.1^{+7.0}_{-6.6} \pm 1.6$ | 700 | ABLIKIM | 18E | BES3 $e^+ e^-$ at 4.6 GeV |

$\Delta A_{CP} = A_{CP}(\Lambda_c^+ \rightarrow pK^+ K^-) - A_{CP}(\Lambda_c^+ \rightarrow p\pi^+ \pi^-)$

| VALUE (%) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------------|------|--------------------|
| $0.30 \pm 0.91 \pm 0.61$ | ¹ AAIJ | 18R | LHCB pp 7, 8 TeV |

¹ AAIJ 18R applies phase-space-dependent weights to the $\Lambda_c^+ \rightarrow p\pi^+ \pi^-$ sample to align its kinematics with the $\Lambda_c^+ \rightarrow pK^+ K^-$ sample.

Λ_c^+ REFERENCES

We have omitted some papers that have been superseded by later experiments. The omitted papers may be found in our 1992 edition (Physical Review **D45**, 1 June, Part II) or in earlier editions.

| | | | | |
|----------------|------|----------------|-----------------------------------|-----------------------|
| AAIJ | 18N | PR D97 091101 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 18R | JHEP 1803 182 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 18V | JHEP 1803 043 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| ABLIKIM | 18AF | PRL 121 251801 | M. Ablikim <i>et al.</i> | (BES III Collab.) |
| ABLIKIM | 18E | PRL 121 062003 | M. Ablikim <i>et al.</i> | (BES III Collab.) |
| ABLIKIM | 18Y | PL B783 200 | M. Ablikim <i>et al.</i> | (BES III Collab.) |
| BERGER | 18 | PR D98 112006 | M. Berger <i>et al.</i> | (BELLE Collab.) |
| ABLIKIM | 17D | PL B767 42 | M. Ablikim <i>et al.</i> | (BES III Collab.) |
| ABLIKIM | 17H | PRL 118 112001 | M. Ablikim <i>et al.</i> | (BES III Collab.) |
| ABLIKIM | 17Q | PR D95 111102 | M. Ablikim <i>et al.</i> | (BES III Collab.) |
| ABLIKIM | 17Y | PL B772 388 | M. Ablikim <i>et al.</i> | (BES III Collab.) |
| PAL | 17 | PR D96 051102 | B. Pal <i>et al.</i> | (BELLE Collab.) |
| ABLIKIM | 16 | PRL 116 052001 | M. Ablikim <i>et al.</i> | (BES III Collab.) |
| ABLIKIM | 16U | PRL 117 232002 | M. Ablikim <i>et al.</i> | (BES III Collab.) |
| YANG | 16 | PRL 117 011801 | S.B. Yang <i>et al.</i> | (BELLE Collab.) |
| ABLIKIM | 15Y | PRL 115 221805 | M. Ablikim <i>et al.</i> | (BES III Collab.) |
| ZUPANC | 14 | PRL 113 042002 | A. Zupanc <i>et al.</i> | (BELLE Collab.) |
| LEES | 11G | PR D84 072006 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| VAZQUEZ-JA... | 08 | PL B666 299 | E. Vazquez-Jauregui <i>et al.</i> | (SELEX Collab.) |
| AUBERT | 07U | PR D75 052002 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| LINK | 06A | PL B634 165 | J.M. Link <i>et al.</i> | (FNAL FOCUS Collab.) |
| AUBERT,B | 05S | PR D72 052006 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| HINSON | 05 | PRL 94 191801 | J.V. Hinson <i>et al.</i> | (CLEO Collab.) |
| LINK | 05F | PL B624 22 | J.M. Link <i>et al.</i> | (FNAL FOCUS Collab.) |
| CRONIN-HEN... | 03 | PR D67 012001 | D. Cronin-Hennessy <i>et al.</i> | (CLEO Collab.) |
| KAYIS-TOPAK... | 03 | PL B555 156 | A. Kayis-Topaksu <i>et al.</i> | (CERN CHORUS Collab.) |
| ABE | 02C | PL B524 33 | K. Abe <i>et al.</i> | (KEK BELLE Collab.) |
| LINK | 02C | PRL 88 161801 | J.M. Link <i>et al.</i> | (FNAL FOCUS Collab.) |
| LINK | 02G | PL B540 25 | J.M. Link <i>et al.</i> | (FNAL FOCUS Collab.) |

| | | | | |
|-------------|-----|------------------------------|----------------------------------|---------------------------|
| PDG | 02 | PR D66 010001 | K. Hagiwara <i>et al.</i> | (PDG Collab.) |
| KUSHNIR... | 01 | PRL 86 5243 | A. KUSHNIRENKO <i>et al.</i> | (FNAL SELEX Collab.) |
| MAHMOOD | 01 | PRL 86 2232 | A.H. Mahmood <i>et al.</i> | (CLEO Collab.) |
| AITALA | 00 | PL B471 449 | E.M. Aitala <i>et al.</i> | (FNAL E791 Collab.) |
| ALAM | 98 | PR D57 4467 | M.S. Alam <i>et al.</i> | (CLEO Collab.) |
| ALBRECHT | 96E | PRPL 276 223 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| ALEXANDER | 96C | PR D53 1013 | J.P. Alexander <i>et al.</i> | (CLEO Collab.) |
| ALBRECHT | 95B | PL B342 397 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| AMMAR | 95 | PRL 74 3534 | R. Ammar <i>et al.</i> | (CLEO Collab.) |
| BISHAI | 95 | PL B350 256 | M. Bishai <i>et al.</i> | (CLEO Collab.) |
| CRAWFORD | 95 | PRL 75 624 | G. Crawford <i>et al.</i> | (CLEO Collab.) |
| KODAMA | 95 | PL B345 85 | K. Kodama <i>et al.</i> | (FNAL E653 Collab.) |
| ALBRECHT | 94B | PL B326 320 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| ALEEV | 94 | PAN 57 1370 | A.N. Aleev <i>et al.</i> | (Serpukhov BIS-2 Collab.) |
| | | Translated from YF 57 1443. | | |
| AVERY | 94 | PL B325 257 | P. Avery <i>et al.</i> | (CLEO Collab.) |
| BERGFELD | 94 | PL B323 219 | T. Bergfeld <i>et al.</i> | (CLEO Collab.) |
| FRABETTI | 94E | PL B328 193 | P.L. Frabetti <i>et al.</i> | (FNAL E687 Collab.) |
| AVERY | 93 | PRL 71 2391 | P. Avery <i>et al.</i> | (CLEO Collab.) |
| BOZEK | 93 | PL B312 247 | A. Bozek <i>et al.</i> | (CERN NA32 Collab.) |
| FRABETTI | 93D | PRL 70 1755 | P.L. Frabetti <i>et al.</i> | (FNAL E687 Collab.) |
| FRABETTI | 93H | PL B314 477 | P.L. Frabetti <i>et al.</i> | (FNAL E687 Collab.) |
| KUBOTA | 93 | PRL 71 3255 | Y. Kubota <i>et al.</i> | (CLEO Collab.) |
| ALBRECHT | 92 | PL B274 239 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| BARLAG | 92 | PL B283 465 | S. Barlag <i>et al.</i> | (ACCMOR Collab.) |
| CRAWFORD | 92 | PR D45 752 | G. Crawford <i>et al.</i> | (CLEO Collab.) |
| JEZABEK | 92 | PL B286 175 | M. Jezabek, K. Rybicki, R. Rylko | (CRAC) |
| ALBRECHT | 91G | PL B269 234 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| AVERY | 91 | PR D43 3599 | P. Avery <i>et al.</i> | (CLEO Collab.) |
| ALVAREZ | 90 | ZPHY C47 539 | M.P. Alvarez <i>et al.</i> | (CERN NA14/2 Collab.) |
| ALVAREZ | 90B | PL B246 256 | M.P. Alvarez <i>et al.</i> | (CERN NA14/2 Collab.) |
| ANJOS | 90 | PR D41 801 | J.C. Anjos <i>et al.</i> | (FNAL E691 Collab.) |
| AVERY | 90B | PRL 65 2842 | P. Avery <i>et al.</i> | (CLEO Collab.) |
| BARLAG | 90D | ZPHY C48 29 | S. Barlag <i>et al.</i> | (ACCMOR Collab.) |
| FRABETTI | 90 | PL B251 639 | P.L. Frabetti <i>et al.</i> | (FNAL E687 Collab.) |
| BARLAG | 89 | PL B218 374 | S. Barlag <i>et al.</i> | (ACCMOR Collab.) |
| AGUILAR-... | 88B | ZPHY C40 321 | M. Aguilar-Benitez <i>et al.</i> | (LEBC-EHS Collab.) |
| Also | | PL B189 254 | M. Aguilar-Benitez <i>et al.</i> | (LEBC-EHS Collab.) |
| Also | | PL B199 462 | M. Aguilar-Benitez <i>et al.</i> | (LEBC-EHS Collab.) |
| Also | | SJNP 48 833 | M. Begalli <i>et al.</i> | (LEBC-EHS Collab.) |
| | | Translated from YAF 48 1310. | | |
| ALBRECHT | 88C | PL B207 109 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| ANJOS | 88B | PRL 60 1379 | J.C. Anjos <i>et al.</i> | (FNAL E691 Collab.) |
| AMENDOLIA | 87 | ZPHY C36 513 | S.R. Amendolia <i>et al.</i> | (CERN NA1 Collab.) |
| JONES | 87 | ZPHY C36 593 | G.T. Jones <i>et al.</i> | (CERN WA21 Collab.) |
| BOSETTI | 82 | PL 109B 234 | P.C. Bosetti <i>et al.</i> | (AACH3, BONN, CERN+) |
| BASILE | 81B | NC 62A 14 | M. Basile <i>et al.</i> | (CERN, BGNA, PGIA, FRAS) |
| CALICCHIO | 80 | PL 93B 521 | M. Calicchio <i>et al.</i> | (BARI, BIRM, BRUX+) |

OTHER RELATED PAPERS

| | | | | |
|-----------|----|---------------|---------------------------|--|
| MIGLIOZZI | 99 | PL B462 217 | P. Migliozi <i>et al.</i> | |
| DUNIETZ | 98 | PR D58 094010 | I. Dunietz | |