

$\phi(1680)$ 

$$I^G(J^{PC}) = 0^-(1^{--})$$

## $\phi(1680)$ MASS

### $e^+e^-$ PRODUCTION

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1680±20 OUR ESTIMATE</b>				
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1700±23	2k	<sup>1</sup> ACHASOV	18A SND	1.3–2.0 $e^+e^- \rightarrow K_S^0 K_L^0 \pi^0$
1674±12±6	6.2k	<sup>2</sup> LEES	14H BABR	$e^+e^- \rightarrow K_S^0 K_L^0 \gamma$
1733±10±10		<sup>3</sup> LEES	12F BABR	10.6 $e^+e^- \rightarrow \phi \pi^+ \pi^- \gamma$
1689±7±10	4.8k	<sup>4</sup> SHEN	09 BELL	10.6 $e^+e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$
1709±20±43		<sup>5</sup> AUBERT	08S BABR	10.6 $e^+e^- \rightarrow$ hadrons
1623±20	948	<sup>6</sup> AKHMETSHIN	03 CMD2	1.05–1.38 $e^+e^- \rightarrow K_L^0 K_S^0$
~1500		<sup>7</sup> ACHASOV	98H RVUE	$e^+e^- \rightarrow \pi^+ \pi^- \pi^0, \omega \pi^+ \pi^-,$ $K^+ K^-$
~1900		<sup>8</sup> ACHASOV	98H RVUE	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$
1700±20		<sup>9</sup> CLEGG	94 RVUE	$e^+e^- \rightarrow K^+ K^-, K_S^0 K \pi$
1657±27	367	BISELLO	91C DM2	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$
1655±17		<sup>10</sup> BISELLO	88B DM2	$e^+e^- \rightarrow K^+ K^-$
1680±10		<sup>11</sup> BUON	82 DM1	$e^+e^- \rightarrow$ hadrons
1677±12		<sup>12</sup> MANE	82 DM1	$e^+e^- \rightarrow K_S^0 K \pi$

<sup>1</sup> Assuming the  $K \bar{K}^*(892) + c.c.$  dynamics. Systematic uncertainties not estimated.

<sup>2</sup> Using a vector meson dominance model with contribution from  $\phi(1020)$ ,  $\phi(1680)$ , and higher mass excitations of  $\rho(770)$  and  $\omega(782)$ .

<sup>3</sup> Using events with  $\pi\pi$  invariant mass less than 0.85 GeV.

<sup>4</sup> From a fit with two incoherent Breit-Wigners.

<sup>5</sup> From the simultaneous fit to the  $K \bar{K}^*(892) + c.c.$  and  $\phi\eta$  data from AUBERT 08S using the results of AUBERT 07AK.

<sup>6</sup> From the combined fit of AKHMETSHIN 03 and MANE 81 also including  $\rho$ ,  $\omega$ , and  $\phi$ . Neither isospin nor flavor structure known.

<sup>7</sup> Using data from IVANOV 81, BARKOV 87, BISELLO 88B, DOLINSKY 91, and ANTONELLI 92.

<sup>8</sup> Using the data from BISELLO 91C.

<sup>9</sup> Using BISELLO 88B and MANE 82 data.

<sup>10</sup> From global fit including  $\rho$ ,  $\omega$ ,  $\phi$  and  $\rho(1700)$  assume mass 1570 MeV and width 510 MeV for  $\rho$  radial excitation.

<sup>11</sup> From global fit of  $\rho$ ,  $\omega$ ,  $\phi$  and their radial excitations to channels  $\omega \pi^+ \pi^-$ ,  $K^+ K^-$ ,  $K_S^0 K_L^0$ ,  $K_S^0 K^\pm \pi^\mp$ . Assume mass 1570 MeV and width 510 MeV for  $\rho$  radial excitations, mass 1570 and width 500 MeV for  $\omega$  radial excitation.

<sup>12</sup> Fit to one channel only, neglecting interference with  $\omega$ ,  $\rho(1700)$ .

### PHOTOPRODUCTION

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1753±3	<sup>1</sup> LINK	02K FOCS	20–160 $\gamma p \rightarrow K^+ K^- p$
1726±22	<sup>1</sup> BUSENITZ	89 TPS	$\gamma p \rightarrow K^+ K^- X$
1760±20	<sup>1</sup> ATKINSON	85C OMEG	20–70 $\gamma p \rightarrow K \bar{K} X$
1690±10	<sup>1</sup> ASTON	81F OMEG	25–70 $\gamma p \rightarrow K^+ K^- X$

<sup>1</sup> We list here a state decaying into  $K^+ K^-$  possibly different from  $\phi(1680)$ .

**$\rho\bar{\rho}$  ANNIHILATION**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1700±8	<sup>1</sup> AMSLER	06	CBAR 0.9 $\bar{p}p \rightarrow K^+ K^- \pi^0$
<sup>1</sup> Could also be $\rho(1700)$ .			

 **$\phi(1680)$  WIDTH** **$e^+e^-$  PRODUCTION**

VALUE (MeV)	EVS	DOCUMENT ID	TECN	COMMENT
<b>150±50 OUR ESTIMATE</b>				This is only an educated guess; the error given is larger than the error on the average of the published values.
• • • We do not use the following data for averages, fits, limits, etc. • • •				
300±50	2k	<sup>1</sup> ACHASOV	18A	SND 1.3–2.0 $e^+e^- \rightarrow K_S^0 K_L^0 \pi^0$
165±38±70	6.2k	<sup>2</sup> LEES	14H	BABR $e^+e^- \rightarrow K_S^0 K_L^0 \gamma$
300±15±37		<sup>3</sup> LEES	12F	BABR 10.6 $e^+e^- \rightarrow \phi \pi^+ \pi^- \gamma$
211±14±19	4.8k	<sup>4</sup> SHEN	09	BELL 10.6 $e^+e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$
322±77±160		<sup>5</sup> AUBERT	08S	BABR 10.6 $e^+e^- \rightarrow$ hadrons
139±60	948	<sup>6</sup> AKHMETSHIN	03	CMD2 1.05–1.38 $e^+e^- \rightarrow K_L^0 K_S^0$
300±60		<sup>7</sup> CLEGG	94	RVUE $e^+e^- \rightarrow K^+ K^-, K_S^0 K \pi$
146±55	367	BISELLO	91C	DM2 $e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$
207±45		<sup>8</sup> BISELLO	88B	DM2 $e^+e^- \rightarrow K^+ K^-$
185±22		<sup>9</sup> BUON	82	DM1 $e^+e^- \rightarrow$ hadrons
102±36		<sup>10</sup> MANE	82	DM1 $e^+e^- \rightarrow K_S^0 K \pi$

<sup>1</sup> Assuming the  $K\bar{K}^*(892) + c.c.$  dynamics. Systematic uncertainties not estimated.

<sup>2</sup> Using a vector meson dominance model with contribution from  $\phi(1020)$ ,  $\phi(1680)$ , and higher mass excitations of  $\rho(770)$  and  $\omega(782)$ .

<sup>3</sup> Using events with  $\pi\pi$  invariant mass less than 0.85 GeV.

<sup>4</sup> From a fit with two incoherent Breit-Wigners.

<sup>5</sup> From the simultaneous fit to the  $K\bar{K}^*(892) + c.c.$  and  $\phi\eta$  data from AUBERT 08S using the results of AUBERT 07AK.

<sup>6</sup> From the combined fit of AKHMETSHIN 03 and MANE 81 also including  $\rho$ ,  $\omega$ , and  $\phi$ . Neither isospin nor flavor structure known.

<sup>7</sup> Using BISELLO 88B and MANE 82 data.

<sup>8</sup> From global fit including  $\rho$ ,  $\omega$ ,  $\phi$  and  $\rho(1700)$ .

<sup>9</sup> From global fit of  $\rho$ ,  $\omega$ ,  $\phi$  and their radial excitations to channels  $\omega\pi^+\pi^-$ ,  $K^+K^-$ ,  $K_S^0 K_L^0$ ,  $K_S^0 K^\pm \pi^\mp$ . Assume mass 1570 MeV and width 510 MeV for  $\rho$  radial excitations, mass 1570 and width 500 MeV for  $\omega$  radial excitation.

<sup>10</sup> Fit to one channel only, neglecting interference with  $\omega$ ,  $\rho(1700)$ .

**PHOTOPRODUCTION**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
122±63	<sup>1</sup> LINK	02K	FOCS 20–160 $\gamma p \rightarrow K^+ K^- p$
121±47	<sup>1</sup> BUSENITZ	89	TPS $\gamma p \rightarrow K^+ K^- X$
80±40	<sup>1</sup> ATKINSON	85C	OMEG 20–70 $\gamma p \rightarrow K\bar{K}X$
100±40	<sup>1</sup> ASTON	81F	OMEG 25–70 $\gamma p \rightarrow K^+ K^- X$

<sup>1</sup> We list here a state decaying into  $K^+ K^-$  possibly different from  $\phi(1680)$ .

## $\rho\bar{\rho}$ ANNIHILATION

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$143 \pm 24$	<sup>1</sup> AMSLER	06	CBAR $0.9 \bar{\rho}\rho \rightarrow K^+ K^- \pi^0$

<sup>1</sup> Could also be  $\rho(1700)$ .

## $\phi(1680)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $K\bar{K}^*(892) + \text{c.c.}$	seen
$\Gamma_2$ $K_S^0 K \pi$	seen
$\Gamma_3$ $K\bar{K}$	seen
$\Gamma_4$ $K_L^0 K_S^0$	
$\Gamma_5$ $e^+ e^-$	seen
$\Gamma_6$ $\omega \pi \pi$	not seen
$\Gamma_7$ $\phi \pi \pi$	
$\Gamma_8$ $K^+ K^- \pi^+ \pi^-$	seen
$\Gamma_9$ $\eta \phi$	seen
$\Gamma_{10}$ $\eta \gamma$	seen
$\Gamma_{11}$ $K^+ K^- \pi^0$	

## $\phi(1680) \Gamma(i)\Gamma(e^+ e^-)/\Gamma(\text{total})$

This combination of a partial width with the partial width into  $e^+ e^-$  and with the total width is obtained from the integrated cross section into channel (I) in  $e^+ e^-$  annihilation. We list only data that have not been used to determine the partial width  $\Gamma(I)$  or the branching ratio  $\Gamma(I)/\text{total}$ .

## $\Gamma(K_L^0 K_S^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_4 \Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
$14.3 \pm 2.4 \pm 6.2$	6.2k	<sup>1</sup> LEES	14H	BABR $e^+ e^- \rightarrow K_S^0 K_L^0 \gamma$

<sup>1</sup> Using a vector meson dominance model with contribution from  $\phi(1020)$ ,  $\phi(1680)$ , and higher mass excitations of  $\rho(770)$  and  $\omega(782)$ .

## $\Gamma(\phi \pi \pi) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_7 \Gamma_5/\Gamma$

VALUE ( $10^{-2}$ keV)	DOCUMENT ID	TECN	COMMENT
$4.2 \pm 0.2 \pm 0.3$	LEES	12F	BABR $10.6 e^+ e^- \rightarrow \phi \pi^+ \pi^- \gamma$

### $\phi(1680) \Gamma(i)\Gamma(e^+e^-)/\Gamma^2(\text{total})$

This combination of a branching ratio into channel (*i*) and branching ratio into  $e^+e^-$  is directly measured and obtained from the cross section at the peak. We list only data that have not been used to determine the branching ratio into (*i*) or  $e^+e^-$ .

$$\Gamma(K_L^0 K_S^0)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_4/\Gamma \times \Gamma_5/\Gamma$$

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.131 \pm 0.059$	948	<sup>1</sup> AKHMETSHIN 03	CMD2	$1.05\text{--}1.38 e^+e^- \rightarrow K_L^0 K_S^0$
-------------------	-----	----------------------------	------	--

<sup>1</sup> From the combined fit of AKHMETSHIN 03 and MANE 81 also including  $\rho$ ,  $\omega$ , and  $\phi$ . Neither isospin nor flavor structure known. Recalculated by us.

$$\Gamma(K\bar{K}^*(892) + \text{c.c.})/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_1/\Gamma \times \Gamma_5/\Gamma$$

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.15 \pm 0.16 \pm 0.01$		<sup>1</sup> AUBERT 08S	BABR	$10.6 e^+e^- \rightarrow K\bar{K}^*(892)\gamma + \text{c.c.}$
$3.29 \pm 1.57$	367	<sup>2</sup> BISELLO 91C	DM2	$1.35\text{--}2.40 e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$

<sup>1</sup> From the simultaneous fit to the  $K\bar{K}^*(892) + \text{c.c.}$  and  $\phi\eta$  data from AUBERT 08S using the results of AUBERT 07AK.

<sup>2</sup> Recalculated by us with the published value of  $B(K\bar{K}^*(892) + \text{c.c.}) \times \Gamma(e^+e^-)$ .

$$\Gamma(\phi\pi\pi)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_7/\Gamma \times \Gamma_5/\Gamma$$

VALUE (units $10^{-7}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.86 \pm 0.14 \pm 0.21$	4.8k	<sup>1</sup> SHEN 09	BELL	$10.6 e^+e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$
--------------------------	------	----------------------	------	--

<sup>1</sup> Multiplied by 3/2 to take into account the  $\phi\pi^0\pi^0$  mode. Using  $B(\phi \rightarrow K^+ K^-) = (49.2 \pm 0.6)\%$ .

$$\Gamma(\eta\phi)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_9/\Gamma \times \Gamma_5/\Gamma$$

VALUE (units $10^{-6}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.43 \pm 0.10 \pm 0.09$	<sup>1</sup> AUBERT 08S	BABR	$10.6 e^+e^- \rightarrow \phi\eta\gamma$
--------------------------	-------------------------	------	--

<sup>1</sup> From the simultaneous fit to the  $K\bar{K}^*(892) + \text{c.c.}$  and  $\phi\eta$  data from AUBERT 08S using the results of AUBERT 07AK.

### $\phi(1680)$ BRANCHING RATIOS

$$\Gamma(K\bar{K}^*(892) + \text{c.c.})/\Gamma(K_S^0 K\pi) \quad \Gamma_1/\Gamma_2$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

dominant	MANE 82	DM1	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$
----------	---------	-----	--

$$\Gamma(K\bar{K})/\Gamma(K\bar{K}^*(892) + \text{c.c.}) \quad \Gamma_3/\Gamma_1$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.07 \pm 0.01$	BUON 82	DM1	$e^+e^-$
-----------------	---------	-----	----------

$\Gamma(\omega\pi\pi)/\Gamma(K\bar{K}^*(892)+c.c.)$   $\Gamma_6/\Gamma_1$

VALUE	DOCUMENT ID	TECN	COMMENT
<0.10	BUON 82	DM1	$e^+e^-$

$\Gamma(\eta\phi)/\Gamma_{total}$   $\Gamma_9/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	35	<sup>1</sup> ACHASOV 14	SND	1.15–2.00 $e^+e^- \rightarrow \eta\gamma$

<sup>1</sup>From a phenomenological model based on vector meson dominance with  $\rho(1450)$  and  $\phi(1680)$  masses and widths from the PDG 12.

$\Gamma(\eta\phi)/\Gamma(K\bar{K}^*(892)+c.c.)$   $\Gamma_9/\Gamma_1$

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			

$\approx 0.37$  <sup>1</sup>AUBERT 08s BABR 10.6  $e^+e^- \rightarrow$  hadrons

<sup>1</sup>From the fit including data from AUBERT 07AK.

$\Gamma(\eta\gamma)/\Gamma_{total}$   $\Gamma_{10}/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	35	<sup>1</sup> ACHASOV 14	SND	1.15–2.00 $e^+e^- \rightarrow \eta\gamma$

<sup>1</sup>From a phenomenological model based on vector meson dominance with  $\rho(1450)$  and  $\phi(1680)$  masses and widths from the PDG 12.

**$\phi(1680)$  REFERENCES**

ACHASOV 18A	PR D97 032011	M.N. Achasov <i>et al.</i>	(SND Collab.)
ACHASOV 14	PR D90 032002	M.N. Achasov <i>et al.</i>	(SND Collab.)
LEES 14H	PR D89 092002	J.P. Lees <i>et al.</i>	(BABAR Collab)
LEES 12F	PR D86 012008	J.P. Lees <i>et al.</i>	(BABAR Collab.)
PDG 12	PR D86 010001	J. Beringer <i>et al.</i>	(PDG Collab.)
SHEN 09	PR D80 031101	C.P. Shen <i>et al.</i>	(BELLE Collab.)
AUBERT 08S	PR D77 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT 07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)
AMSLER 06	PL B639 165	C. Amsler <i>et al.</i>	(CBAR Collab.)
AKHMETSHIN 03	PL B551 27	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
Also	PAN 65 1222	E.V. Anashkin, V.M. Aulchenko, R.R. Akhmetshin	
	Translated from YAF 65 1255.		
LINK 02K	PL B545 50	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
ACHASOV 98H	PR D57 4334	N.N. Achasov, A.A. Kozhevnikov	
CLEGG 94	ZPHY C62 455	A.B. Clegg, A. Donnachie	(LANC, MCHS)
ANTONELLI 92	ZPHY C56 15	A. Antonelli <i>et al.</i>	(DM2 Collab.)
BISELLO 91C	ZPHY C52 227	D. Bisello <i>et al.</i>	(DM2 Collab.)
DOLINSKY 91	PRPL 202 99	S.I. Dolinsky <i>et al.</i>	(NOVO)
BUSENITZ 89	PR D40 1	J.K. Busenitz <i>et al.</i>	(ILL, FNAL)
BISELLO 88B	ZPHY C39 13	D. Bisello <i>et al.</i>	(PADO, CLER, FRAS+)
BARKOV 87	JETPL 46 164	L.M. Barkov <i>et al.</i>	(NOVO)
	Translated from ZETFP 46 132.		
ATKINSON 85C	ZPHY C27 233	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)
BUON 82	PL 118B 221	J. Buon <i>et al.</i>	(LALO, MONP)
MANE 82	PL 112B 178	F. Mane <i>et al.</i>	(LALO)
ASTON 81F	PL 104B 231	D. Aston	(BONN, CERN, EPOL, GLAS, LANC+)
IVANOV 81	PL 107B 297	P.M. Ivanov <i>et al.</i>	(NOVO)
MANE 81	PL 99B 261	F. Mane <i>et al.</i>	(ORSAY)