

$\phi(2170)$ 

$$I^G(J^{PC}) = 0^-(1^{--})$$

Observed by AUBERT, BE 06D in the initial-state radiation process  
 $e^+e^- \rightarrow \phi f_0(980)\gamma$ .

### $\phi(2170)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
2135 ± 8 ± 9	95	ABLIKIM	19I BES3	$e^+e^- \rightarrow \eta\phi f_0(980)$
2239.2 ± 7.1 ± 11.3		<sup>1</sup> ABLIKIM	19L BES3	$e^+e^- \rightarrow K^+K^-$
2200 ± 6 ± 5	471	ABLIKIM	15H BES3	$J/\psi \rightarrow \eta\phi\pi^+\pi^-$
2180 ± 8 ± 8		<sup>2,3</sup> LEES	12F BABR	10.6 $e^+e^- \rightarrow \phi\pi^+\pi^-\gamma$
2079 ± 13 <sup>+79</sup> <sub>-28</sub>	4.8k	<sup>4</sup> SHEN	09 BELL	10.6 $e^+e^- \rightarrow$ $K^+K^-\pi^+\pi^-\gamma$
2186 ± 10 ± 6	52	ABLIKIM	08F BES	$J/\psi \rightarrow \eta\phi f_0(980)$
2125 ± 22 ± 10	483	AUBERT	08S BABR	10.6 $e^+e^- \rightarrow \phi\eta\gamma$
2192 ± 14	116	<sup>5</sup> AUBERT	07AK BABR	10.6 $e^+e^- \rightarrow$ $K^+K^-\pi^+\pi^-\gamma$
2169 ± 20	149	<sup>5</sup> AUBERT	07AK BABR	10.6 $e^+e^- \rightarrow$ $K^+K^-\pi^0\pi^0\gamma$
2175 ± 10 ± 15	201	<sup>3,6</sup> AUBERT, BE	06D BABR	10.6 $e^+e^- \rightarrow$ $K^+K^-\pi\pi\gamma$

<sup>1</sup> The observed structure can be due to both the  $\phi(2170)$  and  $\rho(2150)$ .

<sup>2</sup> Fit includes interference with the  $\phi(1680)$ .

<sup>3</sup> From the  $\phi f_0(980)$  component.

<sup>4</sup> From a fit with two incoherent Breit-Wigners.

<sup>5</sup> From the  $K^+K^- f_0(980)$  component.

<sup>6</sup> Superseded by LEES 12F.

### $\phi(2170)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
104 ± 24 ± 12	95	ABLIKIM	19I BES3	$e^+e^- \rightarrow \eta\phi f_0(980)$
139.8 ± 12.3 ± 20.6		<sup>7</sup> ABLIKIM	19L BES3	$e^+e^- \rightarrow K^+K^-$
104 ± 15 ± 15	471	ABLIKIM	15H BES3	$J/\psi \rightarrow \eta\phi\pi^+\pi^-$
77 ± 15 ± 10		<sup>8,9</sup> LEES	12F BABR	10.6 $e^+e^- \rightarrow \phi\pi^+\pi^-\gamma$
192 ± 23 <sup>+25</sup> <sub>-61</sub>	4.8k	<sup>10</sup> SHEN	09 BELL	10.6 $e^+e^- \rightarrow$ $K^+K^-\pi^+\pi^-\gamma$
65 ± 23 ± 17	52	ABLIKIM	08F BES	$J/\psi \rightarrow \eta\phi f_0(980)$
61 ± 50 ± 13	483	AUBERT	08S BABR	10.6 $e^+e^- \rightarrow \phi\eta\gamma$
71 ± 21	116	<sup>11</sup> AUBERT	07AK BABR	10.6 $e^+e^- \rightarrow$ $K^+K^-\pi^+\pi^-\gamma$
102 ± 27	149	<sup>11</sup> AUBERT	07AK BABR	10.6 $e^+e^- \rightarrow$ $K^+K^-\pi^0\pi^0\gamma$
58 ± 16 ± 20	201	<sup>9,12</sup> AUBERT, BE	06D BABR	10.6 $e^+e^- \rightarrow$ $K^+K^-\pi\pi\gamma$

<sup>7</sup> The observed structure can be due to both the  $\phi(2170)$  and  $\rho(2150)$ .

<sup>8</sup> Fit includes interference with the  $\phi(1680)$ .

<sup>9</sup> From the  $\phi f_0(980)$  component.

<sup>10</sup> From a fit with two incoherent Breit-Wigners.

<sup>11</sup> From the  $K^+ K^- f_0(980)$  component.

<sup>12</sup> Superseded by LEES 12F.

## $\phi(2170)$ DECAY MODES

	Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$	$e^+ e^-$	seen
$\Gamma_2$	$\phi\eta$	
$\Gamma_3$	$\phi\pi\pi$	
$\Gamma_4$	$\phi f_0(980)$	seen
$\Gamma_5$	$K^+ K^- \pi^+ \pi^-$	
$\Gamma_6$	$K^+ K^- f_0(980) \rightarrow K^+ K^- \pi^+ \pi^-$	seen
$\Gamma_7$	$K^+ K^- \pi^0 \pi^0$	
$\Gamma_8$	$K^+ K^- f_0(980) \rightarrow K^+ K^- \pi^0 \pi^0$	seen
$\Gamma_9$	$K^{*0} K^\pm \pi^\mp$	not seen
$\Gamma_{10}$	$K^*(892)^0 \bar{K}^*(892)^0$	not seen

### $\phi(2170) \Gamma(i)\Gamma(e^+ e^-)/\Gamma(\text{total})$

$\Gamma(\phi\eta) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$   $\Gamma_2\Gamma_1/\Gamma$

<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-------------------	-------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.7±0.7±1.3	483	AUBERT	08S	BABR 10.6 $e^+ e^- \rightarrow \phi\eta\gamma$
-------------	-----	--------	-----	--

$\Gamma(\phi f_0(980)) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$   $\Gamma_4\Gamma_1/\Gamma$

<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-------------------	-------------	--------------------	-------------	----------------

**2.3±0.3±0.3**      13,14 LEES      12F BABR 10.6  $e^+ e^- \rightarrow \phi\pi^+\pi^-\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.5±0.8±0.4	201	14,15 AUBERT,BE	06D	BABR 10.6 $e^+ e^- \rightarrow K^+ K^- \pi\pi\gamma$
-------------	-----	-----------------	-----	--

<sup>13</sup> From a fit with constructive interference with the  $\phi(1680)$ . In a fit with destructive interference, the value is larger by a factor of 12.

<sup>14</sup> From the  $\phi f_0(980)$  component.

<sup>15</sup> Superseded by LEES 12F.

### $\phi(2170) \Gamma(i)\Gamma(e^+ e^-)/\Gamma^2(\text{total})$

$\Gamma(\phi\pi\pi)/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$   $\Gamma_3/\Gamma \times \Gamma_1/\Gamma$

<u>VALUE (units 10<sup>-7</sup>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------------------------------	-------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.65±0.15±0.18	4.8k	16 SHEN	09 BELL	10.6 $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$
----------------	------	---------	---------	---

<sup>16</sup> Multiplied by 3/2 to take into account the  $\phi\pi^0\pi^0$  mode. Using  $B(\phi \rightarrow K^+ K^-) = (49.2 \pm 0.6)\%$ .

### $\phi(2170)$ BRANCHING RATIOS

$$\Gamma(K^+ K^- f_0(980) \rightarrow K^+ K^- \pi^+ \pi^-) / \Gamma_{\text{total}} \quad \Gamma_6 / \Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>seen</b>	AUBERT	07AK BABR	10.6 $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$

$$\Gamma(K^+ K^- f_0(980) \rightarrow K^+ K^- \pi^0 \pi^0) / \Gamma_{\text{total}} \quad \Gamma_8 / \Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>seen</b>	AUBERT	07AK BABR	10.6 $e^+ e^- \rightarrow K^+ K^- \pi^0 \pi^0 \gamma$

$$\Gamma(K^{*0} K^\pm \pi^\mp) / \Gamma_{\text{total}} \quad \Gamma_9 / \Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>not seen</b>	AUBERT	07AK BABR	10.6 GeV $e^+ e^-$

$$\Gamma(K^*(892)^0 \bar{K}^*(892)^0) / \Gamma_{\text{total}} \quad \Gamma_{10} / \Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>not seen</b>	ABLIKIM	10C BES2	$J/\psi \rightarrow \eta K^+ \pi^- K^- \pi^+$

### $\phi(2170)$ REFERENCES

ABLIKIM	19I	PR D99 012014	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	19L	PR D99 032001	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABLIKIM	15H	PR D91 052017	M. Ablikim <i>et al.</i>	(BES III Collab.)
LEES	12F	PR D86 012008	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	10C	PL B685 27	M. Ablikim <i>et al.</i>	(BES II Collab.)
SHEN	09	PR D80 031101	C.P. Shen <i>et al.</i>	(BELLE Collab.)
ABLIKIM	08F	PRL 100 102003	M. Ablikim <i>et al.</i>	(BES Collab.)
AUBERT	08S	PR D77 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,BE	06D	PR D74 091103	B. Aubert <i>et al.</i>	(BABAR Collab.)