

$\psi(4660)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

 I needs confirmation.also known as $Y(4660)$; was $X(4660)$

This state shows properties different from a conventional $q\bar{q}$ state.
A candidate for an exotic structure. See the review on non- $q\bar{q}$ states.

Seen in radiative return from e^+e^- collisions at $\sqrt{s} = 9.54\text{--}10.58$ GeV by WANG 07D. Also obtained in a combined fit of WANG 07D, AUBERT 07S, and LEES 14F. See also the review on "Spectroscopy of mesons containing two heavy quarks."

 $\psi(4660)$ MASS

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------------------------------------------------------------------------------------|--------------------|-------------------------------------|------|-----------------------------------------------------|
| 4643 ± 9 | OUR AVERAGE | Error includes scale factor of 1.2. | | |
| 4652 ± 10 ± 11 | 279 | ¹ WANG 15A | BELL | 10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$ |
| 4669 ± 21 ± 3 | 37 | ² LEES 14F | BABR | 10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$ |
| 4634 $\begin{smallmatrix} +8 \\ -7 \end{smallmatrix}$ $\begin{smallmatrix} +5 \\ -8 \end{smallmatrix}$ | 142 | ³ PAKHLOVA 08B | BELL | $e^+e^- \rightarrow \Lambda_c^+\Lambda_c^-$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| 4652.5 ± 3.4 ± 1.1 | | ⁴ DAI 17 | RVUE | $e^+e^- \rightarrow \Lambda_c^+\Lambda_c^-$ |
| 4645.2 ± 9.5 ± 6.0 | | ⁵ ZHANG 17B | RVUE | $e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$ |
| 4646.4 ± 9.7 ± 4.8 | | ⁶ ZHANG 17C | RVUE | $e^+e^- \rightarrow \pi^+\pi^-J/\psi$ or $\psi(2S)$ |
| 4661 $\begin{smallmatrix} +9 \\ -8 \end{smallmatrix}$ ± 6 | 44 | ⁷ LIU 08H | RVUE | 10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$ |
| 4664 ± 11 ± 5 | 44 | WANG 07D | BELL | 10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$ |

¹ From a two-resonance fit. Supersedes WANG 07D.² From a two-resonance fit.³ The $\pi^+\pi^-\psi(2S)$ and $\Lambda_c^+\Lambda_c^-$ states are not necessarily the same.⁴ The pole parameters are extracted from the speed plot.⁵ From a three-resonance fit.⁶ From a combined fit of BELLE, BABAR and BES3 $e^+e^- \rightarrow \pi^+\pi^-J/\psi$ and $e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$ data.⁷ From a combined fit of AUBERT 07S and WANG 07D data with two resonances. **$\psi(4660)$ WIDTH**

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------------------------------------------------------------------------------------------|--------------------|---------------------------|------|-----------------------------------------------------|
| 72 ± 11 | OUR AVERAGE | | | |
| 68 ± 11 ± 5 | 279 | ¹ WANG 15A | BELL | 10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$ |
| 104 ± 48 ± 10 | 37 | ² LEES 14F | BABR | 10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$ |
| 92 $\begin{smallmatrix} +40 \\ -24 \end{smallmatrix}$ $\begin{smallmatrix} +10 \\ -21 \end{smallmatrix}$ | 142 | ³ PAKHLOVA 08B | BELL | $e^+e^- \rightarrow \Lambda_c^+\Lambda_c^-$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| 62.6 ± 5.6 ± 4.3 | | ⁴ DAI 17 | RVUE | $e^+e^- \rightarrow \Lambda_c^+\Lambda_c^-$ |
| 113.8 ± 18.1 ± 3.4 | | ⁵ ZHANG 17B | RVUE | $e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$ |
| 103.5 ± 15.6 ± 4.0 | | ⁶ ZHANG 17C | RVUE | $e^+e^- \rightarrow \pi^+\pi^-J/\psi$ or $\psi(2S)$ |
| 42 $\begin{smallmatrix} +17 \\ -12 \end{smallmatrix}$ ± 6 | 44 | ⁷ LIU 08H | RVUE | 10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$ |
| 48 ± 15 ± 3 | 44 | WANG 07D | BELL | 10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$ |

- ¹ From a two-resonance fit. Supersedes WANG 07D.
² From a two-resonance fit.
³ The $\pi^+\pi^-\psi(2S)$ and $\Lambda_c^+\Lambda_c^-$ states are not necessarily the same.
⁴ The pole parameters are extracted from the speed plot.
⁵ From a three-resonance fit.
⁶ From a combined fit of BELLE, BABAR and BES3 $e^+e^- \rightarrow \pi^+\pi^-J/\psi$ and $e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$ data.
⁷ From a combined fit of AUBERT 07S and WANG 07D data with two resonances.

$\psi(4660)$ DECAY MODES

| Mode | Fraction (Γ_i/Γ) |
|-------------------------------------|--------------------------------|
| Γ_1 e^+e^- | |
| Γ_2 $\psi(2S)\pi^+\pi^-$ | seen |
| Γ_3 $J/\psi\eta$ | |
| Γ_4 $D^0D^{*-}\pi^+$ | |
| Γ_5 $\chi_{c1}\gamma$ | |
| Γ_6 $\chi_{c2}\gamma$ | |
| Γ_7 $\Lambda_c^+\Lambda_c^-$ | |

$\psi(4660) \Gamma(i) \times \Gamma(e^+e^-)/\Gamma(\text{total})$

$\Gamma(\psi(2S)\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_2\Gamma_1/\Gamma$

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------------------------------------------------------------------------------|------|-------------------|----------|-----------------------------------------------------|
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| $2.0 \pm 0.3 \pm 0.2$ | 279 | ¹ WANG | 15A BELL | $10.58 e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$ |
| $8.1 \pm 1.1 \pm 1.0$ | 279 | ² WANG | 15A BELL | $10.58 e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$ |
| $2.7 \pm 1.3 \pm 0.5$ | 37 | ³ LEES | 14F BABR | $10.58 e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$ |
| $7.5 \pm 1.7 \pm 0.7$ | 37 | ⁴ LEES | 14F BABR | $10.58 e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$ |
| $2.2^{+0.7}_{-0.6}$ | 44 | ⁵ LIU | 08H RVUE | $10.58 e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$ |
| 5.9 ± 1.6 | 44 | ⁶ LIU | 08H RVUE | $10.58 e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$ |
| $3.0 \pm 0.9 \pm 0.3$ | 44 | ³ WANG | 07D BELL | $10.58 e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$ |
| $7.6 \pm 1.8 \pm 0.8$ | 44 | ⁴ WANG | 07D BELL | $10.58 e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$ |

- ¹ Solution I of two equivalent solutions from a fit using two interfering resonances. Supersedes WANG 07D.
² Solution II of two equivalent solutions from a fit using two interfering resonances. Supersedes WANG 07D.
³ Solution I of two equivalent solutions in a fit using two interfering resonances.
⁴ Solution II of two equivalent solutions in a fit using two interfering resonances.
⁵ Solution I in a combined fit of AUBERT 07S and WANG 07D data with two resonances.
⁶ Solution II in a combined fit of AUBERT 07S and WANG 07D data with two resonances.

$\Gamma(J/\psi\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_3\Gamma_1/\Gamma$

| VALUE (eV) | CL% | DOCUMENT ID | TECN | COMMENT |
|-------------------------------------------------------------------------------|-----|-------------|----------|---------------------------------------|
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| <0.94 | 90 | WANG | 13B BELL | $e^+e^- \rightarrow J/\psi\eta\gamma$ |

| $\Gamma(\chi_{c1}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ | | | | | $\Gamma_5\Gamma_1/\Gamma$ |
|-----------------------------------------------------------------------|-----|------------------|------|---------|--------------------------------------------|
| VALUE (eV) | CL% | DOCUMENT ID | TECN | COMMENT | |
| <0.45 | 90 | ¹ HAN | 15 | BELL | 10.58 $e^+e^- \rightarrow \chi_{c1}\gamma$ |

¹ Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.

| $\Gamma(\chi_{c2}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ | | | | | $\Gamma_6\Gamma_1/\Gamma$ |
|-----------------------------------------------------------------------|-----|------------------|------|---------|--------------------------------------------|
| VALUE (eV) | CL% | DOCUMENT ID | TECN | COMMENT | |
| <2.1 | 90 | ¹ HAN | 15 | BELL | 10.58 $e^+e^- \rightarrow \chi_{c2}\gamma$ |

¹ Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.

$\psi(4660)$ BRANCHING RATIOS

| $\Gamma(D^0 D^{*-} \pi^+)/\Gamma(\psi(2S)\pi^+\pi^-)$ | | | | | Γ_4/Γ_2 |
|-------------------------------------------------------|-----|-------------|------|---------------------------------------|---------------------|
| VALUE | CL% | DOCUMENT ID | TECN | COMMENT | |
| <10 | 90 | PAKHLOVA 09 | BELL | $e^+e^- \rightarrow D^0 D^{*-} \pi^+$ | |

| $\Gamma(D^0 D^{*-} \pi^+)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ | | | | | $\Gamma_4/\Gamma \times \Gamma_1/\Gamma$ |
|----------------------------------------------------------------------------------------------|-----|--------------------------|------|---------------------------------------|------------------------------------------|
| VALUE | CL% | DOCUMENT ID | TECN | COMMENT | |
| <0.37 $\times 10^{-6}$ | 90 | ¹ PAKHLOVA 09 | BELL | $e^+e^- \rightarrow D^0 D^{*-} \pi^+$ | |

¹ Using $4664 \pm 11 \pm 5$ MeV for the mass of $\psi(4660)$.

| $\Gamma(\Lambda_c^+ \Lambda_c^-)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ | | | | | $\Gamma_7/\Gamma \times \Gamma_1/\Gamma$ |
|-----------------------------------------------------------------------------------------------------|------|---------------------------|------|----------------------------------------------|------------------------------------------|
| VALUE (units 10^{-6}) | EVTS | DOCUMENT ID | TECN | COMMENT | |
| $0.68^{+0.16+0.29}_{-0.15-0.30}$ | 142 | ¹ PAKHLOVA 08B | BELL | $e^+e^- \rightarrow \Lambda_c^+ \Lambda_c^-$ | |

¹ The $\pi^+\pi^-\psi(2S)$ and $\Lambda_c^+\Lambda_c^-$ states are not necessarily the same.

$\psi(4660)$ REFERENCES

| | | | | |
|----------|-----|----------------|-------------------------------------------|-----------------|
| DAI | 17 | PR D96 116001 | L.-Y. Dai, J. Haidenbauer, U.-G. Meissner | (JULI+) |
| ZHANG | 17B | PR D96 054008 | J. Zhang, J. Zhang | |
| ZHANG | 17C | EPJ C77 727 | J. Zhang, L. Yuan | |
| HAN | 15 | PR D92 012011 | Y.L. Han <i>et al.</i> | (BELLE Collab.) |
| WANG | 15A | PR D91 112007 | X.L. Wang <i>et al.</i> | (BELLE Collab.) |
| LEES | 14F | PR D89 111103 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| WANG | 13B | PR D87 051101 | X.L. Wang <i>et al.</i> | (BELLE Collab.) |
| PAKHLOVA | 09 | PR D80 091101 | G. Pakhlova <i>et al.</i> | (BELLE Collab.) |
| LIU | 08H | PR D78 014032 | Z.Q. Liu, X.S. Qin, C.Z. Yuan | |
| PAKHLOVA | 08B | PRL 101 172001 | C. Pakhlova <i>et al.</i> | (BELLE Collab.) |
| AUBERT | 07S | PRL 98 212001 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| WANG | 07D | PRL 99 142002 | X.L. Wang <i>et al.</i> | (BELLE Collab.) |