

$\Sigma(1770) 1/2^+$ $I(J^P) = 1(\frac{1}{2}^+)$ Status: *

OMITTED FROM SUMMARY TABLE

Evidence for this state now rests solely on solution 1 of BAILLON 75, (see the footnotes) but the $\Lambda\pi$ partial-wave amplitudes of this solution are in disagreement with amplitudes from most other $\Lambda\pi$ analyses. ZHANG 13A finds no evidence for this state.

 $\Sigma(1770)$ POLE POSITION**REAL PART**

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|--------------------|---------------------|------|-------------------|
| 1706^{+67}_{-60} | ¹ KAMANO | 15 | DPWA Multichannel |

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹From the preferred solution A in KAMANO 15. Solution B Reports two poles at 1605^{+2}_{-4} and 2014^{+6}_{-13} MeV.

−2×IMAGINARY PART

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|--------------------|---------------------|------|-------------------|
| 101^{+158}_{-84} | ¹ KAMANO | 15 | DPWA Multichannel |

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹From the preferred solution A in KAMANO 15. Solution B reports two poles with 192^{+2}_{-10} and 140^{+28}_{-2} MeV width.

 $\Sigma(1770)$ POLE RESIDUES

The normalized residue is the residue divided by $\Gamma_{pole}/2$.

Normalized residue in $N\bar{K} \rightarrow \Sigma(1770) \rightarrow N\bar{K}$

| MODULUS | PHASE (°) | DOCUMENT ID | TECN | COMMENT |
|---------|-----------|---------------------|------|-------------------|
| 0.0268 | 91 | ¹ KAMANO | 15 | DPWA Multichannel |

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹From the preferred solution A in KAMANO 15.

Normalized residue in $N\bar{K} \rightarrow \Sigma(1770) \rightarrow \Sigma\pi$

| MODULUS | PHASE (°) | DOCUMENT ID | TECN | COMMENT |
|---------|-----------|---------------------|------|-------------------|
| 0.145 | −171 | ¹ KAMANO | 15 | DPWA Multichannel |

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹From the preferred solution A in KAMANO 15.

Normalized residue in $N\bar{K} \rightarrow \Sigma(1770) \rightarrow \Lambda\pi$

| MODULUS | PHASE (°) | DOCUMENT ID | TECN | COMMENT |
|---------|-----------|---------------------|------|-------------------|
| 0.117 | −76 | ¹ KAMANO | 15 | DPWA Multichannel |

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹From the preferred solution A in KAMANO 15.

Normalized residue in $N\bar{K} \rightarrow \Sigma(1770) \rightarrow \Sigma(1385)\pi$

| <u>MODULUS</u> | <u>PHASE ($^\circ$)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------|------------------------------------|--------------------|-------------|----------------|
|----------------|------------------------------------|--------------------|-------------|----------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|--------|------|---------------------|----|-------------------|
| 0.0722 | -128 | ¹ KAMANO | 15 | DPWA Multichannel |
|--------|------|---------------------|----|-------------------|

¹From the preferred solution A in KAMANO 15.

 $\Sigma(1770)$ MASS

| <u>VALUE (MeV)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------|--------------------|-------------|----------------|
|--------------------|--------------------|-------------|----------------|

 ≈ 1770 OUR ESTIMATE

| | | | |
|---------------|----------------------|----|--|
| 1738 ± 10 | ¹ GOPAL | 77 | DPWA $\bar{K}N$ multichannel |
| 1770 ± 20 | ² BAILLON | 75 | IPWA $\bar{K}N \rightarrow \Lambda\pi$ |
| 1772 | ³ KANE | 72 | DPWA $K^-p \rightarrow \Sigma\pi$ |

¹Required to fit the isospin-1 total cross section of CARROLL 76 in the $\bar{K}N$ channel. The addition of new K^-p polarization and K^-n differential cross-section data in GOPAL 80 find it to be more consistent with the $\Sigma(1660) P_{11}$.

²From solution 1 of BAILLON 75; not present in solution 2.

³Not required in KANE 74, which supersedes KANE 72.

 $\Sigma(1770)$ WIDTH

| <u>VALUE (MeV)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------|--------------------|-------------|----------------|
|--------------------|--------------------|-------------|----------------|

| | | | |
|-------------|----------------------|----|--|
| 72 ± 10 | ¹ GOPAL | 77 | DPWA $\bar{K}N$ multichannel |
| 80 ± 30 | ² BAILLON | 75 | IPWA $\bar{K}N \rightarrow \Lambda\pi$ |
| 80 | ³ KANE | 72 | DPWA $K^-p \rightarrow \Sigma\pi$ |

¹Required to fit the isospin-1 total cross section of CARROLL 76 in the $\bar{K}N$ channel. The addition of new K^-p polarization and K^-n differential cross-section data in GOPAL 80 find it to be more consistent with the $\Sigma(1660) P_{11}$.

²From solution 1 of BAILLON 75; not present in solution 2.

³Not required in KANE 74, which supersedes KANE 72.

 $\Sigma(1770)$ DECAY MODES

| | <u>Mode</u> |
|------------|---|
| Γ_1 | $N\bar{K}$ |
| Γ_2 | $\Lambda\pi$ |
| Γ_3 | $\Sigma\pi$ |
| Γ_4 | $\Sigma(1385)\pi$ |
| Γ_5 | $N\bar{K}^*(892)$, $S=1/2$, P -wave |
| Γ_6 | $N\bar{K}^*(892)$, $S=3/2$, P -wave |

$\Sigma(1770)$ BRANCHING RATIOS

See “Sign conventions for resonance couplings” in the Note on Λ and Σ Resonances.

 $\Gamma(N\bar{K})/\Gamma_{\text{total}}$ Γ_1/Γ

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------|---------------------|-------------|------------------------------|
| 0.14 \pm 0.04 | ¹ GOPAL | 77 | DPWA $\bar{K}N$ multichannel |
| 0.016 | ² KAMANO | 15 | DPWA Multichannel |

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ Required to fit the isospin-1 total cross section of CARROLL 76 in the $\bar{K}N$ channel. The addition of new K^-p polarization and K^-n differential cross-section data in GOPAL 80 find it to be more consistent with the $\Sigma(1660) P_{11}$.

² From the preferred solution A in KAMANO 15.

 $\Gamma(\Lambda\pi)/\Gamma_{\text{total}}$ Γ_2/Γ

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------|---------------------|-------------|-------------------|
| 0.283 | ¹ KAMANO | 15 | DPWA Multichannel |

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ From the preferred solution A in KAMANO 15.

 $\Gamma(\Sigma\pi)/\Gamma_{\text{total}}$ Γ_3/Γ

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------|---------------------|-------------|-------------------|
| 0.595 | ¹ KAMANO | 15 | DPWA Multichannel |

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ From the preferred solution A in KAMANO 15.

 $\Gamma(\Sigma(1385)\pi)/\Gamma_{\text{total}}$ Γ_4/Γ

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------|---------------------|-------------|-------------------|
| 0.103 | ¹ KAMANO | 15 | DPWA Multichannel |

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ From the preferred solution A in KAMANO 15.

 $\Gamma(N\bar{K}^*(892), S=1/2, P\text{-wave})/\Gamma_{\text{total}}$ Γ_5/Γ

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------|---------------------|-------------|-------------------|
| 0.004 | ¹ KAMANO | 15 | DPWA Multichannel |

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ From the preferred solution A in KAMANO 15.

 $\Gamma(N\bar{K}^*(892), S=3/2, P\text{-wave})/\Gamma_{\text{total}}$ Γ_6/Γ

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------|---------------------|-------------|-------------------|
| not seen | ¹ KAMANO | 15 | DPWA Multichannel |

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ From the preferred solution A in KAMANO 15.

| $(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(1770) \rightarrow \Lambda\pi$ | | | | $(\Gamma_1 \Gamma_2)^{1/2} / \Gamma$ |
|---|----------------------|------|---------|--------------------------------------|
| VALUE | DOCUMENT ID | TECN | COMMENT | |
| < 0.04 | GOPAL | 77 | DPWA | $\bar{K}N$ multichannel |
| -0.08 ± 0.02 | ¹ BAILLON | 75 | IPWA | $\bar{K}N \rightarrow \Lambda\pi$ |

¹ From solution 1 of BAILLON 75; not present in solution 2.

| $(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(1770) \rightarrow \Sigma\pi$ | | | | $(\Gamma_1 \Gamma_3)^{1/2} / \Gamma$ |
|--|-------------------|------|---------|--------------------------------------|
| VALUE | DOCUMENT ID | TECN | COMMENT | |
| < 0.04 | GOPAL | 77 | DPWA | $\bar{K}N$ multichannel |
| -0.108 | ¹ KANE | 72 | DPWA | $K^- p \rightarrow \Sigma\pi$ |

¹ Not required in KANE 74, which supersedes KANE 72.

$\Sigma(1770)$ REFERENCES

| | | | | |
|---------|-----|-------------------|-------------------------------|------------------|
| KAMANO | 15 | PR C92 025205 | H. Kamano <i>et al.</i> | (ANL, OSAK) |
| ZHANG | 13A | PR C88 035205 | H. Zhang <i>et al.</i> | (KSU) |
| GOPAL | 80 | Toronto Conf. 159 | G.P. Gopal | (RHEL) |
| GOPAL | 77 | NP B119 362 | G.P. Gopal <i>et al.</i> | (LOIC, RHEL) IJP |
| CARROLL | 76 | PRL 37 806 | A.S. Carroll <i>et al.</i> | (BNL) I |
| BAILLON | 75 | NP B94 39 | P.H. Baillon, P.J. Litchfield | (CERN, RHEL) IJP |
| KANE | 74 | LBL-2452 | D.F. Kane | (LBL) IJP |
| KANE | 72 | PR D5 1583 | D.F.J. Kane | (LBL) |