

$f_2(1270)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

$f_2(1270)$ MASS

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-------|--------------------------|----------|--|
| 1275.5 ± 0.8 OUR AVERAGE | | | | |
| 1275.8 ± 1.0 ± 0.4 | | ¹ BOGOLYUB... | 13 SPEC | $7\pi^+(K^+,p)A \rightarrow n\gamma + X$ |
| 1262 $\pm \frac{1}{2}$ ± 8 | | ² ABLIKIM | 06v BES2 | $e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$ |
| 1275 ± 15 | | ABLIKIM | 05 BES2 | $J/\psi \rightarrow \phi\pi^+\pi^-$ |
| 1283 ± 5 | | ALDE | 98 GAM4 | $100\pi^-p \rightarrow \pi^0\pi^0n$ |
| 1278 ± 5 | | ³ BERTIN | 97c OBLX | $0.0\bar{p}p \rightarrow \pi^+\pi^-\pi^0$ |
| 1272 ± 8 | 200k | PROKOSHKIN | 94 GAM2 | $38\pi^-p \rightarrow \pi^0\pi^0n$ |
| 1269.7 ± 5.2 | 5730 | AUGUSTIN | 89 DM2 | $e^+e^- \rightarrow 5\pi$ |
| 1283 ± 8 | 400 | ⁴ ALDE | 87 GAM4 | $100\pi^-p \rightarrow 4\pi^0n$ |
| 1274 ± 5 | | ⁴ AUGUSTIN | 87 DM2 | $J/\psi \rightarrow \gamma\pi^+\pi^-$ |
| 1283 ± 6 | | ⁵ LONGACRE | 86 MPS | $22\pi^-p \rightarrow n2K_S^0$ |
| 1276 ± 7 | | COURAU | 84 DLCO | $e^+e^- \rightarrow e^+e^-\pi^+\pi^-$ |
| 1273.3 ± 2.3 | | ⁶ CHABAUD | 83 ASPK | $17\pi^-p$ polarized |
| 1280 ± 4 | | ⁷ CASON | 82 STRC | $8\pi^+p \rightarrow \Delta^{++}\pi^0\pi^0$ |
| 1281 ± 7 | 11600 | GIDAL | 81 MRK2 | J/ψ decay |
| 1282 ± 5 | | ⁸ CORDEN | 79 OMEG | $12-15\pi^-p \rightarrow n2\pi$ |
| 1269 ± 4 | 10k | APEL | 75 NICE | $40\pi^-p \rightarrow n2\pi^0$ |
| 1272 ± 4 | 4600 | ENGLER | 74 DBC | $6\pi^+n \rightarrow \pi^+\pi^-p$ |
| 1277 ± 4 | 5300 | FLATTE | 71 HBC | $7.0\pi^+p$ |
| 1273 ± 8 | | ⁴ STUNTEBECK | 70 HBC | $8\pi^-p, 5.4\pi^+d$ |
| 1265 ± 8 | | BOESEBECK | 68 HBC | $8\pi^+p$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| 1263.3 ± 0.2 ± 1.5 | | ⁹ ALBRECHT | 20 RVUE | $0.9\bar{p}p \rightarrow \pi^0\pi^0\eta, \pi^0\eta\eta, \pi^0K^+K^-$ |
| 1259 ± 4 ± 4 | 1.7k | ^{10,11} DOBBS | 15 | $J/\psi \rightarrow \gamma\pi^+\pi^-$ |
| 1267 ± 4 ± 3 | 1.5k | ^{10,11} DOBBS | 15 | $\psi(2S) \rightarrow \gamma\pi^+\pi^-$ |
| 1270 ± 8 | | ¹² ANISOVICH | 09 RVUE | $0.0\bar{p}p, \pi N$ |
| 1277 ± 6 | 870 | ¹³ SCHEGELSKY | 06A RVUE | $\gamma\gamma \rightarrow K_S^0K_S^0$ |
| 1251 ± 10 | | TIKHOMIROV | 03 SPEC | $40.0\pi^-C \rightarrow K_S^0K_S^0K_L^0X$ |
| 1260 ± 10 | | ¹⁴ ALDE | 97 GAM2 | $450pp \rightarrow pp\pi^0\pi^0$ |
| 1278 ± 6 | | ¹⁴ GRYGOREV | 96 SPEC | $40\pi^-N \rightarrow K_S^0K_S^0X$ |
| 1262 ± 11 | | AGUILAR-... | 91 EHS | $400pp$ |
| 1275 ± 10 | | AKER | 91 CBAR | $0.0\bar{p}p \rightarrow 3\pi^0$ |
| 1220 ± 10 | | BREAKSTONE | 90 SFM | $pp \rightarrow pp\pi^+\pi^-$ |
| 1288 ± 12 | | ABACHI | 86B HRS | $e^+e^- \rightarrow \pi^+\pi^-X$ |
| 1284 ± 30 | 3k | BINON | 83 GAM2 | $38\pi^-p \rightarrow n2\eta$ |
| 1280 ± 20 | 3k | APEL | 82 CNTR | $25\pi^-p \rightarrow n2\pi^0$ |
| 1284 ± 10 | 16000 | DEUTSCH... | 76 HBC | $16\pi^+p$ |
| 1258 ± 10 | 600 | TAKAHASHI | 72 HBC | $8\pi^-p \rightarrow n2\pi$ |
| 1275 ± 13 | | ARMENISE | 70 HBC | $9\pi^+n \rightarrow p\pi^+\pi^-$ |
| 1261 ± 5 | 1960 | ⁴ ARMENISE | 68 DBC | $5.1\pi^+n \rightarrow p\pi^+MM^-$ |

| | | | | | |
|-----------|-----|-----------------------|----|-----|--|
| 1270 ± 10 | 360 | ⁴ ARMENISE | 68 | DBC | 5.1 π ⁺ n → p π ⁰ MM |
| 1268 ± 6 | | ¹⁵ JOHNSON | 68 | HBC | 3.7–4.2 π ⁻ p |

¹ Averaged over six nuclear targets, no statistically significant dependence on target nucleus observed.

² Breit-Wigner mass.

³ T-matrix pole.

⁴ Mass errors enlarged by us to Γ/√N; see the note with the K*(892) mass.

⁵ From a partial-wave analysis of data using a K-matrix formalism with 5 poles.

⁶ From an energy-independent partial-wave analysis.

⁷ From an amplitude analysis of the reaction π⁺π⁻ → 2π⁰.

⁸ From an amplitude analysis of π⁺π⁻ → π⁺π⁻ scattering data.

⁹ T-matrix pole, 5 poles, 5 channels, including scattering data from HYAMS 75 (ππ), LONGACRE 86 (K \bar{K}), BINON 83 (ηη).

¹⁰ Using CLEO-c data but not authored by the CLEO Collaboration.

¹¹ From a fit to a Breit-Wigner line shape with fixed Γ = 185 MeV.

¹² 4-poles, 5-channel K matrix fit.

¹³ From analysis of L3 data at 91 and 183–209 GeV.


¹⁴ Systematic uncertainties not estimated.

¹⁵ JOHNSON 68 includes BONDAR 63, LEE 64, DERADO 65, EISNER 67.

f₂(1270) WIDTH

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-------|--------------------------|----------|--|
| 186.7⁺_{-2.5} | | OUR FIT | | Error includes scale factor of 1.4. |
| 185.9⁺_{-2.1} | | OUR AVERAGE | | Error includes scale factor of 1.6. See the ideogram below. |
| 190.3 ± 1.9 ± 1.8 | | ¹ BOGOLYUB... | 13 SPEC | 7π ⁺ (K ⁺ ,p)A → nγ + X |
| 175 ⁺ ₋₄ ± 10 | | ² ABLIKIM | 06v BES2 | e ⁺ e ⁻ → J/ψ → γπ ⁺ π ⁻ |
| 190 ± 20 | | ABLIKIM | 05 BES2 | J/ψ → φπ ⁺ π ⁻ |
| 171 ± 10 | | ALDE | 98 GAM4 | 100 π ⁻ p → π ⁰ π ⁰ n |
| 204 ± 20 | | ³ BERTIN | 97c OBLX | 0.0 $\bar{p}p$ → π ⁺ π ⁻ π ⁰ |
| 192 ± 5 | 200k | PROKOSHKIN | 94 GAM2 | 38 π ⁻ p → π ⁰ π ⁰ n |
| 180 ± 24 | | AGUILAR-... | 91 EHS | 400 pp |
| 169 ± 9 | 5730 | ⁴ AUGUSTIN | 89 DM2 | e ⁺ e ⁻ → 5π |
| 150 ± 30 | 400 | ⁴ ALDE | 87 GAM4 | 100 π ⁻ p → 4π ⁰ n |
| 186 ⁺ ₋₂ | | ⁵ LONGACRE | 86 MPS | 22 π ⁻ p → n2K _S ⁰ |
| 179.2 ⁺ _{-6.6} | | ⁶ CHABAUD | 83 ASPK | 17 π ⁻ p polarized |
| 160 ± 11 | | DENNEY | 83 LASS | 10 π ⁺ N |
| 196 ± 10 | 3k | APEL | 82 CNTR | 25 π ⁻ p → n2π ⁰ |
| 152 ± 9 | | ⁷ CASON | 82 STRC | 8 π ⁺ p → Δ ⁺⁺ π ⁰ π ⁰ |
| 186 ± 27 | 11600 | GIDAL | 81 MRK2 | J/ψ decay |
| 216 ± 13 | | ⁸ CORDEN | 79 OMEG | 12–15 π ⁻ p → n2π |
| 190 ± 10 | 10k | APEL | 75 NICE | 40 π ⁻ p → n2π ⁰ |
| 192 ± 16 | 4600 | ENGLER | 74 DBC | 6 π ⁺ n → π ⁺ π ⁻ p |
| 183 ± 15 | 5300 | FLATTE | 71 HBC | 7 π ⁺ p → Δ ⁺⁺ f ₂ |
| 196 ± 30 | | ⁴ STUNTEBECK | 70 HBC | 8 π ⁻ p, 5.4 π ⁺ d |

| | | | | | |
|---|-------|---------------|-----|------|--|
| 216 ± 20 | 1960 | 4 ARMENISE | 68 | DBC | 5.1 $\pi^+ n \rightarrow p \pi^+ \text{MM}^-$ |
| 128 ± 27 | | 4 BOESEBECK | 68 | HBC | 8 $\pi^+ p$ |
| 176 ± 21 | | 4,9 JOHNSON | 68 | HBC | 3.7-4.2 $\pi^- p$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | | |
| 193.7 ± 0.4 ± 1.6 | | 10 ALBRECHT | 20 | RVUE | 0.9 $\bar{p} p \rightarrow \pi^0 \pi^0 \eta, \pi^0 \eta \eta, \pi^0 K^+ K^-$ |
| 194 ± 36 | | 11 ANISOVICH | 09 | RVUE | 0.0 $\bar{p} p, \pi N$ |
| 195 ± 15 | 870 | 12 SCHEGELSKY | 06A | RVUE | $\gamma \gamma \rightarrow K_S^0 K_S^0$ |
| 121 ± 26 | | TIKHOMIROV | 03 | SPEC | 40.0 $\pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$ |
| 187 ± 20 | | 13 ALDE | 97 | GAM2 | 450 $pp \rightarrow pp \pi^0 \pi^0$ |
| 184 ± 10 | | 13 GRYGOREV | 96 | SPEC | 40 $\pi^- N \rightarrow K_S^0 K_S^0 X$ |
| 200 ± 10 | | AKER | 91 | CBAR | 0.0 $\bar{p} p \rightarrow 3\pi^0$ |
| 240 ± 40 | 3k | BINON | 83 | GAM2 | 38 $\pi^- p \rightarrow n 2\eta$ |
| 187 ± 30 | 650 | 4 ANTIPOV | 77 | CIBS | 25 $\pi^- p \rightarrow p 3\pi$ |
| 225 ± 38 | 16000 | DEUTSCH... | 76 | HBC | 16 $\pi^+ p$ |
| 166 ± 28 | 600 | 4 TAKAHASHI | 72 | HBC | 8 $\pi^- p \rightarrow n 2\pi$ |
| 173 ± 53 | | 4 ARMENISE | 70 | HBC | 9 $\pi^+ n \rightarrow p \pi^+ \pi^-$ |



- ¹ Averaged over six nuclear targets, no statistically significant dependence on target nucleus observed.
- ² Breit-Wigner width
- ³ T-matrix pole.
- ⁴ Width errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.
- ⁵ From a partial-wave analysis of data using a K-matrix formalism with 5 poles.
- ⁶ From an energy-independent partial-wave analysis.
- ⁷ From an amplitude analysis of the reaction $\pi^+\pi^-\rightarrow 2\pi^0$.
- ⁸ From an amplitude analysis of $\pi^+\pi^-\rightarrow \pi^+\pi^-$ scattering data.
- ⁹ JOHNSON 68 includes BONDAR 63, LEE 64, DERADO 65, EISNER 67.
- ¹⁰ T-matrix pole, 4 poles, 4 channels, including scattering data from HYAMS 75 ($\pi\pi$), LONGACRE 86 ($K\bar{K}$), BINON 83 ($\eta\eta$).
- ¹¹ 4-poles, 5-channel K matrix fit.
- ¹² From analysis of L3 data at 91 and 183–209 GeV.
- ¹³ Systematic uncertainties not estimated.

$f_2(1270)$ DECAY MODES

| Mode | Fraction (Γ_i/Γ) | Scale factor/ Confidence level |
|--|--------------------------------|-----------------------------------|
| Γ_1 $\pi\pi$ | $(84.2^{+2.9}_{-0.9})\%$ | S=1.1 |
| Γ_2 $\pi^+\pi^-2\pi^0$ | $(7.7^{+1.1}_{-3.2})\%$ | S=1.2 |
| Γ_3 $K\bar{K}$ | $(4.6^{+0.5}_{-0.4})\%$ | S=2.7 |
| Γ_4 $2\pi^+2\pi^-$ | $(2.8\pm 0.4)\%$ | S=1.2 |
| Γ_5 $\eta\eta$ | $(4.0\pm 0.8)\times 10^{-3}$ | S=2.1 |
| Γ_6 $4\pi^0$ | $(3.0\pm 1.0)\times 10^{-3}$ | |
| Γ_7 $\gamma\gamma$ | $(1.42\pm 0.24)\times 10^{-5}$ | S=1.4 |
| Γ_8 $\eta\pi\pi$ | $< 8 \times 10^{-3}$ | CL=95% |
| Γ_9 $K^0K^-\pi^+ + \text{c.c.}$ | $< 3.4 \times 10^{-3}$ | CL=95% |
| Γ_{10} e^+e^- | $< 6 \times 10^{-10}$ | CL=90% |

CONSTRAINED FIT INFORMATION

An overall fit to the total width, 4 partial widths, a combination of partial widths obtained from integrated cross sections, and 6 branching ratios uses 45 measurements and one constraint to determine 8 parameters. The overall fit has a $\chi^2 = 83.0$ for 38 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i/\Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

| | | | | | | | |
|----------|-------|-------|-------|-------|-------|-------|-------|
| x_2 | -90 | | | | | | |
| x_3 | 10 | -39 | | | | | |
| x_4 | 10 | -38 | 1 | | | | |
| x_5 | 1 | -6 | 0 | 0 | | | |
| x_6 | 0 | -7 | 0 | 0 | 0 | | |
| x_7 | 3 | 1 | -15 | 0 | 0 | 0 | |
| Γ | -71 | 65 | -10 | -7 | -1 | 0 | -6 |
| | x_1 | x_2 | x_3 | x_4 | x_5 | x_6 | x_7 |

| Mode | Rate (MeV) | Scale factor |
|-------------------------------|------------------------|--------------|
| Γ_1 $\pi\pi$ | 157.2 $^{+4.0}_{-1.1}$ | |
| Γ_2 $\pi^+\pi^-2\pi^0$ | 14.4 $^{+2.1}_{-6.0}$ | 1.2 |
| Γ_3 $K\bar{K}$ | 8.5 ± 0.8 | 2.8 |
| Γ_4 $2\pi^+2\pi^-$ | 5.2 ± 0.7 | 1.2 |
| Γ_5 $\eta\eta$ | 0.75 ± 0.14 | 2.1 |
| Γ_6 $4\pi^0$ | 0.56 ± 0.19 | |
| Γ_7 $\gamma\gamma$ | 0.0026 ± 0.0005 | 1.4 |

$f_2(1270)$ PARTIAL WIDTHS

$\Gamma(\pi\pi)$ Γ_1

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------------|------|-------------|------|---------|
|-------------|------|-------------|------|---------|

157.2 $^{+4.0}_{-1.1}$ OUR FIT

157.0 $^{+6.0}_{-1.0}$ ¹ LONGACRE 86 MPS 22 $\pi^-p \rightarrow n2K_S^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

152 ± 8 870 ² SCHEGELSKY 06A RVUE $\gamma\gamma \rightarrow K_S^0 K_S^0$

¹ From a partial-wave analysis of data using a K-matrix formalism with 5 poles.

² From analysis of L3 data at 91 and 183–209 GeV and using SU(3) relations.

$\Gamma(K\bar{K})$ Γ_3

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------------|------|-------------|------|---------|
|-------------|------|-------------|------|---------|

8.5 ± 0.8 OUR FIT Error includes scale factor of 2.8.

9.0 $^{+0.7}_{-0.3}$ ¹ LONGACRE 86 MPS 22 $\pi^-p \rightarrow n2K_S^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

7.5 ± 2.0 870 ² SCHEGELSKY 06A RVUE $\gamma\gamma \rightarrow K_S^0 K_S^0$

¹ From a partial-wave analysis of data using a K-matrix formalism with 5 poles.

² From analysis of L3 data at 91 and 183–209 GeV and using SU(3) relations.

$\Gamma(\eta\eta)$ Γ_5

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------------|------|-------------|------|---------|
|-------------|------|-------------|------|---------|

0.75±0.14 OUR FIT Error includes scale factor of 2.1.**1.0 ±0.1** ¹ LONGACRE 86 MPS 22 $\pi^- p \rightarrow n 2K_S^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.8 ±0.4 870 ² SCHEGELSKY 06A RVUE $\gamma\gamma \rightarrow K_S^0 K_S^0$ ¹ From a partial-wave analysis of data using a K-matrix formalism with 5 poles.² From analysis of L3 data at 91 and 183–209 GeV and using SU(3) relations. $\Gamma(\gamma\gamma)$ Γ_7

The value of this width depends on the theoretical model used. Unitary approaches with scalars typically (with exception of PENNINGTON 08) give values clustering around 2.6 keV; without an S-wave contribution, values are systematically higher (typically around 3 keV).

| VALUE (keV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------------|------|-------------|------|---------|
|-------------|------|-------------|------|---------|

2.6 ±0.5 OUR FIT Error includes scale factor of 1.4.**2.93±0.40** ¹ DAI 14A RVUE Compilation

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.14±0.20 ^{2,3} PENNINGTON 08 RVUE Compilation3.82±0.30 ^{3,4} PENNINGTON 08 RVUE Compilation2.55±0.15 870 ⁵ SCHEGELSKY 06A RVUE $\gamma\gamma \rightarrow K_S^0 K_S^0$ 2.84±0.35 BOGLIONE 99 RVUE $\gamma\gamma \rightarrow \pi^+ \pi^-, \pi^0 \pi^0$ 2.93±0.23±0.32 ⁶ YABUKI 95 VNS2.58±0.13 ^{+0.36}_{-0.27} ⁷ BEHREND 92 CELL $e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$ 3.10±0.35±0.35 ⁸ BLINOV 92 MD1 $e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$ 2.27±0.47±0.11 ADACHI 90D TOPZ $e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$ 3.15±0.04±0.39 BOYER 90 MRK2 $e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$ 3.19±0.16 ^{+0.29}_{-0.28} MARSISKE 90 CBAL $e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$ 2.35±0.65 ⁹ MORGAN 90 RVUE $\gamma\gamma \rightarrow \pi^+ \pi^-, \pi^0 \pi^0$ 3.19±0.09 ^{+0.22}_{-0.38} 2177 OEST 90 JADE $e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$ 3.2 ±0.1 ±0.4 ¹⁰ AIHARA 86B TPC $e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$ 2.5 ±0.1 ±0.5 BEHREND 84B CELL $e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$ 2.85±0.25±0.5 ¹¹ BERGER 84 PLUT $e^+ e^- \rightarrow e^+ e^- 2\pi$ 2.70±0.05±0.20 COURAU 84 DLCO $e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$ 2.52±0.13±0.38 ¹² SMITH 84C MRK2 $e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$ 2.7 ±0.2 ±0.6 EDWARDS 82F CBAL $e^+ e^- \rightarrow e^+ e^- 2\pi^0$ 2.9 ^{+0.6}_{-0.4} ±0.6 ¹³ EDWARDS 82F CBAL $e^+ e^- \rightarrow e^+ e^- 2\pi^0$ 3.2 ±0.2 ±0.6 BRANDELIK 81B TASS $e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$ 3.6 ±0.3 ±0.5 ROUSSARIE 81 MRK2 $e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$ 2.3 ±0.8 ¹⁴ BERGER 80B PLUT $e^+ e^-$ ¹ Based on a K-matrix analysis of BELLE data from MORI 07, UEHARA 08A, UEHARA 09 and UEHARA 13. The width is derived for the pole on the third sheet which is closest to the physical axis. Supersedes PENNINGTON 08.² Solution A (preferred solution based on χ^2 -analysis).³ Dispersion theory based amplitude analysis of BOYER 90, MARSISKE 90, BEHREND 92, and MORI 07.

- ⁴ Solution B (worse than solution A; still acceptable when systematic uncertainties are included).
⁵ From analysis of L3 data at 91 and 183–209 GeV and using SU(3) relations.
⁶ With a narrow scalar state around 1220 MeV.
⁷ Using a unitarized model with a 300 - 500 keV wide scalar at 1100 MeV.
⁸ Using the unitarized model of LYTH 85.
⁹ Error includes spread of different solutions. Data of MARK2 and CRYSTAL BALL used in the analysis. Authors report strong correlations with $\gamma\gamma$ width of $f_0(1370)$: $\Gamma(f_2) + 1/4 \Gamma(f^0) = 3.6 \pm 0.3$ KeV.
¹⁰ Radiative corrections modify the partial widths; for instance the COURAU 84 value becomes 2.66 ± 0.21 in the calculation of LANDRO 86.
¹¹ Using the MENNESSIER 83 model.
¹² Superseded by BOYER 90.
¹³ If helicity = 2 assumption is not made.
¹⁴ Using mass, width and $B(f_2(1270) \rightarrow 2\pi)$ from PDG 78.

| $\Gamma(e^+e^-)$ | | | | | Γ_{10} |
|---|-----|-------------|------|---------|---------------------------------|
| VALUE (eV) | CL% | DOCUMENT ID | TECN | COMMENT | |
| <0.11 | 90 | ACHASOV | 00K | SND | $e^+e^- \rightarrow \pi^0\pi^0$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| <1.7 | 90 | VOROBYEV | 88 | ND | $e^+e^- \rightarrow \pi^0\pi^0$ |

$f_2(1270) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

| $\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ | | | | | $\Gamma_3\Gamma_7/\Gamma$ |
|---|-------------------------------------|-----------------------|------|---------|-----------------------------------|
| VALUE (keV) | | DOCUMENT ID | TECN | COMMENT | |
| 0.121±0.020 OUR FIT | Error includes scale factor of 1.3. | | | | |
| 0.091±0.007±0.027 | | ¹ ALBRECHT | 90G | ARG | $e^+e^- \rightarrow e^+e^-K^+K^-$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| 0.104±0.007±0.072 | | ² ALBRECHT | 90G | ARG | $e^+e^- \rightarrow e^+e^-K^+K^-$ |
| ¹ Using an incoherent background. | | | | | |
| ² Using a coherent background. | | | | | |

| $\Gamma(\eta\eta) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ | | | | | $\Gamma_5\Gamma_7/\Gamma$ |
|---|--|---------------------|------|---------|--|
| VALUE (eV) | | DOCUMENT ID | TECN | COMMENT | |
| 11.5^{+1.8+4.5}_{-2.0-3.7} | | ¹ UEHARA | 10A | BELL | 10.6 $e^+e^- \rightarrow e^+e^-\eta\eta$ |
| ¹ Including interference with the $f_2'(1525)$ (parameters fixed to the values from the 2008 edition of this review, PDG 08) and $f_0(\Upsilon)$. | | | | | |

Helicity-0/Helicity-2 RATIO IN $\gamma\gamma \rightarrow f_2(1270) \rightarrow \pi\pi$

| VALUE (units 10^{-2}) | | DOCUMENT ID | TECN | COMMENT | |
|---|--|---------------------------|------|---------|--|
| 3.7±0.3^{+15.9}_{-2.9} | | UEHARA | 08A | BELL | 10.6 $e^+e^- \rightarrow e^+e^-\pi^0\pi^0$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| 9.5±1.8 | | ¹ DAI | 14A | RVUE | Compilation |
| 13 | | ^{2,3} PENNINGTON | 08 | RVUE | Compilation |
| 26 | | ^{3,4} PENNINGTON | 08 | RVUE | Compilation |

¹ Based on a K -matrix analysis of BELLE data from MORI 07, UEHARA 08A, UEHARA 09 and UEHARA 13. The width is derived for the pole on the third sheet which is closest to the physical axis.

² Solution A (preferred solution based on χ^2 -analysis).

³ Dispersion theory based amplitude analysis of BOYER 90, MARSISKE 90, BEHREND 92, and MORI 07.

⁴ Solution B (worse than solution A; still acceptable when systematic uncertainties are included).

$f_2(1270)$ BRANCHING RATIOS

$\Gamma(\pi\pi)/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE EVTS DOCUMENT ID TECN COMMENT

0.842^{+0.029}_{-0.009} **OUR FIT** Error includes scale factor of 1.1.

0.837 \pm 0.020 **OUR AVERAGE**

| | | | | | |
|-------------------|-----|---------|----|------|---|
| 0.849 \pm 0.025 | | CHABAUD | 83 | ASPK | 17 $\pi^- p$ polarized |
| 0.85 \pm 0.05 | 250 | BEAUPRE | 71 | HBC | 8 $\pi^+ p \rightarrow \Delta^{++} f_2^-$ |
| 0.8 \pm 0.04 | 600 | OH | 70 | HBC | 1.26 $\pi^- p \rightarrow \pi^+ \pi^- n$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|------------------------------|--|-----------------------|----|------|---|
| 0.856 \pm 0.001 \pm 0.05 | | ¹ ALBRECHT | 20 | RVUE | 0.9 $\bar{p} p \rightarrow \pi^0 \pi^0 \eta,$ $\pi^0 \eta \eta, \pi^0 K^+ K^-$ |
|------------------------------|--|-----------------------|----|------|---|

¹ Residue from T-matrix pole, 4 poles, 4 channels, including scattering data from HYAMS 75 ($\pi\pi$), LONGACRE 86 ($K\bar{K}$), BINON 83 ($\eta\eta$).

$\Gamma(\pi^+ \pi^- 2\pi^0)/\Gamma(\pi\pi)$ Γ_2/Γ_1

Should be twice $\Gamma(2\pi^+ 2\pi^-)/\Gamma(\pi\pi)$ if decay is $\rho\rho$. (See ASCOLI 68D.)

VALUE EVTS DOCUMENT ID TECN COMMENT

0.091^{+0.014}_{-0.040} **OUR FIT** Error includes scale factor of 1.2.

| | | | | | |
|----------------------------------|-----|-----------|----|-----|---|
| 0.15 \pm0.06 | 600 | EISENBERG | 74 | HBC | 4.9 $\pi^+ p \rightarrow \Delta^{++} f_2^-$ |
|----------------------------------|-----|-----------|----|-----|---|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|------|--|------|-----|-----|---------------------------------|
| 0.07 | | EMMS | 75D | DBC | 4 $\pi^+ n \rightarrow p f_2^-$ |
|------|--|------|-----|-----|---------------------------------|

$\Gamma(K\bar{K})/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|-------------------------------|--|-----------------------|----|------|---|
| 0.033 \pm 0.001 \pm 0.005 | | ¹ ALBRECHT | 20 | RVUE | 0.9 $\bar{p} p \rightarrow \pi^0 \pi^0 \eta,$ $\pi^0 \eta \eta, \pi^0 K^+ K^-$ |
|-------------------------------|--|-----------------------|----|------|---|

¹ Residue from T-matrix pole, 4 poles, 4 channels, including scattering data from HYAMS 75 ($\pi\pi$), LONGACRE 86 ($K\bar{K}$), BINON 83 ($\eta\eta$).

$\Gamma(K\bar{K})/\Gamma(\pi\pi)$

Γ_3/Γ_1

We average only experiments which either take into account $f_2(1270)$ - $a_2(1320)$ interference explicitly or demonstrate that $a_2(1320)$ production is negligible.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------|------|-------------|------|---------|
|-------|------|-------------|------|---------|

0.054^{+0.005}_{-0.006} OUR FIT Error includes scale factor of 2.7.

0.041^{+0.004}_{-0.005} OUR AVERAGE

| | | | | | |
|---|----|--------------------------|-----|------|---|
| 0.045±0.01 | | ¹ BARGIOTTI | 03 | OBLX | $\bar{p}p$ |
| 0.037 ^{+0.008} _{-0.021} | | ETKIN | 82B | MPS | 23 $\pi^- p \rightarrow n 2K_S^0$ |
| 0.045±0.009 | | CHABAUD | 81 | ASPK | 17 $\pi^- p$ polarized |
| 0.039±0.008 | | LOVERRE | 80 | HBC | 4 $\pi^- p \rightarrow K\bar{K}N$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | | |
| 0.052±0.025 | | ABLIKIM | 04E | BES2 | $J/\psi \rightarrow \omega K^+ K^-$ |
| 0.036±0.005 | | ² COSTA | 80 | OMEG | 1-2.2 $\pi^- p \rightarrow K^+ K^- n$ |
| 0.030±0.005 | | ³ MARTIN | 79 | RVUE | |
| 0.027±0.009 | | ⁴ POLYCHRO... | 79 | STRC | 7 $\pi^- p \rightarrow n 2K_S^0$ |
| 0.025±0.015 | | EMMS | 75D | DBC | 4 $\pi^+ n \rightarrow p f_2$ |
| 0.031±0.012 | 20 | ADERHOLZ | 69 | HBC | 8 $\pi^+ p \rightarrow K^+ K^- \pi^+ p$ |

¹ Coupled channel analysis of $\pi^+ \pi^- \pi^0$, $K^+ K^- \pi^0$, and $K^\pm K_S^0 \pi^\mp$.

² Re-evaluated by CHABAUD 83.

³ Includes PAWLICKI 77 data.

⁴ Takes into account the $f_2(1270)$ - $f'_2(1525)$ interference.

$\Gamma(2\pi^+ 2\pi^-)/\Gamma(\pi\pi)$

Γ_4/Γ_1

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------|------|-------------|------|---------|
|-------|------|-------------|------|---------|

0.033±0.005 OUR FIT Error includes scale factor of 1.2.

0.033±0.004 OUR AVERAGE Error includes scale factor of 1.1.

| | | | | | |
|---|-----|--------------------|-----|-----|---|
| 0.024±0.006 | 160 | EMMS | 75D | DBC | 4 $\pi^+ n \rightarrow p f_2$ |
| 0.051±0.025 | 70 | EISENBERG | 74 | HBC | 4.9 $\pi^+ p \rightarrow \Delta^{++} f_2$ |
| 0.043 ^{+0.007} _{-0.011} | 285 | ¹ LOUIE | 74 | HBC | 3.9 $\pi^- p \rightarrow n f_2$ |
| 0.037±0.007 | 154 | ANDERSON | 73 | DBC | 6 $\pi^+ n \rightarrow p f_2$ |
| 0.047±0.013 | | OH | 70 | HBC | 1.26 $\pi^- p \rightarrow \pi^+ \pi^- n$ |

¹ LOUIE 74 was quoted as 0.065 in PDG 74. Factor 2/3 to go from $\pi^+ \pi^- \rightarrow \pi\pi$ forgotten. Mike L.

$\Gamma(\eta\eta)/\Gamma_{\text{total}}$

Γ_5/Γ

| VALUE (units 10 ⁻³) | DOCUMENT ID | TECN | COMMENT |
|---------------------------------|-------------|------|---------|
|---------------------------------|-------------|------|---------|

4.0±0.8 OUR FIT Error includes scale factor of 2.1.

2.9±0.5 OUR AVERAGE

| | | | | |
|---------|-------|-----|------|-------------------------------------|
| 2.7±0.7 | BINON | 05 | GAMS | 33 $\pi^- p \rightarrow \eta\eta n$ |
| 2.8±0.7 | ALDE | 86D | GAM4 | 100 $\pi^- p \rightarrow 2\eta n$ |
| 5.2±1.7 | BINON | 83 | GAM2 | 38 $\pi^- p \rightarrow 2\eta n$ |

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

4.0±1.0±2.0 ¹ ALBRECHT 20 RVUE 0.9 $\bar{p}p \rightarrow \pi^0 \pi^0 \eta, \pi^0 \eta\eta, \pi^0 K^+ K^-$

¹ Residue from T-matrix pole, 4 poles, 4 channels, including scattering data from HYAMS 75 ($\pi\pi$), LONGACRE 86 ($K\bar{K}$), BINON 83 ($\eta\eta$).

$\Gamma(\eta\eta)/\Gamma(\pi\pi)$ Γ_5/Γ_1

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------|------|---|
| 0.003±0.001 | | BARBERIS | 00E | 450 $p p \rightarrow p_f \eta \eta p_S$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| <0.05 | 95 | EDWARDS | 82F | CBAL $e^+ e^- \rightarrow e^+ e^- 2\eta$ |
| <0.016 | 95 | EMMS | 75D | DBC $4 \pi^+ n \rightarrow p f_2$ |
| <0.09 | 95 | EISENBERG | 74 | HBC $4.9 \pi^+ p \rightarrow \Delta^{++} f_2$ |

$\Gamma(4\pi^0)/\Gamma_{total}$ Γ_6/Γ

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------------|----------|-------------|------|---|
| 0.0030±0.0010 OUR FIT | | | | |
| 0.003 ±0.001 | 400 ± 50 | ALDE | 87 | GAM4 $100 \pi^- p \rightarrow 4\pi^0 n$ |

$\Gamma(\gamma\gamma)/\Gamma_{total}$ Γ_7/Γ

| VALUE (units 10^{-5}) | DOCUMENT ID | TECN | COMMENT |
|---|-------------|------|---|
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| $1.57 \pm 0.01^{+1.39}_{-0.14}$ | UEHARA | 08A | BELL $10.6 e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$ |

$\Gamma(\eta\pi\pi)/\Gamma(\pi\pi)$ Γ_8/Γ_1

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|------------------|-----|-------------|------|-----------------------------------|
| <0.010 | 95 | EMMS | 75D | DBC $4 \pi^+ n \rightarrow p f_2$ |

$\Gamma(K^0 K^- \pi^+ + c.c.)/\Gamma(\pi\pi)$ Γ_9/Γ_1

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|------------------|-----|-------------|------|-----------------------------------|
| <0.004 | 95 | EMMS | 75D | DBC $4 \pi^+ n \rightarrow p f_2$ |

$\Gamma(e^+ e^-)/\Gamma_{total}$ Γ_{10}/Γ

| VALUE (units 10^{-10}) | CL% | DOCUMENT ID | TECN | COMMENT |
|---------------------------|-----|-------------|------|---------------------------------------|
| <6 | 90 | ACHASOV | 00K | SND $e^+ e^- \rightarrow \pi^0 \pi^0$ |

$f_2(1270)$ REFERENCES

| | | | | |
|-------------|-----|------------------------------|---------------------------------|--------------------------|
| ALBRECHT | 20 | EPJ C80 453 | M. Albrecht <i>et al.</i> | (Crystal Barrel Collab.) |
| DOBBS | 15 | PR D91 052006 | S. Dobbs <i>et al.</i> | (NWES) |
| DAI | 14A | PR D90 036004 | L.-Y. Dai, M.R. Pennington | (CEBAF) |
| BOGOLYUB... | 13 | PAN 76 1324 | M.Yu. Bogolyubsky <i>et al.</i> | (HYPERON-M Collab.) |
| | | Translated from YAF 76 1389. | | |
| UEHARA | 13 | PTEP 2013 123C01 | S. Uehara <i>et al.</i> | (BELLE Collab.) |
| UEHARA | 10A | PR D82 114031 | S. Uehara <i>et al.</i> | (BELLE Collab.) |
| ANISOVICH | 09 | IJMP A24 2481 | V.V. Anisovich, A.V. Sarantsev | |
| UEHARA | 09 | PR D79 052009 | S. Uehara <i>et al.</i> | (BELLE Collab.) |
| PDG | 08 | PL B667 1 | C. Amsler <i>et al.</i> | (PDG Collab.) |
| PENNINGTON | 08 | EPJ C56 1 | M.R. Pennington <i>et al.</i> | |
| UEHARA | 08A | PR D78 052004 | S. Uehara <i>et al.</i> | (BELLE Collab.) |
| MORI | 07 | PR D75 051101 | T. Mori <i>et al.</i> | (BELLE Collab.) |
| ABLIKIM | 06V | PL B642 441 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| SCHEGELSKY | 06A | EPJ A27 207 | V.A. Schegelsky <i>et al.</i> | |
| ABLIKIM | 05 | PL B607 243 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| BINON | 05 | PAN 68 960 | F. Binon <i>et al.</i> | |
| | | Translated from YAF 68 998. | | |
| ABLIKIM | 04E | PL B603 138 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| BARGIOTTI | 03 | EPJ C26 371 | M. Bargiotti <i>et al.</i> | (OBELIX Collab.) |
| TIKHOMIROV | 03 | PAN 66 828 | G.D. Tikhomirov <i>et al.</i> | |
| | | Translated from YAF 66 860. | | |

| | | | | |
|-------------|-----|--------------------------|--|-----------------------------|
| ACHASOV | 00K | PL B492 8 | M.N. Achasov <i>et al.</i> | (Novosibirsk SND Collab.) |
| BARBERIS | 00E | PL B479 59 | D. Barberis <i>et al.</i> | (WA 102 Collab.) |
| BOGLIONE | 99 | EPJ C9 11 | M. Boggione, M.R. Pennington | |
| ALDE | 98 | EPJ A3 361 | D. Alde <i>et al.</i> | (GAM4 Collab.) |
| Also | | PAN 62 405 | D. Alde <i>et al.</i> | (GAMS Collab.) |
| | | Translated from YAF 62 | 446. | |
| ALDE | 97 | PL B397 350 | D.M. Alde <i>et al.</i> | (GAMS Collab.) |
| BERTIN | 97C | PL B408 476 | A. Bertin <i>et al.</i> | (OBELIX Collab.) |
| GRYGOREV | 96 | PAN 59 2105 | V.K. Grigoriev, O.N. Baloshin, B.P. Barkov | (ITEP) |
| | | Translated from YAF 59 | 2187. | |
| YABUKI | 95 | JPSJ 64 435 | F. Yabuki <i>et al.</i> | (VENUS Collab.) |
| PROKOSHKIN | 94 | PD 39 420 | Y.D. Prokoshkin, A.A. Kondashov | (SERP) |
| | | Translated from DANS 336 | 613. | |
| BEHREND | 92 | ZPHY C56 381 | H.J. Behrend | (CELLO Collab.) |
| BLINOV | 92 | ZPHY C53 33 | A.E. Blinov <i>et al.</i> | (NOVO) |
| AGUILAR-... | 91 | ZPHY C50 405 | M. Aguilar-Benitez <i>et al.</i> | (LEBC-EHS Collab.) |
| AKER | 91 | PL B260 249 | E. Aker <i>et al.</i> | (Crystal Barrel Collab.) |
| ADACHI | 90D | PL B234 185 | I. Adachi <i>et al.</i> | (TOPAZ Collab.) |
| ALBRECHT | 90G | ZPHY C48 183 | H. Albrecht <i>et al.</i> | (ARGUS Collab.) |
| BOYER | 90 | PR D42 1350 | J. Boyer <i>et al.</i> | (Mark II Collab.) |
| BREAKSTONE | 90 | ZPHY C48 569 | A.M. Breakstone <i>et al.</i> | (ISU, BGNA, CERN+) |
| MARSISKE | 90 | PR D41 3324 | H. Marsiske <i>et al.</i> | (Crystal Ball Collab.) |
| MORGAN | 90 | ZPHY C48 623 | D. Morgan, M.R. Pennington | (RAL, DURH) |
| OEST | 90 | ZPHY C47 343 | T. Oest <i>et al.</i> | (JADE Collab.) |
| AUGUSTIN | 89 | NP B320 1 | J.E. Augustin, G. Cosme | (DM2 Collab.) |
| VOROBYEV | 88 | SJNP 48 273 | P.V. Vorobiev <i>et al.</i> | (NOVO) |
| | | Translated from YAF 48 | 436. | |
| ALDE | 87 | PL B198 286 | D.M. Alde <i>et al.</i> | (LANL, BRUX, SERP, LAPP) |
| AUGUSTIN | 87 | ZPHY C36 369 | J.E. Augustin <i>et al.</i> | (LALO, CLER, FRAS+) |
| ABACHI | 86B | PRL 57 1990 | S. Abachi <i>et al.</i> | (PURD, ANL, IND, MICH+) |
| AIHARA | 86B | PRL 57 404 | H. Aihara <i>et al.</i> | (TPC-2 γ Collab.) |
| ALDE | 86D | NP B269 485 | D.M. Alde <i>et al.</i> | (BELG, LAPP, SERP, CERN+) |
| LANDRO | 86 | PL B172 445 | M. Landro, K.J. Mork, H.A. Olsen | (UTRO) |
| LONGACRE | 86 | PL B177 223 | R.S. Longacre <i>et al.</i> | (BNL, BRAN, CUNY+) |
| LYTH | 85 | JP G11 459 | D.H. Lyth | |
| BEHREND | 84B | ZPHY C23 223 | H.J. Behrend <i>et al.</i> | (CELLO Collab.) |
| BERGER | 84 | ZPHY C26 199 | C. Berger <i>et al.</i> | (PLUTO Collab.) |
| COURAU | 84 | PL 147B 227 | A. Courau <i>et al.</i> | (CIT, SLAC) |
| SMITH | 84C | PR D30 851 | J.R. Smith <i>et al.</i> | (SLAC, LBL, HARV) |
| BINON | 83 | NC 78A 313 | F.G. Binon <i>et al.</i> | (BELG, LAPP, SERP+) |
| Also | | SJNP 38 561 | F.G. Binon <i>et al.</i> | (BELG, LAPP, SERP+) |
| | | Translated from YAF 38 | 934. | |
| CHABAUD | 83 | NP B223 1 | V. Chabaud <i>et al.</i> | (CERN, CRAC, MPIM) |
| DENNEY | 83 | PR D28 2726 | D.L. Denney <i>et al.</i> | (IOWA, MICH) |
| MENNESSIER | 83 | ZPHY C16 241 | G. Mennessier | (MONP) |
| APEL | 82 | NP B201 197 | W.D. Apel <i>et al.</i> | (KARLK, KARLE, PISA, SERP+) |
| CASON | 82 | PRL 48 1316 | N.M. Cason <i>et al.</i> | (NDAM, ANL) |
| EDWARDS | 82F | PL 110B 82 | C. Edwards <i>et al.</i> | (CIT, HARV, PRIN+) |
| ETKIN | 82B | PR D25 1786 | A. Etkin <i>et al.</i> | (BNL, CUNY, TUFTS, VAND) |
| BRANDELIK | 81B | ZPHY C10 117 | R. Brandelik <i>et al.</i> | (TASSO Collab.) |
| CHABAUD | 81 | APP B12 575 | V. Chabaud <i>et al.</i> | (CERN, CRAC, MPIM) |
| GIDAL | 81 | PL 107B 153 | G. Gidal <i>et al.</i> | (SLAC, LBL) |
| ROUSSARIE | 81 | PL 105B 304 | A. Roussarie <i>et al.</i> | (SLAC, LBL) |
| BERGER | 80B | PL 94B 254 | C. Berger <i>et al.</i> | (PLUTO Collab.) |
| COSTA | 80 | NP B175 402 | G. Costa <i>et al.</i> | (BARI, BONN, CERN, GLAS+) |
| LOVERRE | 80 | ZPHY C6 187 | P.F. Loverre <i>et al.</i> | (CERN, CDEF, MADR+) |
| CORDEN | 79 | NP B157 250 | M.J. Corden <i>et al.</i> | (BIRM, RHEL, TELA+) |
| MARTIN | 79 | NP B158 520 | A.D. Martin, E.N. Ozmutlu | (DURH) |
| POLYCHRO... | 79 | PR D19 1317 | V.A. Polychronakos <i>et al.</i> | (NDAM, ANL) |
| PDG | 78 | PL 75B 1 | C. Bricman <i>et al.</i> | |
| ANTIPOV | 77 | NP B119 45 | Y.M. Antipov <i>et al.</i> | (SERP, GEVA) |
| PAWLICKI | 77 | PR D15 3196 | A.J. Pawlicki <i>et al.</i> | (ANL) |
| DEUTSCH... | 76 | NP B103 426 | M. Deuschmann <i>et al.</i> | (AACH3, BERL, BONN+) |
| APEL | 75 | PL 57B 398 | W.D. Apel <i>et al.</i> | (KARLK, KARLE, PISA, SERP+) |
| EMMS | 75D | NP B96 155 | M.J. Emms <i>et al.</i> | (BIRM, DURH, RHEL) |
| HYAMS | 75 | NP B100 205 | B.D. Hyams <i>et al.</i> | (CERN, MPIM) |
| EISENBERG | 74 | PL 52B 239 | Y. Eisenberg <i>et al.</i> | (REHO) |
| ENGLER | 74 | PR D10 2070 | A. Engler <i>et al.</i> | (CMU, CASE) |
| LOUIE | 74 | PL 48B 385 | J. Louie <i>et al.</i> | (SACL, CERN) |
| PDG | 74 | PL 50B 1 | V. Chaloupka <i>et al.</i> | |
| ANDERSON | 73 | PRL 31 562 | J.C. Anderson <i>et al.</i> | (CMU, CASE) |

| | | | | |
|------------|-----|-------------|-------------------------------|---------------------------|
| TAKAHASHI | 72 | PR D6 1266 | K. Takahashi <i>et al.</i> | (TOHOK, PENN, NDAM+) |
| BEAUPRE | 71 | NP B28 77 | J.V. Beaupre <i>et al.</i> | (AACH, BERL, CERN) |
| FLATTE | 71 | PL 34B 551 | S.M. Flatte <i>et al.</i> | (LBL) |
| ARMENISE | 70 | LNC 4 199 | N. Armenise <i>et al.</i> | (BARI, BGNA, FIRZ) |
| OH | 70 | PR D1 2494 | B.Y. Oh <i>et al.</i> | (WISC, TNTO) JP |
| STUNTEBECK | 70 | PL 32B 391 | P.H. Stuntebeck <i>et al.</i> | (NDAM) |
| ADERHOLZ | 69 | NP B11 259 | M. Aderholz <i>et al.</i> | (AACH3, BERL, CERN+) |
| ARMENISE | 68 | NC 54A 999 | N. Armenise <i>et al.</i> | (BARI, BGNA, FIRZ+) |
| ASCOLI | 68D | PRL 21 1712 | G. Ascoli <i>et al.</i> | (ILL) |
| BOESEBECK | 68 | NP B4 501 | K. Boesebeck <i>et al.</i> | (AACH, BERL, CERN) |
| JOHNSON | 68 | PR 176 1651 | P.B. Johnson <i>et al.</i> | (NDAM, PURD, SLAC) |
| EISNER | 67 | PR 164 1699 | R.L. Eisner <i>et al.</i> | (PURD) |
| DERADO | 65 | PRL 14 872 | I. Derado <i>et al.</i> | (NDAM) |
| LEE | 64 | PRL 12 342 | Y.Y. Lee <i>et al.</i> | (MICH) |
| BONDAR | 63 | PL 5 153 | L. Bondar <i>et al.</i> | (AACH, BIRM, BONN, DESY+) |
