

$\psi(4260)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

OMITTED FROM SUMMARY TABLE

also known as $Y(4260)$; was $X(4260)$

The state $\psi(4260)$ received its mass label from a Breit-Wigner (BW) fit to the $J/\psi\pi\pi$ data listed below. The symmetric BW placed the mass unavoidably into the center of the distribution. The most recent measurement in the 4260 MeV mass range in the same channel (ABLIKIM 17B), however, revealed that the distribution is asymmetric and that the state has a much lower mass consistent with the entry for particle $\psi(4230)$. Thus, in this edition we merged the measurement of ABLIKIM 17B with the $\psi(4230)$ node and labeled the older measurements of this node as not used. For details see the review on "Spectroscopy of mesons containing two heavy quarks."

 $\psi(4260)$ MASS

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$4209.1 \pm 6.8 \pm 7.0$		¹ ZHANG	17B RVUE	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$
$4223.3 \pm 1.6 \pm 2.5$		² ZHANG	17C RVUE	$e^+e^- \rightarrow \pi^+\pi^-J/\psi$ or $\psi(2S)$
$4258.6 \pm 8.3 \pm 12.1$		³ LIU	13B BELL	$e^+e^- \rightarrow \gamma\pi^+\pi^-J/\psi$
$4245 \pm 5 \pm 4$		⁴ LEES	12AC BABR	$10.58 e^+e^- \rightarrow \gamma\pi^+\pi^-J/\psi$
$4247 \pm 12 \pm^{+17}_{-32}$		^{3,5} YUAN	07 BELL	$10.58 e^+e^- \rightarrow \gamma\pi^+\pi^-J/\psi$
$4284 \pm^{+17}_{-16} \pm 413.6$		HE	06B CLEO	$9.4\text{--}10.6 e^+e^- \rightarrow \gamma\pi^+\pi^-J/\psi$
$4259 \pm 8 \pm^{+2}_{-6} 125$		⁶ AUBERT,B	05I BABR	$10.58 e^+e^- \rightarrow \gamma\pi^+\pi^-J/\psi$

¹ From a three-resonance fit.² From a combined fit of BELLE, BABAR and BES3 $e^+e^- \rightarrow \pi^+\pi^-J/\psi$ and $e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$ data.³ From a two-resonance fit.⁴ From a single-resonance fit. Supersedes AUBERT,B 05I.⁵ Superseded by LIU 13B.⁶ From a single-resonance fit. Two interfering resonances are not excluded. Superseded by LEES 12AC. **$\psi(4260)$ WIDTH**

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$76.6 \pm 14.2 \pm 2.4$		¹ ZHANG	17B RVUE	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$
$54.2 \pm 2.6 \pm 1.0$		² ZHANG	17C RVUE	$e^+e^- \rightarrow \pi^+\pi^-J/\psi$ or $\psi(2S)$
$134.1 \pm 16.4 \pm 5.5$		³ LIU	13B BELL	$e^+e^- \rightarrow \gamma\pi^+\pi^-J/\psi$
$114 \pm^{+16}_{-15} \pm 7$		⁴ LEES	12AC BABR	$10.58 e^+e^- \rightarrow \gamma\pi^+\pi^-J/\psi$
$108 \pm 19 \pm 10$		^{3,5} YUAN	07 BELL	$10.58 e^+e^- \rightarrow \gamma\pi^+\pi^-J/\psi$

73	$\begin{matrix} +39 \\ -25 \end{matrix}$	± 5	13.6	HE	06B	CLEO	9.4–10.6	$e^+e^- \rightarrow \gamma\pi^+\pi^- J/\psi$
88	± 23	$\begin{matrix} +6 \\ -4 \end{matrix}$	125	⁶ AUBERT,B	05I	BABR	10.58	$e^+e^- \rightarrow \gamma\pi^+\pi^- J/\psi$

¹From a three-resonance fit.

²From a combined fit of BELLE, BABAR and BES3 $e^+e^- \rightarrow \pi^+\pi^- J/\psi$ and $e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$ data.

³From a two-resonance fit.

⁴From a single-resonance fit. Supersedes AUBERT,B 05I.

⁵Superseded by LIU 13B.

⁶From a single-resonance fit. Two interfering resonances are not excluded. Superseded by LEES 12AC.

$\psi(4260)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 e^+e^-	
Γ_2 $J/\psi\pi^+\pi^-$	seen
Γ_3 $J/\psi f_0(980), f_0(980) \rightarrow \pi^+\pi^-$	seen
Γ_4 $Z_c(3900)^\pm \pi^\mp, Z_c^\pm \rightarrow J/\psi\pi^\pm$	seen
Γ_5 $J/\psi\pi^0\pi^0$	seen
Γ_6 $J/\psi K^+K^-$	seen
Γ_7 $J/\psi K_S^0 K_S^0$	not seen
Γ_8 $J/\psi\eta$	not seen
Γ_9 $J/\psi\pi^0$	not seen
Γ_{10} $J/\psi\eta'$	not seen
Γ_{11} $J/\psi\pi^+\pi^-\pi^0$	not seen
Γ_{12} $J/\psi\eta\pi^0$	not seen
Γ_{13} $J/\psi\eta\eta$	not seen
Γ_{14} $\psi(2S)\pi^+\pi^-$	not seen
Γ_{15} $\psi(2S)\eta$	not seen
Γ_{16} $\chi_{c0}\omega$	not seen
Γ_{17} $\chi_{c1}\pi^+\pi^-\pi^0$	not seen
Γ_{18} $\chi_{c2}\pi^+\pi^-\pi^0$	not seen
Γ_{19} $h_c(1P)\pi^+\pi^-$	not seen
Γ_{20} $\phi\pi^+\pi^-$	not seen
Γ_{21} $\phi f_0(980) \rightarrow \phi\pi^+\pi^-$	not seen
Γ_{22} $D\bar{D}$	not seen
Γ_{23} $D^0\bar{D}^0$	not seen
Γ_{24} D^+D^-	not seen
Γ_{25} $D^*\bar{D} + \text{c.c.}$	not seen
Γ_{26} $D^*(2007)^0\bar{D}^0 + \text{c.c.}$	not seen
Γ_{27} $D^*(2010)^+D^- + \text{c.c.}$	not seen
Γ_{28} $D^*\bar{D}^*$	not seen
Γ_{29} $D^*(2007)^0\bar{D}^*(2007)^0$	not seen
Γ_{30} $D^*(2010)^+D^*(2010)^-$	not seen

Γ_{31}	$D\bar{D}\pi + \text{c.c.}$	
Γ_{32}	$D^0 D^- \pi^+ + \text{c.c.}$ (excl. $D^*(2007)^0 \bar{D}^{*0} + \text{c.c.},$ $D^*(2010)^+ D^- + \text{c.c.}$)	not seen
Γ_{33}	$D\bar{D}^* \pi + \text{c.c.}$ (excl. $D^* \bar{D}^*$)	not seen
Γ_{34}	$D^0 D^{*-} \pi^+ + \text{c.c.}$ (excl. $D^*(2010)^+ D^*(2010)^-$)	not seen
Γ_{35}	$D^0 D^*(2010)^- \pi^+ + \text{c.c.}$	not seen
Γ_{36}	$D_1(2420) \bar{D} + \text{c.c.}$	not seen
Γ_{37}	$D^* \bar{D}^* \pi$	not seen
Γ_{38}	$D_s^+ D_s^-$	not seen
Γ_{39}	$D_s^{*+} D_s^- + \text{c.c.}$	not seen
Γ_{40}	$D_s^{*+} D_s^{*-}$	not seen
Γ_{41}	$\rho \bar{\rho}$	not seen
Γ_{42}	$\rho \bar{\rho} \pi^0$	not seen
Γ_{43}	$\Xi^- \bar{\Xi}^+$	
Γ_{44}	$K_S^0 K^\pm \pi^\mp$	not seen
Γ_{45}	$K_S^0 K^\pm \pi^\mp \pi^0$	
Γ_{46}	$K_S^0 K^\pm \pi^\mp \eta$	
Γ_{47}	$K^+ K^- \pi^0$	not seen

Radiative decays

Γ_{48}	$\eta_c(1S) \gamma$	possibly seen
Γ_{49}	$\chi_{c1} \gamma$	not seen
Γ_{50}	$\chi_{c2} \gamma$	not seen
Γ_{51}	$\chi_{c1}(3872) \gamma$	seen

$\psi(4260) \Gamma(i) \times \Gamma(e^+ e^-) / \Gamma(\text{total})$

$\Gamma(J/\psi \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$					$\Gamma_2 \Gamma_1 / \Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
9.2 ± 1.0 OUR AVERAGE					
9.2 ± 0.8 ± 0.7		¹ LEES	12AC BABR	10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- J/\psi$	
8.9 ^{+3.9} _{-3.1} ± 1.8	8.1	HE	06B CLEO	9.4–10.6 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- J/\psi$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
6.4 ± 0.8 ± 0.6		² LIU	13B BELL	$e^+ e^- \rightarrow \gamma \pi^+ \pi^- J/\psi$	
20.5 ± 1.4 ± 2.0		³ LIU	13B BELL	$e^+ e^- \rightarrow \gamma \pi^+ \pi^- J/\psi$	
6.0 ± 1.2 ^{+4.7} _{-0.5}		^{2,4} YUAN	07 BELL	10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- J/\psi$	
20.6 ± 2.3 ^{+9.1} _{-1.7}		^{3,4} YUAN	07 BELL	10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- J/\psi$	
5.5 ± 1.0 ^{+0.8} _{-0.7}	125	⁵ AUBERT,B	05I BABR	10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- J/\psi$	

¹ From a single-resonance fit. Supersedes AUBERT,B 05I.

² Solution I of two equivalent solutions in a fit using two interfering resonances.

³ Solution II of two equivalent solutions in a fit using two interfering resonances.

⁴ Superseded by LIU 13B.

⁵ From a single-resonance fit. Two interfering resonances are not excluded. Superseded by LEES 12AC.

$\Gamma(J/\psi K^+ K^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_6 \Gamma_1/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<1.7	90	¹ SHEN	14	BELL 9.4–10.9 $e^+ e^- \rightarrow \gamma K^+ K^- J/\psi$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<1.2	90	² YUAN	08	BELL $e^+ e^- \rightarrow \gamma K^+ K^- J/\psi$

¹ From a fit of the broad $K^+ K^- J/\psi$ enhancement including a coherent $\psi(4260)$ amplitude with mass and width from LIU 13B. Supersedes YUAN 08. The shape of the cross section observed by ABLIKIM 18N between 2.2 and 2.3 GeV is incompatible with that of $e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$ in ABLIKIM 13T and ABLIKIM 17B. They also observe a broad enhancement around 2.5 GeV.

² From a fit of the broad $K^+ K^- J/\psi$ enhancement including a coherent $\psi(4260)$ amplitude with mass and width from YUAN 07.

$\Gamma(J/\psi K_S^0 K_S^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_7 \Gamma_1/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.85	90	¹ SHEN	14	BELL 9.4–10.9 $e^+ e^- \rightarrow \gamma K_S^0 K_S^0 J/\psi$

¹ From a fit of the $K_S^0 K_S^0 J/\psi$ mass range from 4.4 to 5.5 GeV including a coherent $\psi(4260)$ amplitude with mass and width from LIU 13B.

$\Gamma(J/\psi \eta) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_8 \Gamma_1/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<14.2	90	WANG	13B	BELL $e^+ e^- \rightarrow J/\psi \eta \gamma$

$\Gamma(J/\psi \eta') \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_{10} \Gamma_1/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.06 ± 0.03	46	^{1,2} ABLIKIM	20A	BES3 $e^+ e^- \rightarrow \eta' J/\psi$
1.38 ± 0.11	46	^{1,3} ABLIKIM	20A	BES3 $e^+ e^- \rightarrow \eta' J/\psi$

¹ Based on a fit to $\sigma(e^+ e^- \rightarrow \eta' J/\psi)$ from $\sqrt{s} = 4.18$ to 4.60 GeV assuming interfering $\psi(4160)$ and $\psi(4260)$ contributions. At $\sqrt{s} = 4.23$ GeV, $\sigma(e^+ e^- \rightarrow \eta' J/\psi) = 3.6 \pm 0.6 \pm 0.3$ pb.

² Solution I of the fit, corresponding to a phase of -0.03 ± 0.44 rad.

³ Solution II of the fit, corresponding to a phase of 2.54 ± 0.04 rad.

$\Gamma(\psi(2S) \pi^+ \pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_{14} \Gamma_1/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<4.3	90	¹ LIU	08H	RVUE 10.58 $e^+ e^- \rightarrow \psi(2S) \pi^+ \pi^- \gamma$
7.4 ^{+2.1} _{-1.7}		² LIU	08H	RVUE 10.58 $e^+ e^- \rightarrow \psi(2S) \pi^+ \pi^- \gamma$

¹ For constructive interference with the $\psi(4360)$ in a combined fit of AUBERT 07S and WANG 07D data with three resonances.

² For destructive interference with the $\psi(4360)$ in a combined fit of AUBERT 07S and WANG 07D data with three resonances.

$\Gamma(\phi\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{20}\Gamma_1/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.4	90	AUBERT,BE 06D	BABR	10.6 $e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$

$\Gamma(\phi f_0(980) \rightarrow \phi\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{21}\Gamma_1/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.28	90	¹ AUBERT 07AK	BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$

¹AUBERT 07AK reports $[\Gamma(\psi(4260) \rightarrow \phi f_0(980) \rightarrow \phi\pi^+\pi^-) \times \Gamma(\psi(4260) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] < 0.14$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = 49.2 \times 10^{-2}$.

$\Gamma(\Xi^-\Xi^+) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{43}\Gamma_1/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<2.7 × 10 ⁻⁴	90	ABLIKIM 20C	BES3	$e^+e^- \rightarrow \Xi^-\Xi^+$

$\Gamma(K_S^0 K^\pm \pi^\mp) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{44}\Gamma_1/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.04 ± 0.19 ± 0.09	¹ ABLIKIM 19AE	BES3	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$
0.0027 ± 0.0023 ± 0.0001	² ABLIKIM 19AE	BES3	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$
< 0.5 at 90% CL	AUBERT 08S	BABR	10.6 $e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp \gamma$

¹ Solution I of the fit including the $\psi(4160)$ with mass 4191 ± 5 MeV and width 70 ± 10 MeV from PDG 16 and the $\psi(4230)$ with mass $4219.6 \pm 3.3 \pm 5.1$ MeV and width $56.0 \pm 3.6 \pm 6.9$ MeV from GAO 17.

² Solution II of the fit including the $\psi(4160)$ with mass 4191 ± 5 MeV and width 70 ± 10 MeV from PDG 16 and the $\psi(4230)$ with mass $4219.6 \pm 3.3 \pm 5.1$ MeV and width $56.0 \pm 3.6 \pm 6.9$ MeV from GAO 17.

$\Gamma(K_S^0 K^\pm \pi^\mp \pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{45}\Gamma_1/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.05	90	ABLIKIM 19	BES3	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0$

$\Gamma(K_S^0 K^\pm \pi^\mp \eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{46}\Gamma_1/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.19	90	ABLIKIM 19	BES3	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp \eta$

$\Gamma(K^+K^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{47}\Gamma_1/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.6	90	AUBERT 08S	BABR	10.6 $e^+e^- \rightarrow K^+K^-\pi^0\gamma$
------	----	------------	------	---

$\Gamma(\chi_{c1}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{49}\Gamma_1/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<1.4	90	¹ HAN 15	BELL	10.58 $e^+e^- \rightarrow \chi_{c1}\gamma$

¹ Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.

$\Gamma(\chi_{c2}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{50}\Gamma_1/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
<4.0	90	¹ HAN	15	BELL	10.58 $e^+e^- \rightarrow \chi_{c2}\gamma$

¹ Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.

$\psi(4260)$ BRANCHING RATIOS

$\Gamma(J/\psi f_0(980), f_0(980) \rightarrow \pi^+\pi^-)/\Gamma(J/\psi\pi^+\pi^-)$					Γ_3/Γ_2
VALUE		DOCUMENT ID	TECN	COMMENT	
0.17 ± 0.13		¹ LEES	12AC	BABR	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^- J/\psi$

¹ Systematic uncertainties not estimated.

$\Gamma(Z_c(3900)^\pm \pi^\mp, Z_c^\pm \rightarrow J/\psi\pi^\pm)/\Gamma(J/\psi\pi^+\pi^-)$					Γ_4/Γ_2
VALUE		DOCUMENT ID	TECN	COMMENT	
0.215 ± 0.033 ± 0.075		¹ ABLIKIM	13T	BES3	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$
0.29 ± 0.08		² LIU	13B	BELL	$e^+e^- \rightarrow \gamma\pi^+\pi^- J/\psi$

¹ Assuming that the cross section of $e^+e^- \rightarrow \pi^+\pi^- J/\psi$ is fully due to the $\psi(4260)$.
² Systematic error not evaluated.

$\Gamma(J/\psi K_S^0 K_S^0)/\Gamma_{\text{total}}$					Γ_7/Γ
VALUE		DOCUMENT ID	TECN	COMMENT	
not seen		SHEN	14	BELL	9.4–10.9 $e^+e^- \rightarrow \gamma K_S^0 K_S^0 J/\psi$

$\Gamma(J/\psi\eta\pi^0)/\Gamma_{\text{total}}$					Γ_{12}/Γ
VALUE		DOCUMENT ID	TECN	COMMENT	
not seen		ABLIKIM	15Q	BES3	4.0–4.6 $e^+e^- \rightarrow J/\psi\eta\pi^0$

$\Gamma(\psi(2S)\pi^+\pi^-)/\Gamma(J/\psi\pi^+\pi^-)$					Γ_{14}/Γ_2
VALUE		DOCUMENT ID	TECN	COMMENT	
(0.11 ± 0.03 ± 0.03) to (0.55 ± 0.18 ± 0.19)		¹ ZHANG	17C	RVUE	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$ or $\psi(2S)$

¹ From a combined fit of BELLE, BABAR and BES3 $e^+e^- \rightarrow \pi^+\pi^- J/\psi$ and $e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$ data.

$\Gamma(h_c(1P)\pi^+\pi^-)/\Gamma(J/\psi\pi^+\pi^-)$					Γ_{19}/Γ_2
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<1.0	90	¹ PEDLAR	11	CLEO	$e^+e^- \rightarrow h_c(1P)\pi^+\pi^-$

¹ At $\sqrt{s} = 4260$ MeV, PEDLAR 11 measures $\sigma(e^+e^- \rightarrow h_c(1P)\pi^+\pi^-) = 32 \pm 17 \pm 6 \pm 6$ pb, where the errors are statistical, systematic, and due to uncertainty in $B(\psi(2S) \rightarrow \pi^0 h_c(1P))$, respectively.

$\Gamma(D\bar{D})/\Gamma(J/\psi\pi^+\pi^-)$ Γ_{22}/Γ_2

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.0	90	¹ AUBERT	07BE BABR	$e^+e^- \rightarrow D\bar{D}\gamma$
<4.0	90	CRONIN-HEN..09	CLEO	e^+e^-

¹ Using 4259 ± 10 MeV for the mass and 88 ± 24 MeV for the width of $\psi(4260)$. $\Gamma(D^0\bar{D}^0)/\Gamma_{\text{total}}$ Γ_{23}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
not seen	CRONIN-HEN..09	CLEO	$e^+e^- \rightarrow D^0\bar{D}^0$
not seen	AUBERT	09M BABR	$e^+e^- \rightarrow D^0\bar{D}^0\gamma$
not seen	PAKHLOVA	08 BELL	$e^+e^- \rightarrow D^0\bar{D}^0\gamma$

 $\Gamma(D^+D^-)/\Gamma_{\text{total}}$ Γ_{24}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
not seen	CRONIN-HEN..09	CLEO	$e^+e^- \rightarrow D^+D^-$
not seen	AUBERT	09M BABR	$e^+e^- \rightarrow D^+D^-\gamma$
not seen	PAKHLOVA	08 BELL	$e^+e^- \rightarrow D^+D^-\gamma$

 $\Gamma(D^*\bar{D}^0+c.c.)/\Gamma(J/\psi\pi^+\pi^-)$ Γ_{25}/Γ_2

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<34	90	AUBERT	09M BABR	$e^+e^- \rightarrow \gamma D^*\bar{D}^0$
<45	90	CRONIN-HEN..09	CLEO	e^+e^-

 $\Gamma(D^*(2007)^0\bar{D}^0+c.c.)/\Gamma_{\text{total}}$ Γ_{26}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
not seen	CRONIN-HEN..09	CLEO	$e^+e^- \rightarrow D^{*0}\bar{D}^0$
not seen	AUBERT	09M BABR	$e^+e^- \rightarrow D^{*0}\bar{D}^0\gamma$

 $\Gamma(D^*(2010)^+D^-+c.c.)/\Gamma_{\text{total}}$ Γ_{27}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
not seen	CRONIN-HEN..09	CLEO	$e^+e^- \rightarrow D^{*+}D^-$
not seen	PAKHLOVA	07 BELL	$e^+e^- \rightarrow D^{*+}D^-\gamma$
not seen	AUBERT	09M BABR	$e^+e^- \rightarrow D^{*+}D^-\gamma$

 $\Gamma(D^*\bar{D}^*)/\Gamma(J/\psi\pi^+\pi^-)$ Γ_{28}/Γ_2

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<11	90	CRONIN-HEN..09	CLEO	e^+e^-
<40	90	AUBERT	09M BABR	$e^+e^- \rightarrow \gamma D^*\bar{D}^*$

$\Gamma(D^*(2007)^0 \bar{D}^*(2007)^0)/\Gamma_{\text{total}}$ Γ_{29}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^{*0} \bar{D}^{*0}$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
not seen	AUBERT 09M BABR		$e^+ e^- \rightarrow D^{*0} \bar{D}^{*0} \gamma$

$\Gamma(D^*(2010)^+ D^*(2010)^-)/\Gamma_{\text{total}}$ Γ_{30}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^{*+} D^{*-}$
not seen	PAKHLOVA 07 BELL		$e^+ e^- \rightarrow D^{*+} D^{*-} \gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
not seen	AUBERT 09M BABR		$e^+ e^- \rightarrow D^{*+} D^{*-} \gamma$

$\Gamma(D^0 D^- \pi^+ + \text{c.c. (excl. } D^*(2007)^0 \bar{D}^{*0} + \text{c.c., } D^*(2010)^+ D^- + \text{c.c.))}/\Gamma_{\text{total}}$ Γ_{32}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	PAKHLOVA 08A BELL		$10.6 e^+ e^- \rightarrow D^0 D^- \pi^+ \gamma$

$\Gamma(D \bar{D}^* \pi + \text{c.c. (excl. } D^* \bar{D}^*)/\Gamma_{\text{total}}$ Γ_{33}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^* \bar{D} \pi$

$\Gamma(D \bar{D}^* \pi + \text{c.c. (excl. } D^* \bar{D}^*)/\Gamma(J/\psi \pi^+ \pi^-)$ Γ_{33}/Γ_2

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<15	90	CRONIN-HEN..09	CLEO	$e^+ e^-$

$\Gamma(D^0 D^{*-} \pi^+ + \text{c.c. (excl. } D^*(2010)^+ D^*(2010)^-)/\Gamma_{\text{total}}$ Γ_{34}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	PAKHLOVA 09 BELL		$e^+ e^- \rightarrow D^0 D^{*-} \pi^+ \gamma$

$\Gamma(D^0 D^*(2010)^- \pi^+ + \text{c.c.})/\Gamma(J/\psi \pi^+ \pi^-)$ Γ_{35}/Γ_2

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<9	90	PAKHLOVA 09 BELL		$e^+ e^- \rightarrow D^0 D^{*-} \pi^+$

$\Gamma(D^0 D^*(2010)^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_{35}/\Gamma \times \Gamma_1/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.42 × 10 ⁻⁶	90	¹ PAKHLOVA 09 BELL		$e^+ e^- \rightarrow D^0 D^{*-} \pi^+$

¹ Using 4263⁺⁸₋₉ MeV for the mass of $\psi(4260)$.

$\Gamma(D^* \bar{D}^* \pi)/\Gamma_{\text{total}}$ Γ_{37}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^* \bar{D}^* \pi$

$\Gamma(D^* \bar{D}^* \pi)/\Gamma(J/\psi \pi^+ \pi^-)$ Γ_{37}/Γ_2

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<8.2	90	CRONIN-HEN..09	CLEO	$e^+ e^-$

$\Gamma(D_1(2420)\bar{D} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{36}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
not seen	¹ ABLIKIM	19AR BES3	$e^+e^- \rightarrow \pi^+\pi^-D\bar{D}$

¹ Results from a measurement of $\sigma(e^+e^- \rightarrow D_1(2420)\bar{D} + \text{c.c.})$ between $\sqrt{s} = 4.3$ and 4.6 GeV.

$\Gamma(D_s^+ D_s^-)/\Gamma_{\text{total}}$ Γ_{38}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
not seen	DEL-AMO-SA..10N	BABR	$e^+e^- \rightarrow D_s^+ D_s^- \gamma$
not seen	CRONIN-HEN..09	CLEO	$e^+e^- \rightarrow D_s^+ D_s^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen	PAKHLOVA	11	BELL $e^+e^- \rightarrow D_s^+ D_s^- \gamma$
----------	----------	----	--

$\Gamma(D_s^+ D_s^-)/\Gamma(J/\psi\pi^+\pi^-)$ Γ_{38}/Γ_2

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.7	95	DEL-AMO-SA..10N	BABR	10.6 e^+e^-

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.3	90	CRONIN-HEN..09	CLEO	e^+e^-
------	----	----------------	------	----------

$\Gamma(D_s^{*+} D_s^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{39}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
not seen	DEL-AMO-SA..10N	BABR	$e^+e^- \rightarrow D_s^{*+} D_s^- \gamma$
not seen	CRONIN-HEN..09	CLEO	$e^+e^- \rightarrow D_s^{*+} D_s^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen	PAKHLOVA	11	BELL $e^+e^- \rightarrow D_s^{*+} D_s^- \gamma$
----------	----------	----	---

$\Gamma(D_s^{*+} D_s^- + \text{c.c.})/\Gamma(J/\psi\pi^+\pi^-)$ Γ_{39}/Γ_2

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 0.8	90	CRONIN-HEN..09	CLEO	e^+e^-

• • • We do not use the following data for averages, fits, limits, etc. • • •

<44	95	DEL-AMO-SA..10N	BABR	10.6 e^+e^-
-----	----	-----------------	------	---------------

$\Gamma(D_s^{*+} D_s^{*-})/\Gamma_{\text{total}}$ Γ_{40}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
not seen	CRONIN-HEN..09	CLEO	$e^+e^- \rightarrow D_s^{*+} D_s^{*-}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen	PAKHLOVA	11	BELL $e^+e^- \rightarrow D_s^{*+} D_s^{*-} \gamma$
----------	----------	----	--

not seen	DEL-AMO-SA..10N	BABR	$e^+e^- \rightarrow D_s^{*+} D_s^{*-} \gamma$
----------	-----------------	------	---

$\Gamma(D_s^{*+} D_s^{*-})/\Gamma(J/\psi\pi^+\pi^-)$ Γ_{40}/Γ_2

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 9.5	90	CRONIN-HEN..09	CLEO	e^+e^-

• • • We do not use the following data for averages, fits, limits, etc. • • •

<30	95	DEL-AMO-SA..10N	BABR	10.6 e^+e^-
-----	----	-----------------	------	---------------

$\Gamma(p\bar{p})/\Gamma(J/\psi\pi^+\pi^-)$		Γ_{41}/Γ_2		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.13	90	¹ AUBERT	06B BABR	$e^+e^- \rightarrow p\bar{p}\gamma$

¹ Using 4259 ± 10 MeV for the mass and 88 ± 24 MeV for the width of $\psi(4260)$.

$\Gamma(p\bar{p}\pi^0)/\Gamma(J/\psi\pi^+\pi^-)$		Γ_{42}/Γ_2		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 2×10^{-4}	90	ABLIKIM	17F BES3	$e^+e^- \rightarrow \psi(4260) \rightarrow$ hadrons

Radiative decays

$\Gamma(\eta_c(1S)\gamma)/\Gamma_{\text{total}}$		Γ_{48}/Γ		
VALUE		DOCUMENT ID	TECN	COMMENT
possibly seen		¹ ABLIKIM	17W	$e^+e^- \rightarrow \gamma\eta_c(1S)$

¹ Significance ranges from 4.2σ to as low as 1.5σ for a flat component plus $\psi(4260)$ spectrum. Needs confirmation.

$\Gamma(\chi_{c1}(3872)\gamma)/\Gamma_{\text{total}}$		Γ_{51}/Γ		
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	20 ± 5	ABLIKIM	14 BES3	$e^+e^- \rightarrow J/\psi\pi^+\pi^-\gamma$

$\psi(4260)$ REFERENCES

ABLIKIM	20A	PR D101 012008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	20C	PRL 124 032002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19	PR D99 012003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AE	PR D99 072005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AR	PR D100 032005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18N	PR D97 071101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17B	PRL 118 092001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17F	PL B771 45	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17W	PR D96 051101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
GAO	17	PR D95 092007	X.Y. Gao, C.P. Shen, C.Z. Yuan	
ZHANG	17B	PR D96 054008	J. Zhang, J. Zhang	
ZHANG	17C	EPJ C77 727	J. Zhang, L. Yuan	
PDG	16	CP C40 100001	C. Patrignani <i>et al.</i>	(PDG Collab.)
ABLIKIM	15Q	PR D92 012008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
HAN	15	PR D92 012011	Y.L. Han <i>et al.</i>	(BELLE Collab.)
ABLIKIM	14	PRL 112 092001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
SHEN	14	PR D89 072015	C.P. Shen <i>et al.</i>	(BELLE Collab.)
ABLIKIM	13T	PRL 110 252001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LIU	13B	PRL 110 252002	Z.Q. Liu <i>et al.</i>	(BELLE Collab.)
WANG	13B	PR D87 051101	X.L. Wang <i>et al.</i>	(BELLE Collab.)
LEES	12AC	PR D86 051102	J.P. Lees <i>et al.</i>	(BABAR Collab.)
PAKHLOVA	11	PR D83 011101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
PEDLAR	11	PRL 107 041803	T. Pedlar <i>et al.</i>	(CLEO Collab.)
DEL-AMO-SA...	10N	PR D82 052004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
AUBERT	09M	PR D79 092001	B. Aubert <i>et al.</i>	(BABAR Collab.)
CRONIN-HEN...	09	PR D80 072001	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)
PAKHLOVA	09	PR D80 091101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
AUBERT	08S	PR D77 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)
LIU	08H	PR D78 014032	Z.Q. Liu, X.S. Qin, C.Z. Yuan	
PAKHLOVA	08	PR D77 011103	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
PAKHLOVA	08A	PRL 100 062001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
YUAN	08	PR D77 011105	C.Z. Yuan <i>et al.</i>	(BELLE Collab.)
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07BE	PR D76 111105	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07S	PRL 98 212001	B. Aubert <i>et al.</i>	(BABAR Collab.)

PAKHLOVA	07	PRL 98 092001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
WANG	07D	PRL 99 142002	X.L. Wang <i>et al.</i>	(BELLE Collab.)
YUAN	07	PRL 99 182004	C.Z. Yuan <i>et al.</i>	(BELLE Collab.)
AUBERT	06B	PR D73 012005	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,BE	06D	PR D74 091103	B. Aubert <i>et al.</i>	(BABAR Collab.)
HE	06B	PR D74 091104	Q. He <i>et al.</i>	(CLEO Collab.)
AUBERT,B	05I	PRL 95 142001	B. Aubert <i>et al.</i>	(BABAR Collab.)
