

$B_{s1}(5830)^0$

$$I(J^P) = 0(1^+)$$

I, J, P need confirmation.

Quantum numbers shown are quark-model predictions.

 $B_{s1}(5830)^0$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
5828.70 ± 0.20 OUR FIT			
5828.65 ± 0.24 OUR AVERAGE			
5828.78 ± 0.09 ± 0.29	SIRUNYAN	18DF CMS	pp at 8 TeV
5828.40 ± 0.04 ± 0.41	¹ AAIJ	130 LHCb	pp at 7 TeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
5829.4 ± 0.7	² AALTONEN	08k CDF	Repl. by AALTONEN 14i
¹ Uses $B_{s1}(5830)^0 \rightarrow B^{*+} K^-$ decay.			
² Uses two-body decays into K^- and B^+ mesons reconstructed as $B^+ \rightarrow J/\psi K^+$, $J/\psi \rightarrow \mu^+ \mu^-$ or $B^+ \rightarrow \bar{D}^0 \pi^+$, $\bar{D}^0 \rightarrow K^+ \pi^-$.			

 $m_{B_{s1}^0} - m_{B^{*+}}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
503.99 ± 0.17 OUR FIT			
504.03 ± 0.12 ± 0.15	¹ AALTONEN	14i CDF	$p\bar{p}$ at 1.96 TeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
504.41 ± 0.21 ± 0.14	² AALTONEN	08k CDF	Repl. by AALTONEN 14i
¹ AALTONEN 14i reports $m_{B_{s1}(5830)^0} - m_{B^{*+}} - m_{K^-} = 10.35 \pm 0.12 \pm 0.15$ MeV which we adjusted by the K^- mass.			
² Uses two-body decays into K^- and B^+ mesons reconstructed as $B^+ \rightarrow J/\psi K^+$, $J/\psi \rightarrow \mu^+ \mu^-$ or $B^+ \rightarrow \bar{D}^0 \pi^+$, $\bar{D}^0 \rightarrow K^+ \pi^-$.			

 $B_{s1}(5830)^0$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
0.5 ± 0.3 ± 0.3	AALTONEN	14i CDF	$p\bar{p}$ at 1.96 TeV

 $B_{s1}(5830)^0$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $B^{*+} K^-$	seen
Γ_2 $B^{*0} K_S^0$	

 $B_{s1}(5830)^0$ BRANCHING RATIOS

$\Gamma(B^{*+} K^-)/\Gamma_{\text{total}}$	Γ_1/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
seen	AALTONEN	08k CDF	$p\bar{p}$ at 1.96 TeV

$$\Gamma(B^{*0} K_S^0) / \Gamma(B^{*+} K^-) \qquad \Gamma_2 / \Gamma_1$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.49 ± 0.12 ± 0.07	¹ SIRUNYAN	18DF CMS	<i>pp</i> at 8 TeV

¹ With the branching fractions $B(B^+ \rightarrow J/\psi K^+) = (1.026 \pm 0.031) \times 10^{-3}$ and $B(B^0 \rightarrow J/\psi K^{*0}) = (1.28 \pm 0.05) \times 10^{-3}$.

$B_{s1}(5830)^0$ REFERENCES

SIRUNYAN	18DF EPJ C78 939	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
AALTONEN	14I PR D90 012013	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AAIJ	13O PRL 110 151803	R. Aaij <i>et al.</i>	(LHCb Collab.)
AALTONEN	08K PRL 100 082001	T. Aaltonen <i>et al.</i>	(CDF Collab.)