

**$a_2(1700)$** 

$$I^G(J^{PC}) = 1^-(2^{++})$$

 **$a_2(1700)$  MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1698 ± 44</b>		<sup>1</sup> AMSLER 02	CBAR	0.9 $\bar{p}p \rightarrow \pi^0 \eta \eta$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1686 ± 22 $^{+19}_{-7}$		<sup>2</sup> KOPF 21	RVUE	0.9 $p\bar{p} \rightarrow \pi^0 \pi^0 \eta$ , $\pi^0 \eta \eta$ , $\pi^0 K^+ K^-$ and 191 $\pi^- p \rightarrow$ $\pi^- \pi^- \pi^+ p$
1638.9 ± 2.3 $^{+57.4}_{-0.1}$		<sup>3</sup> ALBRECHT 20	RVUE	0.9 $\bar{p}p \rightarrow \pi^0 \pi^0 \eta$ , $\pi^0 \eta \eta$ , $\pi^0 K^+ K^-$
1722 ± 15 ± 67		<sup>4</sup> RODAS 19	JPAC	191 $\pi^- p \rightarrow \eta^{(\prime)} \pi^- p$
1681 $^{+22}_{-35}$	46M	<sup>5,6</sup> AGHASYAN 18B	COMP	190 $\pi^- p \rightarrow$ $\pi^- \pi^+ \pi^- p$
1720 ± 10 ± 60		<sup>7</sup> JACKURA 18	JPAC	$\pi^- p \rightarrow \eta \pi^- p$
1726 ± 12 ± 25		<sup>6</sup> ABLIKIM 17K	BES3	$\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$
1675 ± 25		ANISOVICH 09	RVUE	0.0 $\bar{p}p$ , $\pi N$
1722 ± 9 ± 15	18k	<sup>8</sup> SCHEGELSKY 06	RVUE	$\gamma \gamma \rightarrow \pi^+ \pi^- \pi^0$
1702 ± 7	80k	<sup>9</sup> UMAN 06	E835	5.2 $\bar{p}p \rightarrow \eta \eta \pi^0$
1721 ± 13 ± 44	145k	LU 05	B852	18 $\pi^- p \rightarrow \omega \pi^- \pi^0 p$
1737 ± 5 ± 7		ABE 04	BELL	10.6 $e^+ e^- \rightarrow$ $e^+ e^- K^+ K^-$
1767 ± 14	221	<sup>10</sup> ACCIARRI 01H	L3	$\gamma \gamma \rightarrow K_S^0 K_S^0$ , $E_{\text{cm}}^{\text{ee}} =$ 91, 183–209 GeV
1660 ± 40		<sup>6</sup> ABELE 99B	CBAR	1.94 $\bar{p}p \rightarrow \pi^0 \eta \eta$
~ 1775		<sup>11</sup> GRYGOREV 99	SPEC	40 $\pi^- p \rightarrow K_S^0 K_S^0 n$
1752 ± 21 ± 4		ACCIARRI 97T	L3	$\gamma \gamma \rightarrow \pi^+ \pi^- \pi^0$

<sup>1</sup> T-matrix pole.<sup>2</sup> From T-matrix pole based on combined fit of Crystal Barrel and  $\pi\pi$  scattering data (ALBRECHT 20), and COMPASS data (ADOLPH 15), using a coupled-channel model of  $\eta\pi$ ,  $\eta'\pi$  and  $K\bar{K}$  systems.<sup>3</sup> T-matrix pole, 2 poles, 2 channels ( $\pi\eta$ ,  $K\bar{K}$ ).<sup>4</sup> The coupled-channel analysis of both the  $\eta\pi$  and  $\eta'\pi$  systems using ADOLPH 15 data. The mass is extracted from the T-matrix pole.<sup>5</sup> Statistical error negligible.<sup>6</sup> Breit-Wigner mass.<sup>7</sup> Superseded by RODAS 19.<sup>8</sup> From analysis of L3 data at 183–209 GeV.<sup>9</sup> Statistical error only.<sup>10</sup> Spin 2 dominant, isospin not determined, could also be  $I=1$ .<sup>11</sup> Possibly two  $J^P = 2^+$  resonances with isospins 0 and 1. **$a_2(1700)$  WIDTH**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>265 ± 55</b>		<sup>1</sup> AMSLER 02	CBAR	0.9 $\bar{p}p \rightarrow \pi^0 \eta \eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

421	$\pm 75$	$\begin{matrix} +64 \\ -57 \end{matrix}$	2	KOPF	21	RVUE	0.9	$p\bar{p} \rightarrow \pi^0\pi^0\eta, \pi^0\eta\eta,$ $\pi^0K^+K^-$ and 191 $\pi^-p \rightarrow \pi^-\pi^-\pi^+p$	
224.0	$\pm$	$\begin{matrix} 2.5^+ 1.8 \\ -48.3 \end{matrix}$	3	ALBRECHT	20	RVUE	0.9	$\bar{p}p \rightarrow \pi^0\pi^0\eta,$ $\pi^0\eta\eta, \pi^0K^+K^-$	
247	$\pm 17$	$\pm 63$	4	RODAS	19	JPAC	191	$\pi^-p \rightarrow \eta^{(\prime)}\pi^-p$	
436	$\begin{matrix} + 20 \\ - 16 \end{matrix}$		46M	5,6	AGHASYAN	18B	COMP	190	$\pi^-p \rightarrow \pi^-\pi^+\pi^-p$
280	$\pm 10$	$\pm 70$	7	JACKURA	18	JPAC		$\pi^-p \rightarrow \eta\pi^-p$	
190	$\pm 18$	$\pm 30$	6	ABLIKIM	17K	BES3		$\psi(2S) \rightarrow \gamma\eta\pi^+\pi^-$	
270	$\begin{matrix} + 50 \\ - 20 \end{matrix}$			ANISOVICH	09	RVUE	0.0	$\bar{p}p, \pi N$	
336	$\pm 20$	$\pm 20$	18k	8	SCHEGELSKY	06	RVUE	$\gamma\gamma \rightarrow \pi^+\pi^-\pi^0$	
417	$\pm 19$		80k	9	UMAN	06	E835	5.2	$\bar{p}p \rightarrow \eta\eta\pi^0$
279	$\pm 49$	$\pm 66$	145k		LU	05	B852	18	$\pi^-p \rightarrow \omega\pi^-\pi^0p$
151	$\pm 22$	$\pm 24$			ABE	04	BELL	10.6	$e^+e^- \rightarrow$ $e^+e^-K^+K^-$
187	$\pm 60$		221	10	ACCIARRI	01H	L3	$\gamma\gamma \rightarrow K_S^0K_S^0, E_{\text{cm}}^{\text{ee}} =$ 91, 183–209 GeV	
280	$\pm 70$			6	ABELE	99B	CBAR	1.94	$\bar{p}p \rightarrow \pi^0\eta\eta$
150	$\pm 110$	$\pm 34$			ACCIARRI	97T	L3	$\gamma\gamma \rightarrow \pi^+\pi^-\pi^0$	

<sup>1</sup> T-matrix pole.

<sup>2</sup> From T-matrix pole based on combined fit of Crystal Barrel and  $\pi\pi$  scattering data (ALBRECHT 20), and COMPASS data (ADOLPH 15), using a coupled-channel model of  $\eta\pi, \eta'\pi$  and  $K\bar{K}$  systems.

<sup>3</sup> T-matrix pole, 2 poles, 2 channels ( $\pi\eta, K\bar{K}$ ).

<sup>4</sup> The coupled-channel analysis of both the  $\eta\pi$  and  $\eta'\pi$  systems using ADOLPH 15 data. The width is extracted from the T-matrix pole.

<sup>5</sup> Statistical error negligible.

<sup>6</sup> Breit-Wigner width.

<sup>7</sup> Superseded by RODAS 19.

<sup>8</sup> From analysis of L3 data at 183–209 GeV.

<sup>9</sup> Statistical error only.

<sup>10</sup> Spin 2 dominant, isospin not determined, could also be  $l=1$ .

## $a_2(1700)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\eta\pi$	(3.6 $\pm$ 1.1 ) %
$\Gamma_2$ $\eta'\pi$	
$\Gamma_3$ $\gamma\gamma$	(1.13 $\pm$ 0.30) $\times 10^{-6}$
$\Gamma_4$ $\rho\pi$	seen
$\Gamma_5$ $f_2(1270)\pi$	seen
$\Gamma_6$ $K\bar{K}$	(1.9 $\pm$ 1.2 ) %
$\Gamma_7$ $\omega\pi^-\pi^0$	seen
$\Gamma_8$ $\omega\rho$	seen

**$a_2(1700)$  PARTIAL WIDTHS** **$\Gamma(\eta\pi)$   $\Gamma_1$** 

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>9.5 \pm 2.0</math></b>	870	<sup>1</sup> SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$

<sup>1</sup>From analysis of L3 data at 91 and 183–209 GeV, using  $a_2(1700)$  mass of 1730 MeV and width of 340 MeV, and SU(3) relations.

 **$\Gamma(\gamma\gamma)$   $\Gamma_3$** 

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.30 \pm 0.05</math></b>	870	<sup>1</sup> SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$

<sup>1</sup>From analysis of L3 data at 91 and 183–209 GeV, using  $a_2(1700)$  mass of 1730 MeV and width of 340 MeV, and SU(3) relations.

 **$\Gamma(K\bar{K})$   $\Gamma_6$** 

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>5.0 \pm 3.0</math></b>	870	<sup>1</sup> SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$

<sup>1</sup>From analysis of L3 data at 91 and 183–209 GeV, using  $a_2(1700)$  mass of 1730 MeV and width of 340 MeV, and SU(3) relations.

 **$a_2(1700)$   $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$**  **$[\Gamma(\rho\pi) + \Gamma(f_2(1270)\pi)] \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$   $(\Gamma_4 + \Gamma_5)\Gamma_3/\Gamma$** 

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.29 \pm 0.04 \pm 0.02</math></b>		ACCIARRI	97T L3	$\gamma\gamma \rightarrow \pi^+ \pi^- \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.37^{+0.12}_{-0.08} \pm 0.10$	18k	<sup>1</sup> SCHEGELSKY 06	RVUE	$\gamma\gamma \rightarrow \pi^+ \pi^- \pi^0$
---------------------------------	-----	----------------------------	------	--

<sup>1</sup>From analysis of L3 data at 183–209 GeV.

 **$\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$   $\Gamma_6\Gamma_3/\Gamma$** 

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$20.6 \pm 4.2 \pm 4.6$	<sup>1</sup> ABE	04 BELL	$10.6 e^+ e^- \rightarrow e^+ e^- K^+ K^-$
$49 \pm 11 \pm 13$	<sup>2</sup> ACCIARRI	01H L3	$\gamma\gamma \rightarrow K_S^0 K_S^0, E_{\text{cm}}^{ee} = 91, 183\text{--}209 \text{ GeV}$

<sup>1</sup>Assuming spin 2.

<sup>2</sup>Spin 2 dominant, isospin not determined, could also be  $I=1$ .

 **$a_2(1700)$  BRANCHING RATIOS** **$\Gamma(\rho\pi)/\Gamma(f_2(1270)\pi)$   $\Gamma_4/\Gamma_5$** 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$3.4 \pm 0.4 \pm 0.1$	18k	<sup>1</sup> SCHEGELSKY 06	RVUE	$\gamma\gamma \rightarrow \pi^+ \pi^- \pi^0$
-----------------------	-----	----------------------------	------	--

<sup>1</sup>From analysis of L3 data at 183–209 GeV.

$\Gamma(K\bar{K})/\Gamma(\eta\pi)$  $\Gamma_6/\Gamma_1$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.029 \pm 0.04$ $^{+0.011}_{-0.012}$	<sup>1</sup> KOPF	21	RVUE 0.9 $p\bar{p} \rightarrow \pi^0 \pi^0 \eta, \pi^0 \eta \eta,$ $\pi^0 K^+ K^-$ and 191 $\pi^- p \rightarrow$ $\pi^- \pi^- \pi^+ p$
---------------------------------------	-------------------	----	--

$4.134 \pm 0.106$ $^{+4.909}_{-2.988}$	<sup>2</sup> ALBRECHT	20	RVUE 0.9 $p\bar{p} \rightarrow \pi^0 \pi^0 \eta, \pi^0 \eta \eta,$ $\pi^0 K^+ K^-$
--	-----------------------	----	---

<sup>1</sup> From T-matrix pole based on combined fit of Crystal Barrel and  $\pi\pi$  scattering data (ALBRECHT 20), and COMPASS data (ADOLPH 15), using a coupled-channel model of  $\eta\pi, \eta'\pi$  and  $K\bar{K}$  systems.

<sup>2</sup> Residues from T-matrix pole, 2 poles, 2 channels ( $\pi\eta, K\bar{K}$ ).

 $\Gamma(\eta'\pi)/\Gamma(\eta\pi)$  $\Gamma_2/\Gamma_1$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.035 \pm 0.044$ $^{+0.069}_{-0.012}$	<sup>1</sup> KOPF	21	RVUE 0.9 $p\bar{p} \rightarrow \pi^0 \pi^0 \eta, \pi^0 \eta \eta,$ $\pi^0 K^+ K^-$ and 191 $\pi^- p \rightarrow$ $\pi^- \pi^- \pi^+ p$
--	-------------------	----	--

<sup>1</sup> From T-matrix pole based on combined fit of Crystal Barrel and  $\pi\pi$  scattering data (ALBRECHT 20), and COMPASS data (ADOLPH 15), using a coupled-channel model of  $\eta\pi, \eta'\pi$  and  $K\bar{K}$  systems.

 **$a_2(1700)$  REFERENCES**

KOPF	21	EPJ C81 1056	B. Kopf <i>et al.</i>	(BOCH)
ALBRECHT	20	EPJ C80 453	M. Albrecht <i>et al.</i>	(Crystal Barrel Collab.)
RODAS	19	PRL 122 042002	A. Rodas <i>et al.</i>	(JPAC Collab.)
AGHASYAN	18B	PR D98 092003	M. Aghasyan <i>et al.</i>	(COMPASS Collab.)
JACKURA	18	PL B779 464	A. Jackura <i>et al.</i>	(JPAC and COMPASS Collab.)
ABLIKIM	17K	PR D95 032002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ADOLPH	15	PL B740 303	M. Adolph <i>et al.</i>	(COMPASS Collab.)
ANISOVICH	09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev	
SCHEGELSKY	06	EPJ A27 199	V.A. Schegelsky <i>et al.</i>	
SCHEGELSKY	06A	EPJ A27 207	V.A. Schegelsky <i>et al.</i>	
UMAN	06	PR D73 052009	I. Uman <i>et al.</i>	(FNAL E835)
LU	05	PRL 94 032002	M. Lu <i>et al.</i>	(BNL E852 Collab.)
ABE	04	EPJ C32 323	K. Abe <i>et al.</i>	(BELLE Collab.)
AMSLER	02	EPJ C23 29	C. Amsler <i>et al.</i>	
ACCIARRI	01H	PL B501 173	M. Acciarri <i>et al.</i>	(L3 Collab.)
ABELE	99B	EPJ C8 67	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
GRYGOREV	99	PAN 62 470	V.K. Grygorev <i>et al.</i>	
ACCIARRI	97T	PL B413 147	M. Acciarri <i>et al.</i>	(L3 Collab.)