



$$I^G(J^{PC}) = 0^+(0^{-+})$$

We have omitted some results that have been superseded by later experiments. The omitted results may be found in our 1988 edition Physics Letters **B204** (1988).

η MASS

Recent measurements resolve the obvious inconsistency in previous η mass measurements in favor of the higher value first reported by NA48 (LAI 02). We use only precise measurements consistent with this higher mass value for our η mass average.

| <u>VALUE (MeV)</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|-------------------------|-------------|---|
| 547.862±0.017 OUR AVERAGE | | | | |
| 547.865±0.031±0.062 | | NIKOLAEV | 14 | CRYB $\gamma p \rightarrow p\eta$ |
| 547.873±0.005±0.027 | 1M | GOSLAWSKI | 12 | SPEC $dp \rightarrow {}^3\text{He}\eta$ |
| 547.874±0.007±0.029 | | AMBROSINO | 07B | KLOE $e^+e^- \rightarrow \phi \rightarrow \eta\gamma$ |
| 547.785±0.017±0.057 | 16k | MILLER | 07 | CLEO $\psi(2S) \rightarrow J/\psi\eta$ |
| 547.843±0.030±0.041 | 1134 | LAI | 02 | NA48 $\eta \rightarrow 3\pi^0$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| 547.311±0.028±0.032 | | ¹ ABDEL-BARY | 05 | SPEC $dp \rightarrow {}^3\text{He}\eta$ |
| 547.12 ±0.06 ±0.25 | | KRUSCHE | 95D | SPEC $\gamma p \rightarrow \eta p$, threshold |
| 547.30 ±0.15 | | PLOUIN | 92 | SPEC $dp \rightarrow {}^3\text{He}\eta$ |
| 547.45 ±0.25 | | DUANE | 74 | SPEC $\pi^- p \rightarrow n$ neutrals |
| 548.2 ±0.65 | | FOSTER | 65C | HBC |
| 549.0 ±0.7 | 148 | FOELSCH | 64 | HBC |
| 548.0 ±1.0 | 91 | ALFF-... | 62 | HBC |
| 549.0 ±1.2 | 53 | BASTIEN | 62 | HBC |

¹ ABDEL-BARY 05 disagrees significantly with recent measurements of similar or better precision. See comment in the header.

η WIDTH

This is the partial decay rate $\Gamma(\eta \rightarrow \gamma\gamma)$ divided by the fitted branching fraction for that mode. See the note at the start of the $\Gamma(2\gamma)$ data block, next below.

| <u>VALUE (keV)</u> | <u>DOCUMENT ID</u> |
|--------------------------|--------------------|
| 1.31±0.05 OUR FIT | |

η DECAY MODES

| Mode | Fraction (Γ_i/Γ) | Scale factor/ Confidence level |
|--------------------------|--------------------------------|-----------------------------------|
| Neutral modes | | |
| Γ_1 neutral modes | (71.96±0.30) % | S=1.3 |
| Γ_2 2γ | (39.36±0.18) % | S=1.1 |
| Γ_3 $3\pi^0$ | (32.57±0.21) % | S=1.2 |

| | | | |
|------------|------------------|----------------------------------|--------|
| Γ_4 | $\pi^0 2\gamma$ | $(2.55 \pm 0.22) \times 10^{-4}$ | |
| Γ_5 | $2\pi^0 2\gamma$ | $< 1.2 \times 10^{-3}$ | CL=90% |
| Γ_6 | 4γ | $< 2.8 \times 10^{-4}$ | CL=90% |
| Γ_7 | invisible | $< 1.0 \times 10^{-4}$ | CL=90% |

Charged modes

| | | | |
|---------------|---------------------------------------|----------------------------------|--------|
| Γ_8 | charged modes | $(28.04 \pm 0.30) \%$ | S=1.3 |
| Γ_9 | $\pi^+ \pi^- \pi^0$ | $(23.02 \pm 0.25) \%$ | S=1.2 |
| Γ_{10} | $\pi^+ \pi^- \gamma$ | $(4.28 \pm 0.07) \%$ | S=1.1 |
| Γ_{11} | $e^+ e^- \gamma$ | $(6.9 \pm 0.4) \times 10^{-3}$ | S=1.2 |
| Γ_{12} | $\mu^+ \mu^- \gamma$ | $(3.1 \pm 0.4) \times 10^{-4}$ | |
| Γ_{13} | $e^+ e^-$ | $< 7 \times 10^{-7}$ | CL=90% |
| Γ_{14} | $\mu^+ \mu^-$ | $(5.8 \pm 0.8) \times 10^{-6}$ | |
| Γ_{15} | $2e^+ 2e^-$ | $(2.40 \pm 0.22) \times 10^{-5}$ | |
| Γ_{16} | $\pi^+ \pi^- e^+ e^- (\gamma)$ | $(2.68 \pm 0.11) \times 10^{-4}$ | |
| Γ_{17} | $e^+ e^- \mu^+ \mu^-$ | $< 1.6 \times 10^{-4}$ | CL=90% |
| Γ_{18} | $2\mu^+ 2\mu^-$ | $< 3.6 \times 10^{-4}$ | CL=90% |
| Γ_{19} | $\mu^+ \mu^- \pi^+ \pi^-$ | $< 3.6 \times 10^{-4}$ | CL=90% |
| Γ_{20} | $\pi^+ e^- \bar{\nu}_e + \text{c.c.}$ | $< 1.7 \times 10^{-4}$ | CL=90% |
| Γ_{21} | $\pi^+ \pi^- 2\gamma$ | $< 2.1 \times 10^{-3}$ | |
| Γ_{22} | $\pi^+ \pi^- \pi^0 \gamma$ | $< 6 \times 10^{-4}$ | CL=90% |
| Γ_{23} | $\pi^0 \mu^+ \mu^- \gamma$ | $< 3 \times 10^{-6}$ | CL=90% |

Charge conjugation (C), Parity (P), Charge conjugation \times Parity (CP), or Lepton Family number (LF) violating modes

| | | | | |
|---------------|-------------------------|-------|------------------------|--------|
| Γ_{24} | $\pi^0 \gamma$ | C [a] | $< 9 \times 10^{-5}$ | CL=90% |
| Γ_{25} | $\pi^+ \pi^-$ | P, CP | $< 4.4 \times 10^{-6}$ | CL=90% |
| Γ_{26} | $2\pi^0$ | P, CP | $< 3.5 \times 10^{-4}$ | CL=90% |
| Γ_{27} | $2\pi^0 \gamma$ | C | $< 5 \times 10^{-4}$ | CL=90% |
| Γ_{28} | $3\pi^0 \gamma$ | C | $< 6 \times 10^{-5}$ | CL=90% |
| Γ_{29} | 3γ | C | $< 1.6 \times 10^{-5}$ | CL=90% |
| Γ_{30} | $4\pi^0$ | P, CP | $< 6.9 \times 10^{-7}$ | CL=90% |
| Γ_{31} | $\pi^0 e^+ e^-$ | C [b] | $< 8 \times 10^{-6}$ | CL=90% |
| Γ_{32} | $\pi^0 \mu^+ \mu^-$ | C [b] | $< 5 \times 10^{-6}$ | CL=90% |
| Γ_{33} | $\mu^+ e^- + \mu^- e^+$ | LF | $< 6 \times 10^{-6}$ | CL=90% |

[a] Forbidden by angular momentum conservation.

[b] C parity forbids this to occur as a single-photon process.

CONSTRAINED FIT INFORMATION

An overall fit to 2 decay rate and 22 branching ratios uses 54 measurements and one constraint to determine 9 parameters. The overall fit has a $\chi^2 = 46.2$ for 46 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

| | | | | | | | | | |
|----------|-----|-------|-------|-------|-------|----------|----------|----------|----------|
| x_3 | 12 | | | | | | | | |
| x_4 | 4 | 0 | | | | | | | |
| x_9 | -69 | -76 | -3 | | | | | | |
| x_{10} | -48 | -52 | -2 | 53 | | | | | |
| x_{11} | -8 | -7 | 0 | -4 | -3 | | | | |
| x_{12} | -1 | -1 | 0 | 0 | 0 | 0 | | | |
| x_{16} | 0 | 0 | 0 | 0 | 0 | 0 | 0 | | |
| Γ | -13 | -1 | -32 | 9 | 6 | 1 | 0 | 0 | |
| | | x_2 | x_3 | x_4 | x_9 | x_{10} | x_{11} | x_{12} | x_{16} |

| Mode | Rate (keV) | Scale factor |
|--|----------------------------------|--------------|
| Γ_2 2γ | 0.515 ± 0.018 | |
| Γ_3 $3\pi^0$ | 0.426 ± 0.015 | |
| Γ_4 $\pi^0 2\gamma$ | $(3.34 \pm 0.28) \times 10^{-4}$ | |
| Γ_9 $\pi^+ \pi^- \pi^0$ | 0.301 ± 0.011 | |
| Γ_{10} $\pi^+ \pi^- \gamma$ | 0.0559 ± 0.0022 | |
| Γ_{11} $e^+ e^- \gamma$ | 0.0090 ± 0.0006 | 1.2 |
| Γ_{12} $\mu^+ \mu^- \gamma$ | $(4.1 \pm 0.5) \times 10^{-4}$ | |
| Γ_{16} $\pi^+ \pi^- e^+ e^- (\gamma)$ | $(3.50 \pm 0.19) \times 10^{-4}$ | |

η DECAY RATES

$\Gamma(2\gamma)$

Γ_2

See the table immediately above giving the fitted decay rates. Following the advice of NEFKENS 02, we have removed the Primakoff-effect measurement from the average. See also the "Note on the Decay Width $\Gamma(\eta \rightarrow \gamma\gamma)$," in our 1994 edition, Phys. Rev. **D50**, 1 August 1994, Part I, p. 1451, for a discussion of the various measurements.

| VALUE (keV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|------|---|
| 0.515 ± 0.018 OUR FIT | | | | |
| 0.516 ± 0.018 OUR AVERAGE | | | | |
| $0.520 \pm 0.020 \pm 0.013$ | | BABUSCI | 13A | KLOE $e^+ e^- \rightarrow e^+ e^- \eta$ |
| $0.51 \pm 0.12 \pm 0.05$ | 36 | BARU | 90 | MD1 $e^+ e^- \rightarrow e^+ e^- \eta$ |
| $0.490 \pm 0.010 \pm 0.048$ | 2287 | ROE | 90 | ASP $e^+ e^- \rightarrow e^+ e^- \eta$ |
| $0.514 \pm 0.017 \pm 0.035$ | 1295 | WILLIAMS | 88 | CBAL $e^+ e^- \rightarrow e^+ e^- \eta$ |
| $0.53 \pm 0.04 \pm 0.04$ | | BARTEL | 85E | JADE $e^+ e^- \rightarrow e^+ e^- \eta$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|--------------------|----|------------------------|-----|------|---------------------------------|
| 0.476 ± 0.062 | | ¹ RODRIGUES | 08 | CNTR | Reanalysis |
| 0.64 ± 0.14 ± 0.13 | | AIHARA | 86 | TPC | $e^+e^- \rightarrow e^+e^-\eta$ |
| 0.56 ± 0.16 | 56 | WEINSTEIN | 83 | CBAL | $e^+e^- \rightarrow e^+e^-\eta$ |
| 0.324 ± 0.046 | | BROWMAN | 74B | CNTR | Primakoff effect |
| 1.00 ± 0.22 | | ² BEMPORAD | 67 | CNTR | Primakoff effect |

¹ RODRIGUES 08 uses a more sophisticated calculation for the inelastic background due to incoherent photoproduction to reanalyze the η photoproduction data on Be and Cu at 9 GeV from BROWMAN 74B. This brings the value of $\Gamma(\eta \rightarrow 2\gamma)$ in line with direct measurements of the width. The error here is only statistical.

² BEMPORAD 67 gives $\Gamma(2\gamma) = 1.21 \pm 0.26$ keV assuming $\Gamma(2\gamma)/\Gamma(\text{total}) = 0.314$.

Bemporad private communication gives $\Gamma(2\gamma)^2/\Gamma(\text{total}) = 0.380 \pm 0.083$. We evaluate this using $\Gamma(2\gamma)/\Gamma(\text{total}) = 0.38 \pm 0.01$. Not included in average because the uncertainty resulting from the separation of the coulomb and nuclear amplitudes has apparently been underestimated.

| $\Gamma(\pi^0 2\gamma)$ | | | | | Γ_4 |
|-------------------------|------|-------------|------|---------|-------------------------------|
| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT | |
| 0.334 ± 0.028 | | | | | OUR FIT |
| 0.33 ± 0.03 | 1200 | NEFKENS | 14 | CRYB | $\gamma p \rightarrow \eta p$ |

η BRANCHING RATIOS

Neutral modes

| $\Gamma(\text{neutral modes})/\Gamma_{\text{total}}$ | | $\Gamma_1/\Gamma = (\Gamma_2 + \Gamma_3 + \Gamma_4)/\Gamma$ | | | |
|--|------|---|------|---------|--|
| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT | |
| 0.7196 ± 0.0030 | | | | | OUR FIT Error includes scale factor of 1.3. |
| 0.705 ± 0.008 | 16k | BASILE | 71D | CNTR | MM spectrometer |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|-------------|--|----------|----|------|--|
| 0.79 ± 0.08 | | BUNIATOV | 67 | OSPK | |
|-------------|--|----------|----|------|--|

| $\Gamma(2\gamma)/\Gamma_{\text{total}}$ | | Γ_2/Γ | | | |
|---|------|-------------------|------|---------|--|
| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT | |
| 39.36 ± 0.18 | | | | | OUR FIT Error includes scale factor of 1.1. |
| 39.53 ± 0.33 | | | | | OUR AVERAGE |

| | | | | | |
|---------------------|-----|----------------------|----------|---------------------------------|------------------------------------|
| 39.86 ± 0.04 ± 0.99 | 2m | ¹ ABLIKIM | 21AMBES3 | $J/\psi \rightarrow \gamma\eta$ | |
| 39.49 ± 0.17 ± 0.30 | 65k | ABEGG | 96 | SPEC | $pd \rightarrow {}^3\text{He}\eta$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|---------------------|-----|--------------------|----|------|-----------------------------------|
| 38.45 ± 0.40 ± 0.36 | 14k | ² LOPEZ | 07 | CLEO | $\psi(2S) \rightarrow J/\psi\eta$ |
|---------------------|-----|--------------------|----|------|-----------------------------------|

¹ ABLIKIM 21AM normalize the branching ratio ($\eta \rightarrow \gamma\gamma$) to $B(J/\psi \rightarrow \gamma\eta)$, which they measured absolutely.

² Not independent of other results listed for LOPEZ 07. Assuming decays of $\eta \rightarrow \gamma\gamma$, $3\pi^0$, $\pi^+\pi^-\pi^0$, $\pi^+\pi^-\gamma$, and $e^+e^-\gamma$ account for all η decays within a contribution of 0.3% to the systematic error.

| $\Gamma(2\gamma)/\Gamma(\text{neutral modes})$ | | $\Gamma_2/\Gamma_1 = \Gamma_2/(\Gamma_2 + \Gamma_3 + \Gamma_4)$ | | | |
|--|------|---|------|---------|--|
| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT | |
| 0.5470 ± 0.0018 | | | | | OUR FIT |
| 0.548 ± 0.023 | | | | | OUR AVERAGE Error includes scale factor of 1.5. |
| 0.535 ± 0.018 | | BUTTRAM | 70 | OSPK | |
| 0.59 ± 0.033 | | BUNIATOV | 67 | OSPK | |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|--------------|--------------|------------|----|------|---------------|
| 0.52 ±0.09 | 88 | ABROSIMOV | 80 | HLBC | |
| 0.60 ±0.14 | 113 | KENDALL | 74 | OSPK | |
| 0.57 ±0.09 | | STRUGALSKI | 71 | HLBC | |
| 0.579 ±0.052 | | FELDMAN | 67 | OSPK | |
| 0.416 ±0.044 | | DIGIUGNO | 66 | CNTR | Error doubled |
| 0.44 ±0.07 | | GRUNHAUS | 66 | OSPK | |
| 0.39 ±0.06 | ¹ | JONES | 66 | CNTR | |

¹This result from combining cross sections from two different experiments.

$\Gamma(3\pi^0)/\Gamma_{\text{total}}$

Γ_3/Γ

| <u>VALUE (units 10^{-2})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------------------------------|----------------------|-------------|---------------------------------|
| 32.57±0.21 OUR FIT | Error includes scale factor of 1.2. | | | |
| 31.96±0.07±0.84 | 280k | ¹ ABLIKIM | 21AMBES3 | $J/\psi \rightarrow \gamma\eta$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|-----------------|------|--------------------|----|------|-----------------------------------|
| 34.03±0.56±0.49 | 1821 | ² LOPEZ | 07 | CLEO | $\psi(2S) \rightarrow J/\psi\eta$ |
|-----------------|------|--------------------|----|------|-----------------------------------|

¹ ABLIKIM 21AM normalize the branching ratio ($\eta \rightarrow 3\pi^0$) to $B(J/\psi \rightarrow \gamma\eta)$, which they measured absolutely.

² Not independent of other results listed for LOPEZ 07. Assuming decays of $\eta \rightarrow \gamma\gamma$, $3\pi^0$, $\pi^+\pi^-\pi^0$, $\pi^+\pi^-\gamma$, and $e^+e^-\gamma$ account for all η decays within a contribution of 0.3% to the systematic error.

$\Gamma(3\pi^0)/\Gamma(\text{neutral modes})$

$\Gamma_3/\Gamma_1 = \Gamma_3/(\Gamma_2+\Gamma_3+\Gamma_4)$

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | |
|---|-------------|--------------------|-------------|----------------|--|
| 0.4526±0.0019 OUR FIT | | | | | |
| 0.439 ±0.024 | | BUTTRAM | 70 | OSPK | |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| 0.44 ±0.08 | 75 | ABROSIMOV | 80 | HLBC | |
| 0.32 ±0.09 | | STRUGALSKI | 71 | HLBC | |
| 0.41 ±0.033 | | BUNIATOV | 67 | OSPK | Not indep. of $\Gamma(2\gamma)/\Gamma(\text{neutral modes})$ |
| 0.177 ±0.035 | | FELDMAN | 67 | OSPK | |
| 0.209 ±0.054 | | DIGIUGNO | 66 | CNTR | Error doubled |
| 0.29 ±0.10 | | GRUNHAUS | 66 | OSPK | |

$\Gamma(3\pi^0)/\Gamma(2\gamma)$

Γ_3/Γ_2

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------------------|-------------|-------------------------|-------------|---|
| 0.827±0.006 OUR FIT | | | | |
| 0.829±0.007 OUR AVERAGE | | | | |
| 0.884±0.022±0.019 | 1821 | LOPEZ | 07 | CLEO $\psi(2S) \rightarrow J/\psi\eta$ |
| 0.817±0.012±0.032 | 17.4k | ¹ AKHMETSHIN | 05 | CMD2 $e^+e^- \rightarrow \phi \rightarrow \eta\gamma$ |
| 0.826±0.024 | | ACHASOV | 00D | SND $e^+e^- \rightarrow \phi \rightarrow \eta\gamma$ |
| 0.832±0.005±0.012 | | KRUSCHE | 95D | SPEC $\gamma p \rightarrow \eta p$, threshold |
| 0.841±0.034 | | AMSLER | 93 | CBAR $\bar{p}p \rightarrow \pi^+\pi^-\eta$ at rest |
| 0.822±0.009 | | ALDE | 84 | GAM2 |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|-------------------|--|---------|-----|------|-----------------|
| 0.796±0.016±0.016 | | ACHASOV | 00 | SND | See ACHASOV 00D |
| 0.91 ±0.14 | | COX | 70B | HBC | |
| 0.75 ±0.09 | | DEVONS | 70 | OSPK | |

| | | | |
|------------|--------|-----|--------------------------|
| 0.88 ±0.16 | BALTAY | 67D | DBC |
| 1.1 ±0.2 | CENCE | 67 | OSPK |
| 1.25 ±0.39 | BACCI | 63 | CNTR Inverse BR reported |

¹ Uses result from AKHMETSHIN 01B.

$\Gamma(\pi^0 2\gamma)/\Gamma_{\text{total}}$ Γ_4/Γ

Early results are summarized in the review by LANDSBERG 85.

| VALUE (units 10^{-4}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------|------------------------|------|--|
| 2.55±0.22 OUR FIT | | | | | |
| 2.21±0.24±0.47 | | ≈ 500 | ¹ PRAKHOV | 08 | CRYB $\pi^- p \rightarrow \eta n \approx$ threshold |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| 3.5 ±0.7 ±0.6 | | 1.6k | ^{2,3} PRAKHOV | 05 | CRYB See PRAKHOV 08 |
| <8.4 | 90 | 7 | ACHASOV | 01D | SND $e^+ e^- \rightarrow \phi \rightarrow \eta \gamma$ |
| <30 | 90 | 0 | DAVYDOV | 81 | GAM2 $\pi^- p \rightarrow \eta n$ |

¹ PRAKHOV 08 is a reanalysis of the data of PRAKHOV 05, using for the first time the invariant-mass spectrum of the two photons.

² Normalized using $\Gamma(\eta \rightarrow 2\gamma)/\Gamma = 0.3943 \pm 0.0026$.

³ This measurement and the independent analysis of the same data by KNECHT 04 both imply a lower value of $\Gamma(\pi^0 2\gamma)$ than the one obtained by ALDE 84 from $\Gamma(\pi^0 2\gamma)/\Gamma(2\gamma)$.

$\Gamma(\pi^0 2\gamma)/\Gamma(2\gamma)$ Γ_4/Γ_2

| VALUE (units 10^{-3}) | EVTS | DOCUMENT ID | TECN | CHG | COMMENT |
|---|------|-------------|------|------|-------------|
| 0.65±0.06 OUR FIT | | | | | |
| 1.8 ±0.4 | | ALDE | 84 | GAM2 | 0 |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| 2.5 ±0.6 | 70 | BINON | 82 | GAM2 | See ALDE 84 |

$\Gamma(\pi^0 2\gamma)/\Gamma(3\pi^0)$ Γ_4/Γ_3

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|------|---------|
| 7.8±0.7 OUR FIT | | | |

• • • We do not use the following data for averages, fits, limits, etc. • • •

8.3±2.8±1.4 ¹ KNECHT 04 CRYB $\pi^- p \rightarrow n \eta$

¹ Independent analysis of same data as PRAKHOV 05.

$\Gamma(2\pi^0 2\gamma)/\Gamma_{\text{total}}$ Γ_5/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|----------------------|------|--|
| <1.2 × 10⁻³ | 90 | ¹ NEFKENS | 05A | CRYB $p(720 \text{ MeV}/c) \pi^- \rightarrow n \eta$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| <4.0 × 10 ⁻³ | 90 | BLIK | 07 | GAM4 $\pi^- p \rightarrow \eta n$ |

¹ Measurement is done in limited $\gamma\gamma$ energy range.

$\Gamma(4\gamma)/\Gamma_{\text{total}}$ Γ_6/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|----------------------------------|-----|-------------|------|-----------------------------------|
| <2.8 × 10⁻⁴ | 90 | BLIK | 07 | GAM4 $\pi^- p \rightarrow \eta n$ |

$\Gamma(\text{invisible})/\Gamma(2\gamma)$ Γ_7/Γ_2

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|----------------------------------|-----|----------------------|------|-------------------------------------|
| <2.6 × 10⁻⁴ | 90 | ¹ ABLIKIM | 13 | BES3 $J/\psi \rightarrow \phi \eta$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1.65 \times 10^{-3}$ 90 ² ABLIKIM 06Q BES2 $J/\psi \rightarrow \phi\eta$

¹ Based on 225M J/ψ decays.

² Based on 58M J/ψ decays.

————— Charged modes —————

$\Gamma(\text{charged modes})/\Gamma_{\text{total}}$ $\Gamma_8/\Gamma = (\Gamma_9 + \Gamma_{10} + \Gamma_{11} + \Gamma_{12} + \Gamma_{16})/\Gamma$

VALUE DOCUMENT ID
0.2804 ± 0.0030 OUR FIT Error includes scale factor of 1.3.

$\Gamma(\pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE (units 10^{-2}) EVTS DOCUMENT ID TECN COMMENT
23.02 ± 0.25 OUR FIT Error includes scale factor of 1.2.

23.04 ± 0.03 ± 0.54 60k ¹ ABLIKIM 21AMBES3 $J/\psi \rightarrow \gamma\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

22.60 ± 0.35 ± 0.29 3915 ² LOPEZ 07 CLEO $\psi(2S) \rightarrow J/\psi\eta$

¹ ABLIKIM 21AM normalize the branching ratio ($\eta \rightarrow \pi^+ \pi^- \pi^0$) to $B(J/\psi \rightarrow \gamma\eta)$, which they measured absolutely.

² Not independent of other results listed for LOPEZ 07. Assuming decays of $\eta \rightarrow \gamma\gamma$, $3\pi^0$, $\pi^+ \pi^- \pi^0$, $\pi^+ \pi^- \gamma$, and $e^+ e^- \gamma$ account for all η decays within a contribution of 0.3% to the systematic error.

$\Gamma(\text{neutral modes})/\Gamma(\pi^+ \pi^- \pi^0)$ $\Gamma_1/\Gamma_9 = (\Gamma_2 + \Gamma_3 + \Gamma_4)/\Gamma_9$

VALUE EVTS DOCUMENT ID TECN
3.13 ± 0.05 OUR FIT Error includes scale factor of 1.3.

3.26 ± 0.30 OUR AVERAGE

| | | | | |
|-------------|-----|-------------------------|-----|------|
| 2.54 ± 1.89 | 74 | KENDALL | 74 | OSPK |
| 3.4 ± 1.1 | 29 | AGUILAR-... | 72B | HBC |
| 2.83 ± 0.80 | 70 | ¹ BLOODWO... | 72B | HBC |
| 3.6 ± 0.6 | 244 | FLATTE | 67B | HBC |
| 2.89 ± 0.56 | | ALFF-... | 66 | HBC |
| 3.6 ± 0.8 | 50 | KRAEMER | 64 | DBC |
| 3.8 ± 1.1 | | PAULI | 64 | DBC |

¹ Error increased from published value 0.5 by Bloodworth (private communication).

$\Gamma(2\gamma)/\Gamma(\pi^+ \pi^- \pi^0)$ Γ_2/Γ_9

VALUE EVTS DOCUMENT ID TECN COMMENT
1.710 ± 0.025 OUR FIT Error includes scale factor of 1.2.

1.70 ± 0.04 OUR AVERAGE

| | | | | | |
|-----------------------|------|--------------------|-----|------|---|
| 1.704 ± 0.032 ± 0.026 | 3915 | ¹ LOPEZ | 07 | CLEO | $\psi(2S) \rightarrow J/\psi\eta$ |
| 1.61 ± 0.14 | | ABLIKIM | 06E | BES2 | $e^+ e^- \rightarrow J/\psi \rightarrow \eta\gamma$ |
| 1.78 ± 0.10 ± 0.13 | 1077 | AMSLER | 95 | CBAR | $\bar{p}p \rightarrow \pi^+ \pi^- \eta$ at rest |
| 1.72 ± 0.25 | 401 | BAGLIN | 69 | HLBC | |
| 1.61 ± 0.39 | | FOSTER | 65 | HBC | |

¹ LOPEZ 07 reports $\Gamma(\eta \rightarrow \pi^+ \pi^- \pi^0) / \Gamma(\eta \rightarrow 2\gamma) = \Gamma_9/\Gamma_2 = 0.587 \pm 0.011 \pm 0.009$.

$$\Gamma(3\pi^0)/\Gamma(\pi^+\pi^-\pi^0) \qquad \Gamma_3/\Gamma_9$$

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------|------|-------------|------|---------|
|-------|------|-------------|------|---------|

1.415 ± 0.023 OUR FIT Error includes scale factor of 1.2.

1.48 ± 0.05 OUR AVERAGE

| | | | | | |
|--|------|-------------------------|-----|------|--|
| 1.46 ± 0.03 ± 0.09 | | ACHASOV | 06A | SND | $e^+e^- \rightarrow \eta\gamma$ |
| 1.52 ± 0.04 ± 0.08 | 23k | ¹ AKHMETSHIN | 01B | CMD2 | $e^+e^- \rightarrow \phi \rightarrow \eta\gamma$ |
| 1.44 ± 0.09 ± 0.10 | 1627 | AMSLER | 95 | CBAR | $\bar{p}p \rightarrow \pi^+\pi^-\eta$ at rest |
| 1.50 ^{+0.15} _{-0.29} | 199 | BAGLIN | 69 | HLBC | |
| 1.47 ^{+0.20} _{-0.17} | | BULLOCK | 68 | HLBC | |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|-------------|--|----------|-----|------|--|
| 1.3 ± 0.4 | | BAGLIN | 67B | HLBC | |
| 0.90 ± 0.24 | | FOSTER | 65 | HBC | |
| 2.0 ± 1.0 | | FOELSCH | 64 | HBC | |
| 0.83 ± 0.32 | | CRAWFORD | 63 | HBC | |

¹ AKHMETSHIN 01B uses results from AKHMETSHIN 99F.

$$\Gamma(\pi^+\pi^-\pi^0)/[\Gamma(2\gamma) + \Gamma(3\pi^0)] \qquad \Gamma_9/(\Gamma_2+\Gamma_3)$$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

0.320 ± 0.005 OUR FIT Error includes scale factor of 1.2.

0.304 ± 0.012 ACHASOV 00D SND $e^+e^- \rightarrow \phi \rightarrow \eta\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.3141 ± 0.0081 ± 0.0058 ACHASOV 00B SND See ACHASOV 00D

$$\Gamma(\pi^+\pi^-\gamma)/\Gamma_{\text{total}} \qquad \Gamma_{10}/\Gamma$$

| VALUE (units 10 ⁻²) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------------------|------|-------------|------|---------|
|---------------------------------|------|-------------|------|---------|

4.28 ± 0.07 OUR FIT Error includes scale factor of 1.1.

4.38 ± 0.02 ± 0.10 200k ¹ ABLIKIM 21AMBES3 $J/\psi \rightarrow \gamma\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.96 ± 0.14 ± 0.14 859 ² LOPEZ 07 CLEO $\psi(2S) \rightarrow J/\psi\eta$

¹ ABLIKIM 21AM normalize the branching ratio ($\eta \rightarrow \pi^+\pi^-\gamma$) to $B(J/\psi \rightarrow \gamma\eta)$, which they measured absolutely.

² Not independent of other results listed for LOPEZ 07. Assuming decays of $\eta \rightarrow \gamma\gamma$, $3\pi^0$, $\pi^+\pi^-\pi^0$, $\pi^+\pi^-\gamma$, and $e^+e^-\gamma$ account for all η decays within a contribution of 0.3% to the systematic error.

$$\Gamma(\pi^+\pi^-\gamma)/\Gamma(\pi^+\pi^-\pi^0) \qquad \Gamma_{10}/\Gamma_9$$

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------|------|-------------|------|---------|
|-------|------|-------------|------|---------|

0.1858 ± 0.0025 OUR FIT

0.1847 ± 0.0030 OUR AVERAGE Error includes scale factor of 1.1.

0.1856 ± 0.0005 ± 0.0028 200k BABUSCI 13 KLOE $e^+e^- \rightarrow \phi \rightarrow \eta\gamma$

0.175 ± 0.007 ± 0.006 859 LOPEZ 07 CLEO $\psi(2S) \rightarrow J/\psi\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|---------------|------|------------|-----|------|--|
| 0.209 ± 0.004 | 18k | THALER | 73 | ASPK | |
| 0.201 ± 0.006 | 7250 | GORMLEY | 70 | ASPK | |
| 0.28 ± 0.04 | | BALTAY | 67B | DBC | |
| 0.25 ± 0.035 | | LITCHFIELD | 67 | DBC | |
| 0.30 ± 0.06 | | CRAWFORD | 66 | HBC | |
| 0.196 ± 0.041 | | FOSTER | 65C | HBC | |

$\Gamma(e^+e^-\gamma)/\Gamma_{\text{total}}$ Γ_{11}/Γ

| VALUE (units 10^{-3}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|----------|--------------------|------|--|
| 6.9 ± 0.4 OUR FIT | | | | Error includes scale factor of 1.2. |
| 6.7 ± 0.5 OUR AVERAGE | | | | Error includes scale factor of 1.2. |
| 6.6 ± 0.4 ± 0.4 | 1345 | BERGHAUSER 11 | SPEC | $\gamma p \rightarrow p\eta$ |
| 7.8 ± 0.5 ± 0.8 | 435 ± 31 | BERLOWSKI 08 | WASA | $pd \rightarrow {}^3\text{He} \eta$ |
| 5.15 ± 0.62 ± 0.74 | 283 | ACHASOV 01B | SND | $e^+e^- \rightarrow \phi \rightarrow \eta\gamma$ |
| 7.10 ± 0.64 ± 0.46 | 323 | AKHMETSHIN 01 | CMD2 | $e^+e^- \rightarrow \phi \rightarrow \eta\gamma$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 9.4 ± 0.7 ± 0.5 | 172 | ¹ LOPEZ | 07 | CLEO $\psi(2S) \rightarrow J/\psi\eta$ |

¹Not independent of other results listed for LOPEZ 07. Assuming decays of $\eta \rightarrow \gamma\gamma$, $3\pi^0$, $\pi^+\pi^-\pi^0$, $\pi^+\pi^-\gamma$, and $e^+e^-\gamma$ account for all η decays within a contribution of 0.3% to the systematic error.

 $\Gamma(e^+e^-\gamma)/\Gamma(\pi^+\pi^-\gamma)$ Γ_{11}/Γ_{10}

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------------|------|-------------|------|--|
| 0.161 ± 0.010 OUR FIT | | | | Error includes scale factor of 1.2. |
| 0.237 ± 0.021 ± 0.015 | 172 | LOPEZ | 07 | CLEO $\psi(2S) \rightarrow J/\psi\eta$ |

 $\Gamma(e^+e^-\gamma)/\Gamma(\pi^+\pi^-\pi^0)$ Γ_{11}/Γ_9

| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------------|------|-------------|------|-------------------------------------|
| 2.98 ± 0.19 OUR FIT | | | | Error includes scale factor of 1.3. |
| 2.1 ± 0.5 | 80 | JANE | 75B | OSPK See the erratum |

 $\Gamma(\text{neutral modes})/[\Gamma(\pi^+\pi^-\pi^0) + \Gamma(\pi^+\pi^-\gamma) + \Gamma(e^+e^-\gamma)]$
 $\Gamma_1/(\Gamma_9 + \Gamma_{10} + \Gamma_{11}) = (\Gamma_2 + \Gamma_3 + \Gamma_4)/(\Gamma_9 + \Gamma_{10} + \Gamma_{11})$

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|----------------------|------|-------------------------------------|
| 2.57 ± 0.04 OUR FIT | | | | Error includes scale factor of 1.3. |
| 2.64 ± 0.23 | | BALTAY | 67B | DBC |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 4.5 ± 1.0 | 280 | ¹ JAMES | 66 | HBC |
| 3.20 ± 1.26 | 53 | ¹ BASTIEN | 62 | HBC |
| 2.5 ± 1.0 | 10 | ¹ PICKUP | 62 | HBC |

¹These experiments are not used in the averages as they do not separate clearly $\eta \rightarrow \pi^+\pi^-\pi^0$ and $\eta \rightarrow \pi^+\pi^-\gamma$ from each other. The reported values thus probably contain some unknown fraction of $\eta \rightarrow \pi^+\pi^-\gamma$.

 $\Gamma(2\gamma)/[\Gamma(\pi^+\pi^-\pi^0) + \Gamma(\pi^+\pi^-\gamma) + \Gamma(e^+e^-\gamma)]$ $\Gamma_2/(\Gamma_9 + \Gamma_{10} + \Gamma_{11})$

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------------|------|-------------|------|-------------------------------------|
| 1.407 ± 0.020 OUR FIT | | | | Error includes scale factor of 1.2. |
| 1.1 ± 0.4 OUR AVERAGE | | | | |
| 1.51 ± 0.93 | 75 | KENDALL | 74 | OSPK |
| 0.99 ± 0.48 | | CRAWFORD | 63 | HBC |

 $\Gamma(\mu^+\mu^-\gamma)/\Gamma_{\text{total}}$ Γ_{12}/Γ

| VALUE (units 10^{-4}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|---------------|------|------------------------------|
| 3.1 ± 0.4 OUR FIT | | | | |
| 3.1 ± 0.4 | 600 | DZHELYADIN 80 | SPEC | $\pi^- p \rightarrow \eta n$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 1.5 ± 0.75 | 100 | BUSHNIN | 78 | SPEC See DZHELYADIN 80 |

$\Gamma(e^+e^-)/\Gamma_{\text{total}}$ Γ_{13}/Γ

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|-------------------------|-------------|--|
| <7 × 10⁻⁷ | 90 | ACHASOV | 18B | CNTR $e^+e^- \rightarrow \eta$ Inverse reaction |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| <2.3 × 10 ⁻⁶ | 90 | AGAKISHIEV | 14 | $pp \rightarrow \eta + X$ |
| <5.6 × 10 ⁻⁶ | 90 | ¹ AGAKISHIEV | 12A | SPEC $pp \rightarrow \eta + X$ |
| <2.7 × 10 ⁻⁵ | 90 | BERLOWSKI | 08 | WASA $pd \rightarrow {}^3\text{He} \eta$ |
| <0.77 × 10 ⁻⁴ | 90 | BROWDER | 97B | CLE2 $e^+e^- \simeq 10.5 \text{ GeV}$ |
| <2 × 10 ⁻⁴ | 90 | WHITE | 96 | SPEC $pd \rightarrow \eta {}^3\text{He}$ |
| <3 × 10 ⁻⁴ | 90 | DAVIES | 74 | RVUE Uses ESTEN 67 |

¹AGAKISHIEV 12A uses a data sample of 3.5 GeV proton beam collisions on liquid hydrogen target collected by the HADES detector.

 $\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_{14}/Γ

| <u>VALUE (units 10⁻⁶)</u> | <u>CL%</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|-------------|--------------------|-------------|--|
| 5.8±0.8 OUR AVERAGE | | | | | |
| 5.7±0.7±0.5 | | 114 | ABEGG | 94 | SPEC $pd \rightarrow \eta {}^3\text{He}$ |
| 6.5±2.1 | | 27 | DZHELYADIN | 80B | SPEC $\pi^- p \rightarrow \eta n$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| 5.6 ^{+0.6} _{-0.7} ±0.5 | | 100 | KESSLER | 93 | SPEC See ABEGG 94 |
| < 20 | 95 | 0 | WEHMANN | 68 | OSPK |

 $\Gamma(\mu^+\mu^-)/\Gamma(2\gamma)$ Γ_{14}/Γ_2

| <u>VALUE (units 10⁻⁵)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> |
|---|--------------------|-------------|
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | |
| 5.9±2.2 | HYAMS | 69 |

 $\Gamma(2e^+2e^-)/\Gamma_{\text{total}}$ Γ_{15}/Γ

| <u>VALUE (units 10⁻⁵)</u> | <u>CL%</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|-------------|------------------------|-------------|---|
| 2.4±0.2±0.1 | | 362 | ¹ AMBROSINO | 11B | KLOE $e^+e^- \rightarrow \phi \rightarrow \eta\gamma$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| <9.7 | 90 | | BERLOWSKI | 08 | WASA $pd \rightarrow {}^3\text{He} \eta$ |
| <6.9 | 90 | | AKHMETSHIN | 01 | CMD2 $e^+e^- \rightarrow \phi \rightarrow \eta\gamma$ |

¹This measurement is fully inclusive (includes "2e⁺2e⁻γ" channel).

 $\Gamma(\pi^+\pi^-e^+e^-(\gamma))/\Gamma_{\text{total}}$ Γ_{16}/Γ

| <u>VALUE (units 10⁻⁴)</u> | <u>CL%</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|-------------|------------------------|-------------|---|
| 2.68±0.11 OUR FIT | | | | | |
| 2.68±0.09±0.07 | | 1555 ± 52 | ¹ AMBROSINO | 09B | KLOE $e^+e^- \rightarrow \phi \rightarrow \eta\gamma$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| 4.3 ^{+2.0} _{-1.6} ± 0.4 | | 16 | BERLOWSKI | 08 | WASA $pd \rightarrow {}^3\text{He} \eta$ |
| 4.3 ± 1.3 ± 0.4 | | 16 | BARGHOLTZ | 07 | CNTR See BERLOWSKI 08 |
| 3.7 ^{+2.5} _{-1.8} ± 0.3 | | 4 | AKHMETSHIN | 01 | CMD2 $e^+e^- \rightarrow \phi \rightarrow \eta\gamma$ |

¹This AMBROSINO 09B value includes radiative events.

| $\Gamma(e^+ e^- \mu^+ \mu^-)/\Gamma_{\text{total}}$ | | | | | Γ_{17}/Γ |
|---|-----|--------------|--------------------------|---|--|
| VALUE | CL% | DOCUMENT ID | TECN | COMMENT | |
| $<1.6 \times 10^{-4}$ | 90 | BERLOWSKI 08 | WASA | $pd \rightarrow {}^3\text{He } \eta$ | |
| $\Gamma(2\mu^+ 2\mu^-)/\Gamma_{\text{total}}$ | | | | | Γ_{18}/Γ |
| VALUE | CL% | DOCUMENT ID | TECN | COMMENT | |
| $<3.6 \times 10^{-4}$ | 90 | BERLOWSKI 08 | WASA | $pd \rightarrow {}^3\text{He } \eta$ | |
| $\Gamma(\mu^+ \mu^- \pi^+ \pi^-)/\Gamma_{\text{total}}$ | | | | | Γ_{19}/Γ |
| VALUE | CL% | DOCUMENT ID | TECN | COMMENT | |
| $<3.6 \times 10^{-4}$ | 90 | BERLOWSKI 08 | WASA | $pd \rightarrow {}^3\text{He } \eta$ | |
| $\Gamma(\pi^+ e^- \bar{\nu}_e + \text{c.c.})/\Gamma(\pi^+ \pi^- \pi^0)$ | | | | | Γ_{20}/Γ_9 |
| VALUE | CL% | DOCUMENT ID | TECN | COMMENT | |
| $<7.3 \times 10^{-4}$ | 90 | ABLIKIM 13G | BES3 | $J/\psi \rightarrow \phi \eta$ | |
| $\Gamma(\pi^+ \pi^- 2\gamma)/\Gamma(\pi^+ \pi^- \pi^0)$ | | | | | Γ_{21}/Γ_9 |
| VALUE | CL% | DOCUMENT ID | TECN | | |
| $< 9 \times 10^{-3}$ | | PRICE 67 | HBC | | |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| $<16 \times 10^{-3}$ | 95 | BALTAY 67B | DBC | | |
| $\Gamma(\pi^+ \pi^- \pi^0 \gamma)/\Gamma(\pi^+ \pi^- \pi^0)$ | | | | | Γ_{22}/Γ_9 |
| VALUE | CL% | EVTS | DOCUMENT ID | TECN | |
| $<0.24 \times 10^{-2}$ | 90 | 0 | THALER 73 | ASPK | |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| $<1.7 \times 10^{-2}$ | 90 | | ARNOLD 68 | HLBC | |
| $<1.6 \times 10^{-2}$ | 95 | | BALTAY 67B | DBC | |
| $<7.0 \times 10^{-2}$ | | | FLATTE 67 | HBC | |
| $<0.9 \times 10^{-2}$ | | | PRICE 67 | HBC | |
| $\Gamma(\pi^0 \mu^+ \mu^- \gamma)/\Gamma_{\text{total}}$ | | | | | Γ_{23}/Γ |
| VALUE | CL% | DOCUMENT ID | TECN | COMMENT | |
| $<3 \times 10^{-6}$ | 90 | DZHELADIN 81 | SPEC | $\pi^- p \rightarrow \eta n$ | |
| ————— Forbidden modes ————— | | | | | |
| $\Gamma(\pi^0 \gamma)/\Gamma_{\text{total}}$ | | | | | Γ_{24}/Γ |
| Forbidden by angular momentum conservation. | | | | | |
| VALUE | CL% | DOCUMENT ID | TECN | COMMENT | |
| $<9 \times 10^{-5}$ | 90 | NEFKENS 05A | CRYB | $p(720 \text{ MeV}/c) \pi^- \rightarrow n \eta$ | |
| $\Gamma(\pi^+ \pi^-)/\Gamma_{\text{total}}$ | | | | | Γ_{25}/Γ |
| Forbidden by P and CP invariance. | | | | | |
| VALUE | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
| $< 4.4 \times 10^{-6}$ | 90 | 83M | ¹ BABUSCI 20A | KLOE | $e^+ e^- \rightarrow \phi \rightarrow \eta \gamma$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|-------------------------|----|------|---------------------------|------|---|
| $< 5.3 \times 10^{-17}$ | | | ² ZHEVLAKOV 19 | THEO | from nEDM limits |
| $< 1.6 \times 10^{-5}$ | 90 | 25M | AAIJ | 17D | LHCB in $D \rightarrow \pi\pi\pi$ decays |
| $< 3.9 \times 10^{-4}$ | 90 | 225M | ABLIKIM | 11G | BES3 $e^+e^- \rightarrow J/\psi \rightarrow \eta\gamma$ |
| $< 1.3 \times 10^{-5}$ | 90 | 16M | AMBROSINO | 05A | KLOE $e^+e^- \rightarrow \phi \rightarrow \eta\gamma$ |
| $< 3.3 \times 10^{-4}$ | 90 | | AKHMETSHIN | 99B | CMD2 $e^+e^- \rightarrow \phi \rightarrow \eta\gamma$ |
| $< 9 \times 10^{-4}$ | 90 | | AKHMETSHIN | 97C | CMD2 See AKHMETSHIN 99B |
| $< 15 \times 10^{-4}$ | | 0 | THALER | 73 | ASPK |

¹BABUSCI 20A combines new data with the previous AMBROSINO 05A data, and thus supersedes AMBROSINO 05A.

²ZHEVLAKOV 19 derives the value from the experimental limits of nEDM by a calculation using an effective Lagrangian.

$\Gamma(2\pi^0)/\Gamma_{\text{total}}$ Γ_{26}/Γ

Forbidden by P and CP invariance.

| VALUE | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------|-----|------|-------------|------|-----------------------------------|
| $< 3.5 \times 10^{-4}$ | 90 | | BLIK | 07 | GAM4 $\pi^- p \rightarrow \eta n$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|-------------------------|----|------|---------------------------|------|---|
| $< 2.7 \times 10^{-17}$ | | | ¹ ZHEVLAKOV 19 | THEO | from nEDM limits |
| $< 6.9 \times 10^{-4}$ | 90 | 225M | ABLIKIM | 11G | BES3 $e^+e^- \rightarrow J/\psi \rightarrow \eta\gamma$ |
| $< 4.3 \times 10^{-4}$ | 90 | | AKHMETSHIN | 99C | CMD2 $e^+e^- \rightarrow \phi \rightarrow \eta\gamma$ |
| $< 6 \times 10^{-4}$ | 90 | | ² ACHASOV 98 | SND | $e^+e^- \rightarrow \phi \rightarrow \eta\gamma$ |

¹ZHEVLAKOV 19 derives the value from the experimental limits of nEDM by a calculation using an effective Lagrangian.

²ACHASOV 98 observes one event in a $\pm 3\sigma$ region around the η mass, while a Monte Carlo calculation gives 10 ± 5 events. The limit here is the Poisson upper limit for one observed event and no background.

$\Gamma(2\pi^0\gamma)/\Gamma_{\text{total}}$ Γ_{27}/Γ

Forbidden by C invariance.

| VALUE | CL% | DOCUMENT ID | TECN | CHG | COMMENT |
|----------------------|-----|-------------|------|-----|--|
| $< 5 \times 10^{-4}$ | 90 | NEFKENS 05 | CRYB | 0 | p(720 MeV/c) $\pi^- \rightarrow n\eta$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|-----------------------|----|------|----|------|------------------------------|
| $< 17 \times 10^{-4}$ | 90 | BLIK | 07 | GAM4 | $\pi^- p \rightarrow \eta n$ |
|-----------------------|----|------|----|------|------------------------------|

$\Gamma(3\pi^0\gamma)/\Gamma_{\text{total}}$ Γ_{28}/Γ

Forbidden by C invariance.

| VALUE | CL% | DOCUMENT ID | TECN | CHG | COMMENT |
|----------------------|-----|-------------|------|-----|--|
| $< 6 \times 10^{-5}$ | 90 | NEFKENS 05 | CRYB | 0 | p(720 MeV/c) $\pi^- \rightarrow n\eta$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|-----------------------|----|------|----|------|------------------------------|
| $< 24 \times 10^{-5}$ | 90 | BLIK | 07 | GAM4 | $\pi^- p \rightarrow \eta n$ |
|-----------------------|----|------|----|------|------------------------------|

$\Gamma(3\gamma)/\Gamma_{\text{total}}$ Γ_{29}/Γ

Forbidden by C invariance.

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-----|-------------|------|---|
| $< 16 \times 10^{-5}$ | 90 | BLIK | 07 | GAM4 $\pi^- p \rightarrow \eta n$ |
| $< 4 \times 10^{-5}$ | 90 | NEFKENS | 05A | CRYB p(720 MeV/c) $\pi^- \rightarrow n\eta$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-----------------------|----|---------|-----|---|
| $< 16 \times 10^{-5}$ | 90 | BLIK | 07 | GAM4 $\pi^- p \rightarrow \eta n$ |
| $< 4 \times 10^{-5}$ | 90 | NEFKENS | 05A | CRYB p(720 MeV/c) $\pi^- \rightarrow n\eta$ |

$\Gamma(3\gamma)/\Gamma(2\gamma)$ Γ_{29}/Γ_2

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>CHG</u> |
|------------------------|------------|--------------------|-------------|------------|
| $< 1.2 \times 10^{-3}$ | 95 | ALDE | 84 | GAM2 0 |

 $\Gamma(3\gamma)/\Gamma(3\pi^0)$ Γ_{29}/Γ_3

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|------------------------|------------|--------------------|-------------|------------------------------------|
| $< 4.9 \times 10^{-5}$ | 90 | ALOISIO | 04 | KLOE $\phi \rightarrow \eta\gamma$ |

 $\Gamma(4\pi^0)/\Gamma_{\text{total}}$ Γ_{30}/Γ Forbidden by P and CP invariance.

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|---|
| $< 6.9 \times 10^{-7}$ | 90 | PRAKHOV | 00 | CRYB $\pi^- p \rightarrow n\eta$, 720 MeV/ c |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| $< 200 \times 10^{-7}$ | 90 | BLIK | 07 | GAM4 $\pi^- p \rightarrow \eta n$ |

 $\Gamma(\pi^0 e^+ e^-)/\Gamma_{\text{total}}$ Γ_{31}/Γ C parity forbids this to occur as a single-photon process.

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|--|
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| $< 7.5 \times 10^{-6}$ | 90 | ADLARSON | 18C | WASA $pd \rightarrow \eta {}^3\text{He}$ |
| $< 1.6 \times 10^{-4}$ | 90 | MARTYNOV | 76 | HLBC |
| $< 8.4 \times 10^{-4}$ | 90 | BAZIN | 68 | DBC |
| $< 70 \times 10^{-4}$ | | RITTENBERG | 65 | HBC |

 $\Gamma(\pi^0 e^+ e^-)/\Gamma(\pi^+ \pi^- \pi^0)$ Γ_{31}/Γ_9 C parity forbids this to occur as a single-photon process.

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|--|
| $< 3.28 \times 10^{-5}$ | 90 | ADLARSON | 18C | WASA $pd \rightarrow \eta {}^3\text{He}$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| $< 1.9 \times 10^{-4}$ | 90 | JANE | 75 | OSPK |
| $< 42 \times 10^{-4}$ | 90 | BAGLIN | 67 | HLBC |
| $< 16 \times 10^{-4}$ | 90 | BILLING | 67 | HLBC |
| $< 77 \times 10^{-4}$ | | FOSTER | 65B | HBC |
| $< 110 \times 10^{-4}$ | | PRICE | 65 | HBC |

 $\Gamma(\pi^0 \mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{32}/Γ C parity forbids this to occur as a single-photon process.

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|-----------------------------------|
| $< 5 \times 10^{-6}$ | 90 | DZHELYADIN | 81 | SPEC $\pi^- p \rightarrow \eta n$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| $< 500 \times 10^{-6}$ | | WEHMANN | 68 | OSPK |

 $[\Gamma(\mu^+ e^-) + \Gamma(\mu^- e^+)]/\Gamma_{\text{total}}$ Γ_{33}/Γ

Forbidden by lepton family number conservation.

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------------|------------|--------------------|-------------|--|
| $< 6 \times 10^{-6}$ | 90 | WHITE | 96 | SPEC $pd \rightarrow \eta {}^3\text{He}$ |

η C-NONCONSERVING DECAY PARAMETERS **$\pi^+ \pi^- \pi^0$ LEFT-RIGHT ASYMMETRY PARAMETER**Measurements with an error $> 1.0 \times 10^{-2}$ have been omitted.

| <u>VALUE (units 10^{-2})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> |
|---|-------------|--------------------|-------------|
|---|-------------|--------------------|-------------|

 $0.09^{+0.11}_{-0.12}$ OUR AVERAGE

| | | | |
|----------------------------------|-------|---------------|------|
| $+0.09 \pm 0.10^{+0.09}_{-0.14}$ | 1.34M | AMBROSINO 08D | KLOE |
| 0.28 ± 0.26 | 165k | JANE 74 | OSPK |
| -0.05 ± 0.22 | 220k | LAYTER 72 | ASPK |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|---------------|-----|--------------------------|------|
| 1.5 ± 0.5 | 37k | ¹ GORMLEY 68C | ASPK |
|---------------|-----|--------------------------|------|

¹The GORMLEY 68C asymmetry is probably due to unmeasured (**E** × **B**) spark chamber effects. New experiments with (**E** × **B**) controls don't observe an asymmetry. **$\pi^+ \pi^- \pi^0$ SEXTANT ASYMMETRY PARAMETER**Measurements with an error $> 2.0 \times 10^{-2}$ have been omitted.

| <u>VALUE (units 10^{-2})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> |
|---|-------------|--------------------|-------------|
|---|-------------|--------------------|-------------|

 $0.12^{+0.10}_{-0.11}$ OUR AVERAGE

| | | | |
|----------------------------------|-------|---------------|------|
| $+0.08 \pm 0.10^{+0.08}_{-0.13}$ | 1.34M | AMBROSINO 08D | KLOE |
| 0.20 ± 0.25 | 165k | JANE 74 | OSPK |
| 0.10 ± 0.22 | 220k | LAYTER 72 | ASPK |
| 0.5 ± 0.5 | 37k | GORMLEY 68C | WIRE |

 $\pi^+ \pi^- \pi^0$ QUADRANT ASYMMETRY PARAMETER

| <u>VALUE (units 10^{-2})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> |
|---|-------------|--------------------|-------------|
|---|-------------|--------------------|-------------|

 -0.09 ± 0.09 OUR AVERAGE

| | | | |
|----------------------------------|-------|---------------|------|
| $-0.05 \pm 0.10^{+0.03}_{-0.05}$ | 1.34M | AMBROSINO 08D | KLOE |
| -0.30 ± 0.25 | 165k | JANE 74 | OSPK |
| -0.07 ± 0.22 | 220k | LAYTER 72 | ASPK |

 $\pi^+ \pi^- \gamma$ LEFT-RIGHT ASYMMETRY PARAMETERMeasurements with an error $> 2.0 \times 10^{-2}$ have been omitted.

| <u>VALUE (units 10^{-2})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> |
|---|-------------|--------------------|-------------|
|---|-------------|--------------------|-------------|

 0.9 ± 0.4 OUR AVERAGE

| | | | |
|-----------------|------|------------|------|
| 1.2 ± 0.6 | 35k | JANE 74B | OSPK |
| 0.5 ± 0.6 | 36k | THALER 72 | ASPK |
| 1.22 ± 1.56 | 7257 | GORMLEY 70 | ASPK |

 $\pi^+ \pi^- \gamma$ PARAMETER β (*D*-wave)Sensitive to a *D*-wave contribution: $dN/d\cos\theta = \sin^2\theta (1 + \beta \cos^2\theta)$.

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> |
|--------------|-------------|--------------------|-------------|
|--------------|-------------|--------------------|-------------|

 -0.02 ± 0.07 OUR AVERAGE Error includes scale factor of 1.3.

| | | | |
|--------------------|------|------------|------|
| 0.11 ± 0.11 | 35k | JANE 74B | OSPK |
| -0.060 ± 0.065 | 7250 | GORMLEY 70 | WIRE |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|-----------------|--|------------------------|------|
| 0.12 ± 0.06 | | ¹ THALER 72 | ASPK |
|-----------------|--|------------------------|------|

¹The authors don't believe this indicates D -wave because the dependence of β on the γ energy is inconsistent with the theoretical prediction. A $\cos^2\theta$ dependence can also come from P - and F -wave interference.

η CP-NONCONSERVING DECAY PARAMETER

$\pi^+\pi^-\ e^+e^-$ DECAY-PLANE ASYMMETRY PARAMETER A_ϕ

In the η rest frame, the total momentum of the e^+e^- pair is equal and opposite to that of the $\pi^+\pi^-$ pair. Let \hat{z} be the unit vector along the momentum of the e^+e^- pair; let \hat{n}_{ee} and $\hat{n}_{\pi\pi}$ be the unit vectors normal to the e^+e^- and $\pi^+\pi^-$ planes; and let ϕ be the angle between the two normals. Then

$$\sin\phi \cos\phi = [(\hat{n}_{ee} \times \hat{n}_{\pi\pi}) \cdot \hat{z}] (\hat{n}_{ee} \cdot \hat{n}_{\pi\pi}),$$

and

$$A_\phi \equiv \frac{N_{\sin\phi \cos\phi > 0} - N_{\sin\phi \cos\phi < 0}}{N_{\sin\phi \cos\phi > 0} + N_{\sin\phi \cos\phi < 0}}.$$

| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|---------------|-------------|----------|--|
| $-0.6 \pm 2.5 \pm 1.8$ | 1555 ± 52 | AMBROSINO | 09B KLOE | $e^+e^- \rightarrow \phi \rightarrow \eta\gamma$ |

ENERGY DEPENDENCE OF $\eta \rightarrow 3\pi$ DALITZ PLOTS

PARAMETERS FOR $\eta \rightarrow \pi^+\pi^-\pi^0$

See the "Note on η Decay Parameters," page 1454, in our 1994 edition (Physical Review **D50** 1173 (1994)). The following experiments fit to one or more of the coefficients a, b, c, d, e, f or g for $|\text{matrix element}|^2 = 1 + ay + by^2 + cx + dx^2 + exy + fy^3 + gx^2y$.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|----------|--|
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| 4.7M | 1 | ANASTASI | 16A KLOE | $e^+e^- \rightarrow \phi \rightarrow \eta\gamma$ |
| 79k | | ABLIKIM | 15G BES3 | $e^+e^- \rightarrow J/\psi \rightarrow \gamma\eta$ |
| 174k | | ADLARSON | 14A WASA | $pd \rightarrow \eta\ ^3\text{He}$ |
| 1.34M | | AMBROSINO | 08D KLOE | |
| 3230 | 2 | ABELE | 98D CBAR | $\bar{p}p \rightarrow \pi^0\pi^0\eta$ at rest |
| 1077 | 3 | AMSLER | 95 CBAR | $\bar{p}p \rightarrow \pi^+\pi^-\eta$ at rest |
| 81k | | LAYTER | 73 ASPK | |
| 220k | | LAYTER | 72 ASPK | |
| 1138 | | CARPENTER | 70 HBC | |
| 349 | | DANBURG | 70 DBC | |
| 7250 | | GORMLEY | 70 WIRE | |
| 526 | | BAGLIN | 69 HLBC | |
| 7170 | | CNOPS | 68 OSPK | |
| 37k | | GORMLEY | 68C WIRE | |
| 1300 | | CLPWY | 66 HBC | |
| 705 | | LARRIBE | 66 HBC | |

¹ANASTASI 16A measure the Dalitz parameters a, b, d, f , and g . This is the first measurement of g .

²ABELE 98D obtains $a = -1.22 \pm 0.07$ and $b = 0.22 \pm 0.11$ when c (or d) is fixed at 0.06.

³AMSLER 95 fits to $(1+ay+by^2)$ and obtains $a = -0.94 \pm 0.15$ and $b = 0.11 \pm 0.27$.

α PARAMETER FOR $\eta \rightarrow 3\pi^0$

See the "Note on η Decay Parameters" in our 1994 edition, Phys. Rev. **D50**, 1 August 1994, Part I, p. 1454. The value here is of α in $|\text{matrix element}|^2 = 1 + 2\alpha z$.

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|--------------------|-------------------------------------|----------|--|
| -0.0288 ± 0.0012 | OUR AVERAGE | Error includes scale factor of 1.1. | | |
| $-0.0265 \pm 0.0010 \pm 0.0009$ | 7M | PRAKHOV | 18 CRYB | $\gamma p \rightarrow p\eta$ |
| $-0.055 \pm 0.014 \pm 0.004$ | 33k | ABLIKIM | 15G BES3 | $e^+e^- \rightarrow J/\psi \rightarrow \gamma\eta$ |
| $-0.0301 \pm 0.0035^{+0.0022}_{-0.0035}$ | 512k | AMBROSINO | 10A KLOE | $e^+e^- \rightarrow \phi \rightarrow \eta\gamma$ |
| $-0.027 \pm 0.008 \pm 0.005$ | 120k | ¹ ADOLPH | 09 WASA | $pp \rightarrow pp\eta$ |
| $-0.0322 \pm 0.0012 \pm 0.0022$ | 3M | ² PRAKHOV | 09 CRYB | $\gamma p \rightarrow p\eta$ |
| $-0.032 \pm 0.002 \pm 0.002$ | 1.8M | ² UNVERZAGT | 09 CRYB | $\gamma p \rightarrow p\eta$ |
| $-0.026 \pm 0.010 \pm 0.010$ | 75k | BASHKANOV | 07 WASA | $pp \rightarrow pp\eta$ |
| $-0.010 \pm 0.021 \pm 0.010$ | 12k | ACHASOV | 01C SND | $e^+e^- \rightarrow \phi \rightarrow \eta\gamma$ |
| -0.031 ± 0.004 | 1M | TIPPENS | 01 CRYB | $\pi^- p \rightarrow n\eta$, 720 MeV |
| $-0.052 \pm 0.017 \pm 0.010$ | 98k | ABELE | 98C CBAR | $\bar{p}p \rightarrow 5\pi^0$ |
| -0.022 ± 0.023 | 50k | ALDE | 84 GAM2 | |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|--------------------------------------|-------|------------------------|----------|--|
| $-0.038 \pm 0.003^{+0.012}_{-0.008}$ | 1.34M | ³ AMBROSINO | 08D KLOE | |
| -0.32 ± 0.37 | 192 | BAGLIN | 70 HLBC | |

¹This ADOLPH 09 result is independent of the BASHKANOV 07 result.

²The PRAKHOV 09 and UNVERZAGT 09 results are independent.

³This AMBROSINO 08D value is an indirect result using $\eta \rightarrow \pi^+\pi^0\pi^-$ events and a rescattering matrix that mixes isospin decay amplitudes.

PARAMETER Λ IN $\eta \rightarrow \ell^+\ell^-\gamma$ DECAY

In the pole approximation the electromagnetic transition form factor for a resonance of mass M is given by the expression:

$$|F|^2 = (1 - M_{\ell\ell}^2/\Lambda^2)^{-2},$$

where for the parameter Λ vector dominance predicts $\Lambda \approx 0.770$ GeV.

| VALUE (GeV/c ²) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------------------------------------|--------------------|-----------------------|----------|--|
| 0.716 ± 0.011 | OUR AVERAGE | | | |
| 0.712 ± 0.020 | | ¹ ADLARSON | 17B A2MM | $\gamma p \rightarrow \eta p$ |
| $0.7191 \pm 0.0125 \pm 0.0093$ | | ² ARNALDI | 16 NA60 | 400 GeV p -A collisions |
| $0.716 \pm 0.031 \pm 0.009$ | | ³ ARNALDI | 09 NA60 | 158A In-In collisions |
| 0.72 ± 0.09 | 600 | DZHELYADIN | 80 SPEC | $\pi^- p \rightarrow \eta n$, $\eta \rightarrow \gamma\mu^+\mu^-$ |

¹ADLARSON 17B reports $\Lambda^{-2}(\eta \rightarrow \gamma e^+e^-) = 1.97 \pm 0.11$ (GeV/c²)⁻² which we converted to the quoted Λ value and uncertainty (total=statistical plus systematic).

²ARNALDI 16 reports $\Lambda^{-2}(\eta \rightarrow \gamma\mu^+\mu^-) = 1.934 \pm 0.067 \pm 0.050$ (GeV/c²)⁻² which we converted to the quoted Λ value.

³ARNALDI 09 reports $\Lambda^{-2}(\eta \rightarrow \gamma\mu^+\mu^-) = 1.95 \pm 0.17 \pm 0.05$ (GeV/c²)⁻² which we converted to the quoted Λ value.

η REFERENCES

| | | | | |
|------------|------|--|-------------------------------|-------------------------------|
| ABLIKIM | 21AM | PR D104 092004 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| BABUSCI | 20A | JHEP 2010 047 | D. Babusci <i>et al.</i> | (KLOE-2 Collab.) |
| ZHEVLAKOV | 19 | PR D99 031703 | A.S. Zhevlakov <i>et al.</i> | (TMSK, MAINZ, TUBIN+) |
| ACHASOV | 18B | PR D98 052007 | M.N. Achasov <i>et al.</i> | (SND Collab.) |
| ADLARSON | 18C | PL B784 378 | P. Adlarson <i>et al.</i> | (WASA-at-COSY Collab.) |
| PRAKHOV | 18 | PR C97 065203 | S. Prakhov <i>et al.</i> | (A2 Collab. at MAMI) |
| AAIJ | 17D | PL B764 233 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| ADLARSON | 17B | PR C95 035208 | P. Adlarson <i>et al.</i> | (A2 Collab. at MAMI) |
| ANASTASI | 16A | JHEP 1605 019 | A. Anastasi <i>et al.</i> | (KLOE-2 Collab.) |
| ARNALDI | 16 | PL B757 437 | R. Arnaldi <i>et al.</i> | (NA60 Collab.) |
| ABLIKIM | 15G | PR D92 012014 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ADLARSON | 14A | PR C90 045207 | P. Adlarson <i>et al.</i> | (WASA-at-COSY Collab.) |
| AGAKISHIEV | 14 | PL B731 265 | G. Agakishiev <i>et al.</i> | (HADES Collab.) |
| NEFKENS | 14 | PR C90 025206 | B.M.K. Nefkens <i>et al.</i> | (A2 Collab. at MAMI) |
| NIKOLAEV | 14 | EPJ A50 58 | A. Nikolaev <i>et al.</i> | (MAMI-B, MAINZ, BONN) |
| ABLIKIM | 13 | PR D87 012009 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 13G | PR D87 032006 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| BABUSCI | 13 | PL B718 910 | D. Babusci <i>et al.</i> | (KLOE/KLOE-2 Collab.) |
| BABUSCI | 13A | JHEP 1301 119 | D. Babusci <i>et al.</i> | (KLOE-2 Collab.) |
| AGAKISHIEV | 12A | EPJ A48 64 | G. Agakishiev <i>et al.</i> | (HADES Collab.) |
| GOSLAWSKI | 12 | PR D85 112011 | P. Goslawski <i>et al.</i> | (COSY-ANKE Collab.) |
| ABLIKIM | 11G | PR D84 032006 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| AMBROSINO | 11B | PL B702 324 | F. Ambrosino <i>et al.</i> | (KLOE Collab.) |
| BERGHAUSER | 11 | PL B701 562 | H. Berghausen <i>et al.</i> | (GIES, UCLA, GUTE) |
| AMBROSINO | 10A | PL B694 16 | F. Ambrosino <i>et al.</i> | (KLOE Collab.) |
| ADOLPH | 09 | PL B677 24 | C. Adolph <i>et al.</i> | (WASA-at-COSY Collab.) |
| AMBROSINO | 09B | PL B675 283 | F. Ambrosino <i>et al.</i> | (KLOE Collab.) |
| ARNALDI | 09 | PL B677 260 | R. Arnaldi <i>et al.</i> | (NA60 Collab.) |
| PRAKHOV | 09 | PR C79 035204 | S. Prakhov <i>et al.</i> | (MAMI-C Crystal Ball Collab.) |
| UNVERZAGT | 09 | EPJ A39 169 | M. Unverzagt <i>et al.</i> | (MAMI-B Crystal Ball Collab.) |
| AMBROSINO | 08D | JHEP 0805 006 | F. Ambrosino <i>et al.</i> | (DAPHNE KLOE Collab.) |
| BERLOWSKI | 08 | PR D77 032004 | M. Berlowski <i>et al.</i> | (CELSIUS/WASA Collab.) |
| PRAKHOV | 08 | PR C78 015206 | S. Prakhov <i>et al.</i> | (BNL Crystal Ball Collab.) |
| RODRIGUES | 08 | PRL 101 012301 | T.E. Rodrigues <i>et al.</i> | (USP, FESP, UNESP+) |
| AMBROSINO | 07B | JHEP 0712 073 | F. Ambrosino <i>et al.</i> | (KLOE Collab.) |
| BARGHOLTZ | 07 | PL B644 299 | Chr. Bargholtz <i>et al.</i> | (CELSIUS/WASA Collab.) |
| BASHKANOV | 07 | PR C76 048201 | M. Bashkanov <i>et al.</i> | (CELSIUS/WASA Collab.) |
| BLIK | 07 | PAN 70 693 | A.M. Blik <i>et al.</i> | (GAMS Collab.) |
| LOPEZ | 07 | Translated from YAF 70 724. PRL 99 122001 | A. Lopez <i>et al.</i> | (CLEO Collab.) |
| MILLER | 07 | PRL 99 122002 | D.H. Miller <i>et al.</i> | (CLEO Collab.) |
| ABLIKIM | 06E | PR D73 052008 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ABLIKIM | 06Q | PRL 97 202002 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| ACHASOV | 06A | PR D74 014016 | M.N. Achasov <i>et al.</i> | (SND Collab.) |
| ABDEL-BARY | 05 | PL B619 281 | M. Abdel-Bary <i>et al.</i> | (GEM Collab.) |
| AKHMETSHIN | 05 | PL B605 26 | R.R. Akhmetshin <i>et al.</i> | (Novosibirsk CMD-2 Collab.) |
| AMBROSINO | 05A | PL B606 276 | F. Ambrosino <i>et al.</i> | (KLOE Collab.) |
| NEFKENS | 05 | PRL 94 041601 | B.M.K. Nefkens <i>et al.</i> | (BNL Crystal Ball Collab.) |
| NEFKENS | 05A | PR C72 035212 | B.M.K. Nefkens <i>et al.</i> | (BNL Crystal Ball Collab.) |
| PRAKHOV | 05 | PR C72 025201 | S. Prakhov <i>et al.</i> | (BNL Crystal Ball Collab.) |
| ALOISIO | 04 | PL B591 49 | A. Aloisio <i>et al.</i> | (KLOE Collab.) |
| KNECHT | 04 | PL B589 14 | N. Knecht <i>et al.</i> | |
| LAI | 02 | PL B533 196 | A. Lai <i>et al.</i> | (CERN NA48 Collab.) |
| NEFKENS | 02 | PS T99 114 | B.M.K. Nefkens, J.W. Price | (UCLA) |
| ACHASOV | 01B | PL B504 275 | M.N. Achasov <i>et al.</i> | (Novosibirsk SND Collab.) |
| ACHASOV | 01C | JETPL 73 451 | M.N. Achasov <i>et al.</i> | (Novosibirsk SND Collab.) |
| ACHASOV | 01D | Translated from ZETFP 73 511. NP B600 3 | M.N. Achasov <i>et al.</i> | (Novosibirsk SND Collab.) |
| AKHMETSHIN | 01 | PL B501 191 | R.R. Akhmetshin <i>et al.</i> | (Novosibirsk CMD-2 Collab.) |
| AKHMETSHIN | 01B | PL B509 217 | R.R. Akhmetshin <i>et al.</i> | (Novosibirsk CMD-2 Collab.) |
| TIPPENS | 01 | PRL 87 192001 | W.B. Tippens <i>et al.</i> | (BNL Crystal Ball Collab.) |
| ACHASOV | 00 | EPJ C12 25 | M.N. Achasov <i>et al.</i> | (Novosibirsk SND Collab.) |
| ACHASOV | 00B | JETP 90 17 | M.N. Achasov <i>et al.</i> | (Novosibirsk SND Collab.) |
| ACHASOV | 00D | Translated from ZETF 117 22. JETPL 72 282 | M.N. Achasov <i>et al.</i> | (Novosibirsk SND Collab.) |
| | | Translated from ZETFP 72 411. | | |

| | | | | |
|---------------|-----|-----------------------------------|---|-----------------------------|
| PRAKHOV | 00 | PRL 84 4802 | S. Prakhov <i>et al.</i> | (BNL Crystal Ball Collab.) |
| AKHMETSHIN | 99B | PL B462 371 | R.R. Akhmetshin <i>et al.</i> | (Novosibirsk CMD-2 Collab.) |
| AKHMETSHIN | 99C | PL B462 380 | R.R. Akhmetshin <i>et al.</i> | (Novosibirsk CMD-2 Collab.) |
| AKHMETSHIN | 99F | PL B460 242 | R.R. Akhmetshin <i>et al.</i> | (Novosibirsk CMD-2 Collab.) |
| ABELE | 98C | PL B417 193 | A. Abele <i>et al.</i> | (Crystal Barrel Collab.) |
| ABELE | 98D | PL B417 197 | A. Abele <i>et al.</i> | (Crystal Barrel Collab.) |
| ACHASOV | 98 | PL B425 388 | M.N. Achasov <i>et al.</i> | (Novosibirsk SND Collab.) |
| AKHMETSHIN | 97C | PL B415 452 | R.R. Akhmetshin <i>et al.</i> | (Novosibirsk CMD-2 Collab.) |
| BROWDER | 97B | PR D56 5359 | T.E. Browder <i>et al.</i> | (CLEO Collab.) |
| ABEGG | 96 | PR D53 11 | R. Abegg <i>et al.</i> | (Saturne SPES2 Collab.) |
| WHITE | 96 | PR D53 6658 | D.B. White <i>et al.</i> | (Saturne SPES2 Collab.) |
| AMSLER | 95 | PL B346 203 | C. Amsler <i>et al.</i> | (Crystal Barrel Collab.) |
| KRUSCHE | 95D | ZPHY A351 237 | B. Krusche <i>et al.</i> | (TAPS + A2 Collab.) |
| ABEGG | 94 | PR D50 92 | R. Abegg <i>et al.</i> | (Saturne SPES2 Collab.) |
| PDG | 94 | PR D50 1173 | L. Montanet <i>et al.</i> | (CERN, LBL, BOST+) |
| AMSLER | 93 | ZPHY C58 175 | C. Amsler <i>et al.</i> | (Crystal Barrel Collab.) |
| KESSLER | 93 | PRL 70 892 | R.S. Kessler <i>et al.</i> | (Saturne SPES2 Collab.) |
| PLOUIN | 92 | PL B276 526 | F. Plouin <i>et al.</i> | (Saturne SPES4 Collab.) |
| BARU | 90 | ZPHY C48 581 | S.E. Baru <i>et al.</i> | (MD-1 Collab.) |
| ROE | 90 | PR D41 17 | N.A. Roe <i>et al.</i> | (ASP Collab.) |
| WILLIAMS | 88 | PR D38 1365 | D.A. Williams <i>et al.</i> | (Crystal Ball Collab.) |
| AIHARA | 86 | PR D33 844 | H. Aihara <i>et al.</i> | (TPC-2 γ Collab.) |
| BARTEL | 85E | PL 160B 421 | W. Bartel <i>et al.</i> | (JADE Collab.) |
| LANDSBERG | 85 | PRPL 128 301 | L.G. Landsberg | (SERP) |
| ALDE | 84 | ZPHY C25 225 | D.M. Alde <i>et al.</i> | (SERP, BELG, LAPP) |
| Also | | SJNP 40 918 | D.M. Alde <i>et al.</i> | (SERP, BELG, LAPP) |
| | | Translated from YAF 40 1447. | | |
| WEINSTEIN | 83 | PR D28 2896 | A.J. Weinstein <i>et al.</i> | (Crystal Ball Collab.) |
| BINON | 82 | SJNP 36 391 | F.G. Binon <i>et al.</i> | (SERP, BELG, LAPP+) |
| | | Translated from YAF 36 670. | | |
| Also | | NC 71A 497 | F.G. Binon <i>et al.</i> | (SERP, BELG, LAPP+) |
| DAVYDOV | 81 | LNC 32 45 | V.A. Davydov <i>et al.</i> | (SERP, BELG, LAPP+) |
| Also | | SJNP 33 825 | V.A. Davydov <i>et al.</i> | (SERP, BELG, LAPP+) |
| | | Translated from YAF 33 1534. | | |
| DZHELADIN | 81 | PL 105B 239 | R.I. Dzhelyadin <i>et al.</i> | (SERP) |
| Also | | SJNP 33 822 | R.I. Dzhelyadin <i>et al.</i> | (SERP) |
| | | Translated from YAF 33 1529. | | |
| ABROSIMOV | 80 | SJNP 31 195 | A.T. Abrosimov <i>et al.</i> | (JINR) |
| | | Translated from YAF 31 371. | | |
| DZHELADIN | 80 | PL 94B 548 | R.I. Dzhelyadin <i>et al.</i> | (SERP) |
| Also | | SJNP 32 516 | R.I. Dzhelyadin <i>et al.</i> | (SERP) |
| | | Translated from YAF 32 998. | | |
| DZHELADIN | 80B | PL 97B 471 | R.I. Dzhelyadin <i>et al.</i> | (SERP) |
| Also | | SJNP 32 518 | R.I. Dzhelyadin <i>et al.</i> | (SERP) |
| | | Translated from YAF 32 1002. | | |
| BUSHNIN | 78 | PL 79B 147 | Y.B. Bushnin <i>et al.</i> | (SERP) |
| Also | | SJNP 28 775 | Y.B. Bushnin <i>et al.</i> | (SERP) |
| | | Translated from YAF 28 1507. | | |
| MARTYNOV | 76 | SJNP 23 48 | A.S. Martynov <i>et al.</i> | (JINR) |
| | | Translated from YAF 23 93. | | |
| JANE | 75 | PL 59B 99 | M.R. Jane <i>et al.</i> | (RHEL, LOWC) |
| JANE | 75B | PL 59B 103 | M.R. Jane <i>et al.</i> | (RHEL, LOWC) |
| Also | | PL 73B 503 | M.R. Jane | |
| | | Erratum in private communication. | | |
| BROWMAN | 74B | PRL 32 1067 | A. Browman <i>et al.</i> | (CORN, BING) |
| DAVIES | 74 | NC 24A 324 | J.D. Davies, J.G. Guy, R.K.P. Zia | (BIRM, RHEL+) |
| DUANE | 74 | PRL 32 425 | A. Duane <i>et al.</i> | (LOIC, SHMP) |
| JANE | 74 | PL 48B 260 | M.R. Jane <i>et al.</i> | (RHEL, LOWC, SUSS) |
| JANE | 74B | PL 48B 265 | M.R. Jane <i>et al.</i> | (RHEL, LOWC, SUSS) |
| KENDALL | 74 | NC 21A 387 | B.N. Kendall <i>et al.</i> | (BROW, BARI, MIT) |
| LAYTER | 73 | PR D7 2565 | J.G. Layter <i>et al.</i> | (COLU) |
| THALER | 73 | PR D7 2569 | J.J. Thaler <i>et al.</i> | (COLU) |
| AGUILAR... | 72B | PR D6 29 | M. Aguilar-Benitez <i>et al.</i> | (BNL) |
| BLOODWORTH... | 72B | NP B39 525 | I.J. Bloodworth <i>et al.</i> | (TNTO) |
| LAYTER | 72 | PRL 29 316 | J.G. Layter <i>et al.</i> | (COLU) |
| THALER | 72 | PRL 29 313 | J.J. Thaler <i>et al.</i> | (COLU) |
| BASILE | 71D | NC 3A 796 | M. Basile <i>et al.</i> | (CERN, BGNA, STRB) |
| STRUGALSKI | 71 | NP B27 429 | Z.S. Strugalski <i>et al.</i> | (JINR) |
| BAGLIN | 70 | NP B22 66 | C. Baglin <i>et al.</i> | (EPOL, MADR, STRB) |
| BUTTRAM | 70 | PRL 25 1358 | M.T. Buttram, M.N. Kreisler, R.E. Mischke | (PRIN) |
| CARPENTER | 70 | PR D1 1303 | D.W. Carpenter <i>et al.</i> | (DUKE) |
| COX | 70B | PRL 24 534 | B. Cox, L. Fortney, J.P. Golson | (DUKE) |

| | | | | |
|------------|-----|------------------|---|-------------------------------|
| DANBURG | 70 | PR D2 2564 | J.S. Danburg <i>et al.</i> | (LRL) |
| DEVONS | 70 | PR D1 1936 | S. Devons <i>et al.</i> | (COLU, SYRA) |
| GORMLEY | 70 | PR D2 501 | M. Gormley <i>et al.</i> | (COLU, BNL) |
| Also | | Thesis Nevis 181 | M. Gormley | (COLU) |
| BAGLIN | 69 | PL 29B 445 | C. Baglin <i>et al.</i> | (EPOL, UCB, MADR, STRB) |
| Also | | NP B22 66 | C. Baglin <i>et al.</i> | (EPOL, MADR, STRB) |
| HYAMS | 69 | PL 29B 128 | B.D. Hyams <i>et al.</i> | (CERN, MPIM) |
| ARNOLD | 68 | PL 27B 466 | R.G. Arnold <i>et al.</i> | (STRB, MADR, EPOL+) |
| BAZIN | 68 | PRL 20 895 | M.J. Bazin <i>et al.</i> | (PRIN, UKI) |
| BULLOCK | 68 | PL 27B 402 | F.W. Bullock <i>et al.</i> | (LOUC) |
| CNOPS | 68 | PRL 21 1609 | A.M. Cnops <i>et al.</i> | (BNL, ORNL, UCND+) |
| GORMLEY | 68C | PRL 21 402 | M. Gormley <i>et al.</i> | (COLU, BNL) |
| WEHMANN | 68 | PRL 20 748 | A.W. Wehmann <i>et al.</i> | (HARV, CASE, SLAC+) |
| BAGLIN | 67 | PL 24B 637 | C. Baglin <i>et al.</i> | (EPOL, UCB) |
| BAGLIN | 67B | BAPS 12 567 | C. Baglin <i>et al.</i> | (EPOL, UCB) |
| BALTAY | 67B | PRL 19 1498 | C. Baltay <i>et al.</i> | (COLU, STON) |
| BALTAY | 67D | PRL 19 1495 | C. Baltay <i>et al.</i> | (COLU, BRAN) |
| BEMPORAD | 67 | PL 25B 380 | C. Bemporad <i>et al.</i> | (PISA, BONN) |
| Also | | Private Comm. | I. Ion | |
| BILLING | 67 | PL 25B 435 | K.D. Billing <i>et al.</i> | (LOUC, OXF) |
| BUNIATOV | 67 | PL 25B 560 | S.A. Bunyatov <i>et al.</i> | (CERN, KARL) |
| CENCE | 67 | PRL 19 1393 | R.J. Cence <i>et al.</i> | (HAWA, LRL) |
| ESTEN | 67 | PL 24B 115 | M.J. Esten <i>et al.</i> | (LOUC, OXF) |
| FELDMAN | 67 | PRL 18 868 | M. Feldman <i>et al.</i> | (PENN) |
| FLATTE | 67 | PRL 18 976 | S.M. Flatte | (LRL) |
| FLATTE | 67B | PR 163 1441 | S.M. Flatte, C.G. Wohl | (LRL) |
| LITCHFIELD | 67 | PL 24B 486 | P.J. Litchfield <i>et al.</i> | (RHEL, SACL) |
| PRICE | 67 | PRL 18 1207 | L.R. Price, F.S. Crawford | (LRL) |
| ALFF-... | 66 | PR 145 1072 | C. Alff-Steinberger <i>et al.</i> | (COLU, RUTG) |
| CLPWY | 66 | PR 149 1044 | C. Baltay | (SCUC, LRL, PURD, WISC, YALE) |
| CRAWFORD | 66 | PRL 16 333 | F.S. Crawford, L.R. Price | (LRL) |
| DIGIUGNO | 66 | PRL 16 767 | G. di Giugno <i>et al.</i> | (NAPL, TRST, FRAS) |
| GRUNHAUS | 66 | Thesis | J. Grunhaus | (COLU) |
| JAMES | 66 | PR 142 896 | F.E. James, H.L. Kraybill | (YALE, BNL) |
| JONES | 66 | PL 23 597 | W.G. Jones <i>et al.</i> | (LOIC, RHEL) |
| LARRIBE | 66 | PL 23 600 | A. Larribe <i>et al.</i> | (SACL, RHEL) |
| FOSTER | 65 | PR 138 B652 | M. Foster <i>et al.</i> | (WISC, PURD) |
| FOSTER | 65B | Athens Conf. | M. Foster, M. Good, M. Meer | (WISC) |
| FOSTER | 65C | Thesis | M. Foster | (WISC) |
| PRICE | 65 | PRL 15 123 | L.R. Price, F.S. Crawford | (LRL) |
| RITTENBERG | 65 | PRL 15 556 | A. Rittenberg, G.R. Kalbfleisch | (LRL, BNL) |
| FOELSCH | 64 | PR 134 B1138 | H.W.J. Foelsche, H.L. Kraybill | (YALE) |
| KRAEMER | 64 | PR 136 B496 | R.W. Kraemer <i>et al.</i> | (JHU, NWES, WOOD) |
| PAULI | 64 | PL 13 351 | E. Pauli, A. Muller | (SACL) |
| BACCI | 63 | PRL 11 37 | C. Bacci <i>et al.</i> | (ROMA, FRAS) |
| CRAWFORD | 63 | PRL 10 546 | F.S.Jr. Crawford, L.J. Lloyd, E.C. Fowler | (LRL+) |
| Also | | PRL 16 907 | F.S. Crawford, L.J. Lloyd, E.C. Fowler | (LRL+) |
| ALFF-... | 62 | PRL 9 322 | C. Alff-Steinberger <i>et al.</i> | (COLU, RUTG) |
| BASTIEN | 62 | PRL 8 114 | P.L. Bastien <i>et al.</i> | (LRL) |
| PICKUP | 62 | PRL 8 329 | E. Pickup, D.K. Robinson, E.O. Salant | (CNRC+) |