

**$f_2(1565)$** 

$$I^G(J^{PC}) = 0^+(2^{++})$$


## OMITTED FROM SUMMARY TABLE

Seen mostly in antinucleon-nucleon annihilation. Needs confirmation in other channels. See the review on "Spectroscopy of Light Meson Resonances."

 **$f_2(1565)$  MASS**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b><math>1542 \pm 19</math> OUR AVERAGE</b>	Error includes scale factor of 2.2. See the ideogram below.		
$1552 \pm 13$	<sup>1</sup> AMSLER	02	CBAR $0.9 \bar{p}p \rightarrow \pi^0 \eta \eta, \pi^0 \pi^0 \pi^0$
$1575 \pm 18$	<sup>1</sup> BERTIN	98	OBLX $0.05\text{--}0.405 \bar{n}p \rightarrow \pi^+ \pi^+ \pi^-$
$1507 \pm 15$	<sup>1</sup> BERTIN	97C	OBLX $0.0 \bar{p}p \rightarrow \pi^+ \pi^- \pi^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$1560 \pm 15$	<sup>2</sup> ANISOVICH	09	RVUE $0.0 \bar{p}p, \pi N$
$1590 \pm 10$	<sup>3,4</sup> AMELIN	06	VES $36 \pi^- p \rightarrow \omega \omega n$
$1550 \pm 10 \pm 20$	<sup>4</sup> AMELIN	00	VES $37 \pi^- p \rightarrow \eta \pi^+ \pi^- n$
$1598 \pm 11 \pm 9$	BAKER	99B	SPEC $0 \bar{p}p \rightarrow \omega \omega \pi^0$
$1534 \pm 20$	<sup>5</sup> ABELE	96C	RVUE Compilation
$\sim 1552$	<sup>6</sup> AMSLER	95D	CBAR $0.0 \bar{p}p \rightarrow \pi^0 \pi^0 \pi^0, \pi^0 \eta \eta, \pi^0 \pi^0 \eta$
$1598 \pm 72$	BALOSHIN	95	SPEC $40 \pi^- C \rightarrow K_S^0 K_S^0 X$
$1566^{+80}_{-50}$	<sup>7</sup> ANISOVICH	94	CBAR $0.0 \bar{p}p \rightarrow 3\pi^0, \eta \eta \pi^0$
$1502 \pm 9$	ADAMO	93	OBLX $\bar{n}p \rightarrow \pi^+ \pi^+ \pi^-$
$1488 \pm 10$	<sup>8</sup> ARMSTRONG	93C	E760 $\bar{p}p \rightarrow \pi^0 \eta \eta \rightarrow 6\gamma$
$1508 \pm 10$	<sup>8</sup> ARMSTRONG	93D	E760 $\bar{p}p \rightarrow 3\pi^0 \rightarrow 6\gamma$
$1525 \pm 10$	<sup>8</sup> ARMSTRONG	93D	E760 $\bar{p}p \rightarrow \eta \pi^0 \pi^0 \rightarrow 6\gamma$
$\sim 1504$	<sup>9</sup> WEIDENAUER	93	ASTE $0.0 \bar{p}N \rightarrow 3\pi^- 2\pi^+$
$1540 \pm 15$	<sup>8</sup> ADAMO	92	OBLX $\bar{n}p \rightarrow \pi^+ \pi^+ \pi^-$
$1515 \pm 10$	<sup>10</sup> AKER	91	CBAR $0.0 \bar{p}p \rightarrow 3\pi^0$
$1565 \pm 20$	MAY	90	ASTE $0.0 \bar{p}p \rightarrow \pi^+ \pi^- \pi^0$
$1477 \pm 5$	BRIDGES	86C	DBC $0.0 \bar{p}N \rightarrow 3\pi^- 2\pi^+$

<sup>1</sup> T-matrix pole.<sup>2</sup> On sheet II in a two-pole solution.<sup>3</sup> Supersedes the  $\omega\omega$  state of BELADIDZE 92B earlier assigned to the  $f_2(1640)$ .<sup>4</sup> Breit-Wigner width.<sup>5</sup> T-matrix pole, large coupling to  $\rho\rho$  and  $\omega\omega$ , could be  $f_2(1640)$ .<sup>6</sup> Coupled-channel analysis of AMSLER 95B, AMSLER 95C, and AMSLER 94D.<sup>7</sup> From a simultaneous analysis of the annihilations  $\bar{p}p \rightarrow 3\pi^0, \pi^0 \eta \eta$  including AKER 91 data.<sup>8</sup>  $J^P$  not determined, could be partly  $f_0(1500)$ .<sup>9</sup>  $J^P$  not determined.<sup>10</sup> Superseded by AMSLER 95B.



### $f_2(1565)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>122± 13 OUR AVERAGE</b>			
113± 23	11 AMSLER	02 CBAR	0.9 $\bar{p}p \rightarrow \pi^0 \eta \eta, \pi^0 \pi^0 \pi^0$
119± 24	11 BERTIN	98 OBLX	0.05–0.405 $\bar{n}p \rightarrow \pi^+ \pi^+ \pi^-$
130± 20	11 BERTIN	97C OBLX	0.0 $\bar{p}p \rightarrow \pi^+ \pi^- \pi^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
280± 40	12 ANISOVICH	09 RVUE	0.0 $\bar{p}p, \pi N$
140± 11	13,14 AMELIN	06 VES	36 $\pi^- p \rightarrow \omega \omega n$
130± 20±40	14 AMELIN	00 VES	37 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$
180± 60	15 ABELE	96C RVUE	Compilation
~ 142	16 AMSLER	95D CBAR	0.0 $\bar{p}p \rightarrow \pi^0 \pi^0 \pi^0, \pi^0 \eta \eta, \pi^0 \pi^0 \eta$
263±101	BALOSHIN	95 SPEC	40 $\pi^- C \rightarrow K_S^0 K_S^0 X$
166 <sup>+</sup> <sub>-</sub> 80 20	17 ANISOVICH	94 CBAR	0.0 $\bar{p}p \rightarrow 3\pi^0, \eta \eta \pi^0$
130± 10	18 ADAMO	93 OBLX	$\bar{n}p \rightarrow \pi^+ \pi^+ \pi^-$
148± 27	19 ARMSTRONG	93C E760	$\bar{p}p \rightarrow \pi^0 \eta \eta \rightarrow 6\gamma$
103± 15	19 ARMSTRONG	93D E760	$\bar{p}p \rightarrow 3\pi^0 \rightarrow 6\gamma$
111± 10	19 ARMSTRONG	93D E760	$\bar{p}p \rightarrow \eta \pi^0 \pi^0 \rightarrow 6\gamma$
~ 206	20 WEIDENAUER	93 ASTE	0.0 $\bar{p}N \rightarrow 3\pi^- 2\pi^+$
132± 37	19 ADAMO	92 OBLX	$\bar{n}p \rightarrow \pi^+ \pi^+ \pi^-$
120± 10	21 AKER	91 CBAR	0.0 $\bar{p}p \rightarrow 3\pi^0$
170± 40	MAY	90 ASTE	0.0 $\bar{p}p \rightarrow \pi^+ \pi^- \pi^0$
116± 9	BRIDGES	86C DBC	0.0 $\bar{p}N \rightarrow 3\pi^- 2\pi^+$

- 11 T-matrix pole.  
 12 On sheet II in a two-pole solution.  
 13 Supersedes the  $\omega\omega$  state of BELADIDZE 92B earlier assigned to the  $f_2(1640)$ .  
 14 Breit-Wigner width.  
 15 T-matrix pole, large coupling to  $\rho\rho$  and  $\omega\omega$ , could be  $f_2(1640)$ .  
 16 Coupled-channel analysis of AMSLER 95B, AMSLER 95C, and AMSLER 94D.  
 17 From a simultaneous analysis of the annihilations  $\bar{p}p \rightarrow 3\pi^0, \pi^0\eta\eta$  including AKER 91 data.  
 18 Supersedes ADAMO 92.  
 19  $J^P$  not determined, could be partly  $f_0(1500)$ .  
 20  $J^P$  not determined.  
 21 Superseded by AMSLER 95B.

## $f_2(1565)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\pi\pi$	seen
$\Gamma_2$ $\pi^+\pi^-$	seen
$\Gamma_3$ $\pi^0\pi^0$	seen
$\Gamma_4$ $\rho^0\rho^0$	seen
$\Gamma_5$ $2\pi^+2\pi^-$	seen
$\Gamma_6$ $\eta\eta$	seen
$\Gamma_7$ $\omega\omega$	seen
$\Gamma_8$ $K\bar{K}$	seen
$\Gamma_9$ $\gamma\gamma$	seen

## $f_2(1565)$ PARTIAL WIDTHS

$\Gamma(\eta\eta)$					$\Gamma_6$
VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
••• We do not use the following data for averages, fits, limits, etc. •••					
$1.2 \pm 0.3$	870	<sup>22</sup> SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$	
$\Gamma(K\bar{K})$					$\Gamma_8$
VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
••• We do not use the following data for averages, fits, limits, etc. •••					
$2.0 \pm 1.0$	870	<sup>22</sup> SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$	
$\Gamma(\gamma\gamma)$					$\Gamma_9$
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT	
••• We do not use the following data for averages, fits, limits, etc. •••					
$0.70 \pm 0.14$	870	<sup>22</sup> SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$	
<sup>22</sup> From analysis of L3 data at 91 and 183–209 GeV, using $f_2(1565)$ mass of 1570 MeV, width of 160 MeV, $\Gamma(\pi\pi) = 25$ MeV, and SU(3) relations.					

**$f_2(1565)$  BRANCHING RATIOS** **$\Gamma(\pi\pi)/\Gamma_{\text{total}}$   $\Gamma_1/\Gamma$** 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
seen	BAKER	99B SPEC	$0 \bar{p}p \rightarrow \omega\omega\pi^0$

 **$\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_2/\Gamma$** 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
seen	BERTIN	98 OBLX	$0.05-0.405 \bar{p}p \rightarrow \pi^+\pi^+\pi^-$
not seen	<sup>23</sup> ANISOVICH	94B RVUE	$\bar{p}p \rightarrow \pi^+\pi^-\pi^0$
seen	MAY	89 ASTE	$\bar{p}p \rightarrow \pi^+\pi^-\pi^0$
<sup>23</sup> ANISOVICH 94B is from a reanalysis of MAY 90.			

 **$\Gamma(\pi^0\pi^0)/\Gamma_{\text{total}}$   $\Gamma_3/\Gamma$** 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
seen	AMSLER	95B CBAR	$0.0 \bar{p}p \rightarrow 3\pi^0$

 **$\Gamma(\pi^+\pi^-)/\Gamma(\rho^0\rho^0)$   $\Gamma_2/\Gamma_4$** 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$0.042 \pm 0.013$	BRIDGES	86B DBC	$\bar{p}N \rightarrow 3\pi^- 2\pi^+$

 **$\Gamma(\eta\eta)/\Gamma(\pi^0\pi^0)$   $\Gamma_6/\Gamma_3$** 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$0.024 \pm 0.005 \pm 0.012$	<sup>24</sup> ARMSTRONG	93C E760	$\bar{p}p \rightarrow \pi^0\eta\eta \rightarrow 6\gamma$
<sup>24</sup> $J^P$ not determined, could be partly $f_0(1500)$ .			

 **$\Gamma(\omega\omega)/\Gamma_{\text{total}}$   $\Gamma_7/\Gamma$** 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
seen	BAKER	99B SPEC	$0 \bar{p}p \rightarrow \omega\omega\pi^0$

 **$f_2(1565)$  REFERENCES**

ANISOVICH	09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev	
AMELIN	06	PAN 69 690	D.V. Amelin <i>et al.</i>	(VES Collab.)
		Translated from YAF 69 715.		
SCHEGELSKY	06A	EPJ A27 207	V.A. Schegelsky <i>et al.</i>	
AMSLER	02	EPJ C23 29	C. Amsler <i>et al.</i>	
AMELIN	00	NP A668 83	D. Amelin <i>et al.</i>	(VES Collab.)
BAKER	99B	PL B467 147	C.A. Baker <i>et al.</i>	
BERTIN	98	PR D57 55	A. Bertin <i>et al.</i>	(OBELIX Collab.)
BERTIN	97C	PL B408 476	A. Bertin <i>et al.</i>	(OBELIX Collab.)
ABELE	96C	NP A609 562	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
AMSLER	95B	PL B342 433	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
AMSLER	95C	PL B353 571	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
AMSLER	95D	PL B355 425	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
BALOSHIN	95	PAN 58 46	O.N. Baloshin <i>et al.</i>	(ITEP)
		Translated from YAF 58 50.		

AMSLER	94D	PL B333 277	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
ANISOVICH	94	PL B323 233	V.V. Anisovich <i>et al.</i>	(Crystal Barrel Collab.)
ANISOVICH	94B	PR D50 1972	V.V. Anisovich <i>et al.</i>	(LOQM)
ADAMO	93	NP A558 13C	A. Adamo <i>et al.</i>	(OBELIX Collab.)
ARMSTRONG	93C	PL B307 394	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
ARMSTRONG	93D	PL B307 399	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
WEIDENAUER	93	ZPHY C59 387	P. Weidenauer <i>et al.</i>	(ASTERIX Collab.)
ADAMO	92	PL B287 368	A. Adamo <i>et al.</i>	(OBELIX Collab.)
BELADIDZE	92B	ZPHY C54 367	G.M. Beladidze <i>et al.</i>	(VES Collab.)
AKER	91	PL B260 249	E. Aker <i>et al.</i>	(Crystal Barrel Collab.)
MAY	90	ZPHY C46 203	B. May <i>et al.</i>	(ASTERIX Collab.)
MAY	89	PL B225 450	B. May <i>et al.</i>	(ASTERIX Collab.) IJP
BRIDGES	86B	PRL 56 215	D.L. Bridges <i>et al.</i>	(SYRA, CASE)
BRIDGES	86C	PRL 57 1534	D.L. Bridges <i>et al.</i>	(SYRA)

---