

$\psi(3770)$

$I^G(J^{PC}) = 0^-(1^{--})$

$\psi(3770)$ MASS (MeV)

OUR FIT includes measurements of $m_{\psi(2S)}$, $m_{\psi(3770)}$, and $m_{\psi(3770)} - m_{\psi(2S)}$.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3773.7 ± 0.4 OUR FIT	Error includes scale factor of 1.4.			
3778.1 ± 0.7 OUR AVERAGE				
3778.1 $\pm 0.7 \pm 0.6$	1	AAIJ	19M LHCb	$p p \rightarrow D \bar{D} +$ anything
3779.2 $\pm 1.8 \pm 0.6$ -1.7 -0.8	2	ANASHIN	12A KEDR	$e^+ e^- \rightarrow D \bar{D}$
3775.5 $\pm 2.4 \pm 0.5$	57	AUBERT	08B BABR	$B \rightarrow D \bar{D} K$
3776 $\pm 5 \pm 4$	68	BRODZICKA	08 BELL	$B^+ \rightarrow D^0 \bar{D}^0 K^+$
3778.8 $\pm 1.9 \pm 0.9$		AUBERT	07BE BABR	$e^+ e^- \rightarrow D \bar{D} \gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
3779.8 ± 0.6	3	SHAMOV	17 RVUE	$e^+ e^- \rightarrow D \bar{D}$, hadrons
3772.0 ± 1.9	4,5	ABLIKIM	08D BES2	$e^+ e^- \rightarrow$ hadrons
3778.4 $\pm 3.0 \pm 1.3$	34	CHISTOV	04 BELL	Sup. by BRODZICKA 08

¹ Measured in prompt hadroproduction.

² Taking into account interference between the resonant and non-resonant $D \bar{D}$ production.

³ From the joint analysis of the data on the $D \bar{D}$ and inclusive hadronic cross sections in the $\psi(3770)$ region from BaBar, Belle, BES-II, CLEO and KEDR.


⁴ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = 0^\circ$.

⁵ Interference between the resonant and non-resonant $D \bar{D}$ production not taken into account.

$m_{\psi(3770)} - m_{\psi(2S)}$

OUR FIT includes measurements of $m_{\psi(2S)}$, $m_{\psi(3770)}$, and $m_{\psi(3770)} - m_{\psi(2S)}$.

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
87.6 ± 0.4 OUR FIT	Error includes scale factor of 1.4.		
86.6 ± 0.7 OUR AVERAGE	Error includes scale factor of 2.0. See the ideogram below.		
86.9 ± 0.4	1 ABLIKIM	07E BES2	$e^+ e^- \rightarrow$ hadrons
86.7 ± 0.7	ABLIKIM	06L BES2	$e^+ e^- \rightarrow$ hadrons
80 ± 2	SCHINDLER	80 MRK2	$e^+ e^-$
86 ± 2	2 BACINO	78 DLCO	$e^+ e^-$
88 ± 3	RAPIDIS	77 LGW	$e^+ e^-$



¹ BES-II $\psi(2S)$ mass subtracted (see ABLIKIM 06L).

² SPEAR $\psi(2S)$ mass subtracted (see SCHINDLER 80).

$m_{\psi}(3770) - m_{\psi}(2S)$ (MeV)

$\psi(3770)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
27.2± 1.0 OUR FIT				
27.5± 0.9 OUR AVERAGE				
24.9± 4.6±0.5		¹ ANASHIN	12A KEDR	$e^+ e^- \rightarrow D\bar{D}$
30.4± 8.5		^{2,3} ABLIKIM	08D BES2	$e^+ e^- \rightarrow$ hadrons
27 ±10 ±5	68	BRODZICKA	08 BELL	$B^+ \rightarrow D^0 \bar{D}^0 K^+$
28.5± 1.2±0.2		³ ABLIKIM	07E BES2	$e^+ e^- \rightarrow$ hadrons
23.5± 3.7±0.9		AUBERT	07BE BABR	$e^+ e^- \rightarrow D\bar{D}\gamma$
26.9± 2.4±0.3		³ ABLIKIM	06L BES2	$e^+ e^- \rightarrow$ hadrons
24 ± 5		³ SCHINDLER	80 MRK2	$e^+ e^-$
24 ± 5		³ BACINO	78 DLCO	$e^+ e^-$
28 ± 5		³ RAPIDIS	77 LGW	$e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
25.8± 1.3		⁴ SHAMOV	17 RVUE	$e^+ e^- \rightarrow D\bar{D}$, hadrons

¹ Taking into account interference between the resonant and non-resonant $D\bar{D}$ production.

² Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = 0^\circ$.

³ Interference between the resonant and non-resonant $D\bar{D}$ production not taken into account.

⁴ From the joint analysis of the data on the $D\bar{D}$ and inclusive hadronic cross sections in the $\psi(3770)$ region from BaBar, Belle, BES-II, CLEO and KEDR.

$\psi(3770)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
$\Gamma_1 D\bar{D}$	(93 $\begin{array}{l} +8 \\ -9 \end{array}$) %	S=2.0
$\Gamma_2 D^0\bar{D}^0$	(52 $\begin{array}{l} +4 \\ -5 \end{array}$) %	S=2.0
$\Gamma_3 D^+D^-$	(41 ± 4) %	S=2.0
$\Gamma_4 J/\psi X$	(5.0 ± 2.2) $\times 10^{-3}$	
$\Gamma_5 J/\psi\pi^+\pi^-$	(1.93 ± 0.28) $\times 10^{-3}$	
$\Gamma_6 J/\psi\pi^0\pi^0$	(8.0 ± 3.0) $\times 10^{-4}$	
$\Gamma_7 J/\psi\eta$	(9 ± 4) $\times 10^{-4}$	
$\Gamma_8 J/\psi\pi^0$	< 2.8 $\times 10^{-4}$	CL=90%
$\Gamma_9 e^+e^-$	(9.6 ± 0.7) $\times 10^{-6}$	S=1.3
Decays to light hadrons		
$\Gamma_{10} b_1(1235)\pi$	< 1.4 $\times 10^{-5}$	CL=90%
$\Gamma_{11} \phi\eta'$	< 7 $\times 10^{-4}$	CL=90%
$\Gamma_{12} \omega\eta'$	< 4 $\times 10^{-4}$	CL=90%
$\Gamma_{13} \rho^0\eta'$	< 6 $\times 10^{-4}$	CL=90%
$\Gamma_{14} \phi\eta$	(3.1 ± 0.7) $\times 10^{-4}$	
$\Gamma_{15} \omega\eta$	< 1.4 $\times 10^{-5}$	CL=90%
$\Gamma_{16} \rho^0\eta$	< 5 $\times 10^{-4}$	CL=90%
$\Gamma_{17} \phi\pi^0$	< 3 $\times 10^{-5}$	CL=90%
$\Gamma_{18} \omega\pi^0$	< 6 $\times 10^{-4}$	CL=90%
$\Gamma_{19} \pi^+\pi^-\pi^0$	< 5 $\times 10^{-6}$	CL=90%
$\Gamma_{20} \rho\pi$	< 5 $\times 10^{-6}$	CL=90%
$\Gamma_{21} K^+K^-$		
$\Gamma_{22} K^*(892)^+K^- + \text{c.c.}$	< 1.4 $\times 10^{-5}$	CL=90%
$\Gamma_{23} K^*(892)^0\bar{K}^0 + \text{c.c.}$	< 1.2 $\times 10^{-3}$	CL=90%
$\Gamma_{24} K_S^0K_L^0$	< 1.2 $\times 10^{-5}$	CL=90%
$\Gamma_{25} 2(\pi^+\pi^-)$	< 1.12 $\times 10^{-3}$	CL=90%
$\Gamma_{26} 2(\pi^+\pi^-)\pi^0$	< 1.06 $\times 10^{-3}$	CL=90%
$\Gamma_{27} 2(\pi^+\pi^-\pi^0)$	< 5.85 %	CL=90%
$\Gamma_{28} \omega\pi^+\pi^-$	< 6.0 $\times 10^{-4}$	CL=90%
$\Gamma_{29} 3(\pi^+\pi^-)$	< 9.1 $\times 10^{-3}$	CL=90%
$\Gamma_{30} 3(\pi^+\pi^-)\pi^0$	< 1.37 %	CL=90%
$\Gamma_{31} 3(\pi^+\pi^-)2\pi^0$	< 11.74 %	CL=90%
$\Gamma_{32} \eta\pi^+\pi^-$	< 1.24 $\times 10^{-3}$	CL=90%
$\Gamma_{33} \pi^+\pi^-2\pi^0$	< 8.9 $\times 10^{-3}$	CL=90%
$\Gamma_{34} \rho^0\pi^+\pi^-$	< 6.9 $\times 10^{-3}$	CL=90%
$\Gamma_{35} \eta 3\pi$	< 1.34 $\times 10^{-3}$	CL=90%
$\Gamma_{36} \eta 2(\pi^+\pi^-)$	< 2.43 %	CL=90%
$\Gamma_{37} \eta\rho^0\pi^+\pi^-$	< 1.45 %	CL=90%
$\Gamma_{38} \eta' 3\pi$	< 2.44 $\times 10^{-3}$	CL=90%

Γ_{39}	$K^+ K^- \pi^+ \pi^-$	< 9.0	$\times 10^{-4}$	CL=90%
Γ_{40}	$\phi \pi^+ \pi^-$	< 4.1	$\times 10^{-4}$	CL=90%
Γ_{41}	$K^+ K^- 2\pi^0$	< 4.2	$\times 10^{-3}$	CL=90%
Γ_{42}	$4(\pi^+ \pi^-)$	< 1.67	%	CL=90%
Γ_{43}	$4(\pi^+ \pi^-) \pi^0$	< 3.06	%	CL=90%
Γ_{44}	$\phi f_0(980)$	< 4.5	$\times 10^{-4}$	CL=90%
Γ_{45}	$K^+ K^- \pi^+ \pi^- \pi^0$	< 2.36	$\times 10^{-3}$	CL=90%
Γ_{46}	$K^+ K^- \rho^0 \pi^0$	< 8	$\times 10^{-4}$	CL=90%
Γ_{47}	$K^+ K^- \rho^+ \pi^-$	< 1.46	%	CL=90%
Γ_{48}	$\omega K^+ K^-$	< 3.4	$\times 10^{-4}$	CL=90%
Γ_{49}	$\phi \pi^+ \pi^- \pi^0$	< 3.8	$\times 10^{-3}$	CL=90%
Γ_{50}	$K^{*0} K^- \pi^+ \pi^0 + \text{c.c.}$	< 1.62	%	CL=90%
Γ_{51}	$K^{*+} K^- \pi^+ \pi^- + \text{c.c.}$	< 3.23	%	CL=90%
Γ_{52}	$K^+ K^- \pi^+ \pi^- 2\pi^0$	< 2.67	%	CL=90%
Γ_{53}	$K^+ K^- 2(\pi^+ \pi^-)$	< 1.03	%	CL=90%
Γ_{54}	$K^+ K^- 2(\pi^+ \pi^-) \pi^0$	< 3.60	%	CL=90%
Γ_{55}	$\eta K^+ K^-$	< 4.1	$\times 10^{-4}$	CL=90%
Γ_{56}	$\eta K^+ K^- \pi^+ \pi^-$	< 1.24	%	CL=90%
Γ_{57}	$\rho^0 K^+ K^-$	< 5.0	$\times 10^{-3}$	CL=90%
Γ_{58}	$2(K^+ K^-)$	< 6.0	$\times 10^{-4}$	CL=90%
Γ_{59}	$\phi K^+ K^-$	< 7.5	$\times 10^{-4}$	CL=90%
Γ_{60}	$2(K^+ K^-) \pi^0$	< 2.9	$\times 10^{-4}$	CL=90%
Γ_{61}	$2(K^+ K^-) \pi^+ \pi^-$	< 3.2	$\times 10^{-3}$	CL=90%
Γ_{62}	$K_S^0 K^- \pi^+$	< 3.2	$\times 10^{-3}$	CL=90%
Γ_{63}	$K_S^0 K^- \pi^+ \pi^0$	< 1.33	%	CL=90%
Γ_{64}	$K_S^0 K^- \rho^+$	< 6.6	$\times 10^{-3}$	CL=90%
Γ_{65}	$K_S^0 K^- 2\pi^+ \pi^-$	< 8.7	$\times 10^{-3}$	CL=90%
Γ_{66}	$K_S^0 K^- \pi^+ \rho^0$	< 1.6	%	CL=90%
Γ_{67}	$K_S^0 K^- \pi^+ \eta$	< 1.3	%	CL=90%
Γ_{68}	$K_S^0 K^- 2\pi^+ \pi^- \pi^0$	< 4.18	%	CL=90%
Γ_{69}	$K_S^0 K^- 2\pi^+ \pi^- \eta$	< 4.8	%	CL=90%
Γ_{70}	$K_S^0 K^- \pi^+ 2(\pi^+ \pi^-)$	< 1.22	%	CL=90%
Γ_{71}	$K_S^0 K^- \pi^+ 2\pi^0$	< 2.65	%	CL=90%
Γ_{72}	$K_S^0 K^- K^+ K^- \pi^+$	< 4.9	$\times 10^{-3}$	CL=90%
Γ_{73}	$K_S^0 K^- K^+ K^- \pi^+ \pi^0$	< 3.0	%	CL=90%
Γ_{74}	$K_S^0 K^- K^+ K^- \pi^+ \eta$	< 2.2	%	CL=90%
Γ_{75}	$K^{*0} K^- \pi^+ + \text{c.c.}$	< 9.7	$\times 10^{-3}$	CL=90%
Γ_{76}	$p\bar{p}$			
Γ_{77}	$p\bar{p} \pi^0$	< 4	$\times 10^{-5}$	CL=90%
Γ_{78}	$p\bar{p} \pi^+ \pi^-$	< 5.8	$\times 10^{-4}$	CL=90%
Γ_{79}	$\Lambda \bar{\Lambda}$	< 1.2	$\times 10^{-4}$	CL=90%
Γ_{80}	$p\bar{p} \pi^+ \pi^- \pi^0$	< 1.85	$\times 10^{-3}$	CL=90%
Γ_{81}	$\omega p\bar{p}$	< 2.9	$\times 10^{-4}$	CL=90%

Γ_{82}	$\Lambda\bar{\Lambda}\pi^0$	< 7	$\times 10^{-5}$	CL=90%
Γ_{83}	$p\bar{p}2(\pi^+\pi^-)$	< 2.6	$\times 10^{-3}$	CL=90%
Γ_{84}	$\eta p\bar{p}$	< 5.4	$\times 10^{-4}$	CL=90%
Γ_{85}	$\eta p\bar{p}\pi^+\pi^-$	< 3.3	$\times 10^{-3}$	CL=90%
Γ_{86}	$\rho^0 p\bar{p}$	< 1.7	$\times 10^{-3}$	CL=90%
Γ_{87}	$p\bar{p}K^+K^-$	< 3.2	$\times 10^{-4}$	CL=90%
Γ_{88}	$\eta p\bar{p}K^+K^-$	< 6.9	$\times 10^{-3}$	CL=90%
Γ_{89}	$\pi^0 p\bar{p}K^+K^-$	< 1.2	$\times 10^{-3}$	CL=90%
Γ_{90}	$\phi p\bar{p}$	< 1.3	$\times 10^{-4}$	CL=90%
Γ_{91}	$\Lambda\bar{\Lambda}\pi^+\pi^-$	< 2.5	$\times 10^{-4}$	CL=90%
Γ_{92}	$\Lambda\bar{p}K^+$	< 2.8	$\times 10^{-4}$	CL=90%
Γ_{93}	$\Lambda\bar{p}K^+\pi^+\pi^-$	< 6.3	$\times 10^{-4}$	CL=90%
Γ_{94}	$\Lambda\bar{\Lambda}\eta$	< 1.9	$\times 10^{-4}$	CL=90%
Γ_{95}	$\Sigma^+\bar{\Sigma}^-$	< 1.0	$\times 10^{-4}$	CL=90%
Γ_{96}	$\Sigma^0\bar{\Sigma}^0$	< 4	$\times 10^{-5}$	CL=90%
Γ_{97}	$\Xi^+\bar{\Xi}^-$	< 1.5	$\times 10^{-4}$	CL=90%
Γ_{98}	$\Xi^0\bar{\Xi}^0$	< 1.4	$\times 10^{-4}$	CL=90%

Radiative decays

Γ_{99}	$\gamma\chi_{c2}$	< 6.4	$\times 10^{-4}$	CL=90%
Γ_{100}	$\gamma\chi_{c1}$	(2.49 \pm 0.23)	$\times 10^{-3}$	
Γ_{101}	$\gamma\chi_{c0}$	(6.9 \pm 0.6)	$\times 10^{-3}$	
Γ_{102}	$\gamma\eta_c$	< 7	$\times 10^{-4}$	CL=90%
Γ_{103}	$\gamma\eta_c(2S)$	< 9	$\times 10^{-4}$	CL=90%
Γ_{104}	$\gamma\eta'$	< 1.8	$\times 10^{-4}$	CL=90%
Γ_{105}	$\gamma\eta$	< 1.5	$\times 10^{-4}$	CL=90%
Γ_{106}	$\gamma\pi^0$	< 2	$\times 10^{-4}$	CL=90%

CONSTRAINED FIT INFORMATION

An overall fit to the total width, a partial width, and 3 branching ratios uses 23 measurements and one constraint to determine 5 parameters. The overall fit has a $\chi^2 = 20.1$ for 19 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

$$\begin{array}{c|ccc} & & & \\ & & & \\ x_3 & 99 & & \\ & & & \\ x_9 & 0 & 0 & \\ & & & \\ \Gamma & 0 & 0 & -44 \\ & & & \\ & x_2 & x_3 & x_9 \end{array}$$

Mode		Rate (MeV)	Scale factor
Γ_2	$D^0 \bar{D}^0$	14.0 ± 1.4	1.8
Γ_3	$D^+ D^-$	11.2 ± 1.1	1.7
Γ_9	$e^+ e^-$	$(2.62 \pm 0.18) \times 10^{-4}$	1.4

$\psi(3770)$ PARTIAL WIDTHS

$\Gamma(e^+ e^-)$

Γ_9

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.262 ± 0.018 OUR FIT		Error includes scale factor of 1.4.		
0.256 ± 0.016 OUR AVERAGE		Error includes scale factor of 1.2.		
0.154 ^{+0.079 +0.021} -0.058 -0.027	1,2	ANASHIN	12A	KEDR $e^+ e^- \rightarrow D\bar{D}$
0.22 ± 0.05	3,4	ABLIKIM	08D	BES2 $e^+ e^- \rightarrow$ hadrons
0.277 ± 0.011 ± 0.013	4	ABLIKIM	07E	BES2 $e^+ e^- \rightarrow$ hadrons
0.203 ± 0.003 ^{+0.041} -0.027	1.4M	4,5 BESSON	06	CLEO $e^+ e^- \rightarrow$ hadrons
0.276 ± 0.050	4	SCHINDLER	80	MRK2 $e^+ e^-$
0.18 ± 0.06	4	BACINO	78	DLCO $e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.196 ± 0.018	6	SHAMOV	17	RVUE $e^+ e^- \rightarrow D\bar{D}$, hadrons
0.414 ^{+0.072 +0.093} -0.080 -0.028	2,7	ANASHIN	12A	KEDR $e^+ e^- \rightarrow D\bar{D}$
0.37 ± 0.09	8	RAPIDIS	77	LGW $e^+ e^-$

¹ Solution I of the two solutions.

² Taking into account interference between the resonant and non-resonant $D\bar{D}$ production.

³ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = 0^\circ$.

⁴ Interference between the resonant and non-resonant $D\bar{D}$ production not taken into account.

⁵ BESSON 06 (as corrected in BESSON 10) measure $\sigma(e^+ e^- \rightarrow \psi(3770) \rightarrow$ hadrons) = $6.36 \pm 0.08^{+0.41}_{-0.30}$ nb at $\sqrt{s} = 3773 \pm 1$ MeV, and obtain Γ_{ee} from the Born-level cross section calculated using $\psi(3770)$ mass and width from our 2004 edition, PDG 04.

⁶ From the joint analysis of the data on the $D\bar{D}$ and inclusive hadronic cross sections in the $\psi(3770)$ region from BaBar, Belle, BES-II, CLEO and KEDR.

⁷ Solution II of the two solutions.

⁸ See also $\Gamma(e^+ e^-)/\Gamma_{\text{total}}$ below.

$\psi(3770)$ BRANCHING RATIOS

$\Gamma(D\bar{D})/\Gamma_{\text{total}}$

$\Gamma_1/\Gamma = (\Gamma_2 + \Gamma_3)/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.93^{+0.08} -0.09 OUR FIT		Error includes scale factor of 2.0.		
0.93^{+0.08} -0.09 OUR AVERAGE		Error includes scale factor of 2.1.		
0.849 ± 0.056 ± 0.018	1	ABLIKIM	08B	BES2 $e^+ e^- \rightarrow$ non- $D\bar{D}$
1.033 ± 0.014 ^{+0.048} -0.066	1.427M	2 BESSON	06	CLEO $e^+ e^- \rightarrow$ hadrons

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.836 ± 0.049	³ SHAMOV	17	RVUE	$e^+ e^- \rightarrow D\bar{D}$, hadrons
$0.866 \pm 0.050 \pm 0.036$	^{4,5} ABLIKIM	07K	BES2	$e^+ e^- \rightarrow \text{non-}D\bar{D}$
$0.836 \pm 0.073 \pm 0.042$	⁵ ABLIKIM	06L	BES2	$e^+ e^- \rightarrow D\bar{D}$
$0.855 \pm 0.017 \pm 0.058$	^{5,6} ABLIKIM	06N	BES2	$e^+ e^- \rightarrow D\bar{D}$

¹ Neglecting interference.

² Obtained by comparing a measurement of the total cross section (corrected in BESSON 10) with that of $D\bar{D}$ reported by CLEO in DOBBS 07.

³ From the joint analysis of the data on the $D\bar{D}$ and inclusive hadronic cross sections in the $\psi(3770)$ region from BaBar, Belle, BES-II, CLEO and KEDR.

⁴ Using $\sigma^{obs} = 7.07 \pm 0.58$ nb and neglecting interference.

⁵ Not independent of ABLIKIM 08B.

⁶ From a measurement of $\sigma(e^+ e^- \rightarrow D\bar{D})$ at $\sqrt{s} = 3773$ MeV, using the $\psi(3770)$ resonance parameters measured by ABLIKIM 06L.

$\Gamma(D^0\bar{D}^0)/\Gamma_{\text{total}}$

Γ_2/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.52 ± 0.04 OUR FIT	Error includes scale factor of 2.0.		

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.467 \pm 0.047 \pm 0.023$	ABLIKIM	06L	BES2	$e^+ e^- \rightarrow D^0\bar{D}^0$
$0.499 \pm 0.013 \pm 0.038$	¹ ABLIKIM	06N	BES2	$e^+ e^- \rightarrow D^0\bar{D}^0$

¹ From a measurement of $\sigma(e^+ e^- \rightarrow D\bar{D})$ at $\sqrt{s} = 3773$ MeV, using the $\psi(3770)$ resonance parameters measured by ABLIKIM 06L.

$\Gamma(D^+ D^-)/\Gamma_{\text{total}}$

Γ_3/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.41 ± 0.04 OUR FIT	Error includes scale factor of 2.0.		

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.369 \pm 0.037 \pm 0.028$	ABLIKIM	06L	BES2	$e^+ e^- \rightarrow D^+ D^-$
$0.357 \pm 0.011 \pm 0.034$	¹ ABLIKIM	06N	BES2	$e^+ e^- \rightarrow D^+ D^-$

¹ From a measurement of $\sigma(e^+ e^- \rightarrow D\bar{D})$ at $\sqrt{s} = 3773$ MeV, using the $\psi(3770)$ resonance parameters measured by ABLIKIM 06L.

$\Gamma(D^0\bar{D}^0)/\Gamma(D^+ D^-)$

Γ_2/Γ_3

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.253 ± 0.016 OUR FIT				
1.253 ± 0.016 OUR AVERAGE				

$1.252 \pm 0.009 \pm 0.013$	5.3M	BONVICINI	14	CLEO	$e^+ e^- \rightarrow D\bar{D}$
$1.39 \pm 0.31 \pm 0.12$		PAKHLOVA	08	BELL	$10.6 e^+ e^- \rightarrow D\bar{D}\gamma$
$1.78 \pm 0.33 \pm 0.24$		AUBERT	07BE	BABR	$e^+ e^- \rightarrow D\bar{D}\gamma$
$1.27 \pm 0.12 \pm 0.08$		ABLIKIM	06L	BES2	$e^+ e^- \rightarrow D\bar{D}$
$2.43 \pm 1.50 \pm 0.43$	34	¹ CHISTOV	04	BELL	$B^+ \rightarrow \psi(3770) K^+$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.258 \pm 0.016 \pm 0.014$		² DOBBS	07	CLEO	$e^+ e^- \rightarrow D\bar{D}$
-----------------------------	--	--------------------	----	------	--------------------------------

¹ See ADLER 88C for older measurements of this quantity.

² Superseded by BONVICINI 14.

$\Gamma(J/\psi X)/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE (%)	DOCUMENT ID	TECN	COMMENT
0.5±0.2±0.1	1 ABLIKIM 21Z BES3	e ⁺ e ⁻ → ℓ ⁺ ℓ ⁻ X	

¹ From a fit to the $e^+ e^- \rightarrow J/\psi X$ cross section between 3.645 and 3.891 GeV, with $\psi(2S)$ and $\psi(3770)$ masses, total widths and leptonic widths fixed to the values from the PDG 20. An alternative fit with an improved χ^2 , corresponding to a significance of 5.3σ , uses an additional resonance with a mass of $3766.2 \pm 3.8 \pm 0.4$ MeV/c², a total width of $22.2 \pm 5.9 \pm 1.4$ MeV, and $\Gamma(e^+ e^-) \cdot B(J/\psi X) = 79.4 \pm 85.5 \pm 11.7$ eV, possibly compatible with the results of ABLIKIM 08H.

 $\Gamma(J/\psi \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.93±0.28 OUR AVERAGE				
$1.89 \pm 0.20 \pm 0.20$	231 ± 33	ADAM	06	CLEO $e^+ e^- \rightarrow \psi(3770)$
$3.4 \pm 1.4 \pm 0.9$	17.8 ± 4.8	BAI	05	BES2 $e^+ e^- \rightarrow \psi(3770)$

 $\Gamma(J/\psi \pi^0 \pi^0)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
0.080±0.025±0.016	39 ± 14	ADAM	06	CLEO $e^+ e^- \rightarrow \psi(3770)$

 $\Gamma(J/\psi \eta)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
87±33±22	22 ± 10	ADAM	06	CLEO $e^+ e^- \rightarrow \psi(3770)$

 $\Gamma(J/\psi \pi^0)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<28	90	<10	ADAM	06	CLEO $e^+ e^- \rightarrow \psi(3770)$

 $\Gamma(e^+ e^-)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
0.96±0.07 OUR FIT	Error includes scale factor of 1.3.		
1.3 ±0.2	RAPIDIS	77 LGW	$e^+ e^-$

— DECAYS TO LIGHT HADRONS — $\Gamma(b_1(1235)\pi)/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<1.4	90	1 ADAMS	06	CLEO $e^+ e^- \rightarrow \psi(3770)$

¹ Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb.

 $\Gamma(\phi\eta')/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<7	90	1 ADAMS	06	CLEO $e^+ e^- \rightarrow \psi(3770)$

¹ Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb.

$\Gamma(\omega\eta')/\Gamma_{\text{total}}$ Γ_{12}/Γ

<i>VALUE</i> (units 10^{-4})	<i>CL%</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
<4	90	1 ADAMS 06	CLEO	$e^+ e^- \rightarrow \psi(3770)$

¹ Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb.

 $\Gamma(\rho^0\eta')/\Gamma_{\text{total}}$ Γ_{13}/Γ

<i>VALUE</i> (units 10^{-4})	<i>CL%</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
<6	90	1 ADAMS 06	CLEO	$e^+ e^- \rightarrow \psi(3770)$

¹ Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb.

 $\Gamma(\phi\eta)/\Gamma_{\text{total}}$ Γ_{14}/Γ

<i>VALUE</i> (units 10^{-4})	<i>CL%</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
3.1±0.6±0.3	90	1 ADAMS 06	CLEO	$3.773 e^+ e^- \rightarrow \phi\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<i>VALUE</i> (units 10^{-4})	<i>CL%</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
<19	90	2 ABLIKIM 07B BES2	$e^+ e^- \rightarrow \psi(3770)$	

¹ Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb.

² Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

 $\Gamma(\omega\eta)/\Gamma_{\text{total}}$ Γ_{15}/Γ

<i>VALUE</i> (units 10^{-5})	<i>CL%</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
<1.4	90	1 ADAMS 06	CLEO	$e^+ e^- \rightarrow \psi(3770)$

¹ Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb.

 $\Gamma(\rho^0\eta)/\Gamma_{\text{total}}$ Γ_{16}/Γ

<i>VALUE</i> (units 10^{-4})	<i>CL%</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
<5	90	1 ADAMS 06	CLEO	$e^+ e^- \rightarrow \psi(3770)$

¹ Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb.

 $\Gamma(\phi\pi^0)/\Gamma_{\text{total}}$ Γ_{17}/Γ

<i>VALUE</i> (units 10^{-5})	<i>CL%</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
< 3	90	1 ADAMS 06	CLEO	$e^+ e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<i>VALUE</i> (units 10^{-5})	<i>CL%</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
<50	90	2 ABLIKIM 07B BES2	$e^+ e^- \rightarrow \psi(3770)$	

¹ Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb.

² Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

 $\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$ Γ_{18}/Γ

<i>VALUE</i> (units 10^{-4})	<i>CL%</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
<6	90	1 ADAMS 06	CLEO	$e^+ e^- \rightarrow \psi(3770)$

¹ Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb.

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{19}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<5	90	1,2 ADAMS	06 CLEO	$e^+e^- \rightarrow \psi(3770)$

¹ Data suggest possible destructive interference with continuum.² Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb. $\Gamma(\rho\pi)/\Gamma_{\text{total}}$ Γ_{20}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<5	90	1,2 ADAMS	06 CLEO	$e^+e^- \rightarrow \psi(3770)$

¹ Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb.² Data suggest possible destructive interference with continuum. $\Gamma(K^+K^-)/\Gamma_{\text{total}}$ Γ_{21}/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			

 $\sim 10^{-5}$ ¹ DRUZHININ 15 RVUE $e^+e^- \rightarrow \psi(3770)$ ¹ DRUZHININ 15 uses BABAR and CLEO data taking into account interference of the processes $e^+e^- \rightarrow K^+K^-$ and $e^+e^- \rightarrow K_S^0K_L^0$. $\Gamma(K^*(892)^+K^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{22}/Γ

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.4	90	1 ADAMS	06 CLEO	$e^+e^- \rightarrow \psi(3770)$

¹ Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb. $\Gamma(K^*(892)^0\bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{23}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.2	90	1 ADAMS	06 CLEO	$e^+e^- \rightarrow \psi(3770)$

¹ Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb. $\Gamma(K_S^0K_L^0)/\Gamma_{\text{total}}$ Γ_{24}/Γ

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 1.2	90	1 CRONIN-HEN..06	CLEO	$e^+e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $<21 \quad 90 \quad 2 \text{ ABLIKIM} \quad 04F \text{ BES} \quad e^+e^- \rightarrow \psi(3770)$ ¹ Using $\sigma(e^+e^- \rightarrow \psi(3770) \rightarrow \text{hadrons}) = (6.38 \pm 0.08^{+0.41}_{-0.30})$ nb from BESSON 06 and $B(K_S^0 \rightarrow \pi^+\pi^-) = 0.6895 \pm 0.0014$.² Using $B(K_S^0 \rightarrow \pi^+\pi^-) = 0.6860 \pm 0.0027$. $\Gamma(2(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{25}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<11.2	90	1 HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<48 90 ² ABLIKIM 07B BES2 $e^+ e^- \rightarrow \psi(3770)$

¹ Using $\sigma_{tot}(e^+ e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

² Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

$\Gamma(2(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}$

Γ_{26}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<10.6	90	¹ HUANG	06A CLEO	$e^+ e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<62 90 ² ABLIKIM 07B BES2 $e^+ e^- \rightarrow \psi(3770)$

¹ Using $\sigma_{tot}(e^+ e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

² Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

$\Gamma(2(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}$

Γ_{27}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<58.5	90	305	ABLIKIM	08N BES2	$e^+ e^- \rightarrow \psi(3770)$

$\Gamma(\omega\pi^+\pi^-)/\Gamma_{\text{total}}$

Γ_{28}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 6.0	90	¹ HUANG	06A CLEO	$e^+ e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<55 90 ² ABLIKIM 07I BES2 $3.77 e^+ e^-$

¹ Using $\sigma_{tot}(e^+ e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

² Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

$\Gamma(3(\pi^+\pi^-))/\Gamma_{\text{total}}$

Γ_{29}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<91	90	¹ ABLIKIM	07B BES2	$e^+ e^- \rightarrow \psi(3770)$

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

$\Gamma(3(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}$

Γ_{30}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<137	90	¹ ABLIKIM	07B BES2	$e^+ e^- \rightarrow \psi(3770)$

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

$\Gamma(3(\pi^+\pi^-)2\pi^0)/\Gamma_{\text{total}}$

Γ_{31}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<117.4	90	59	ABLIKIM	08N BES2	$e^+ e^- \rightarrow \psi(3770)$

$\Gamma(\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{32}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.24	90	¹ HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.3	90	² ABLIKIM	10D BES2	$e^+e^- \rightarrow \psi(3770)$
------	----	----------------------	----------	---------------------------------

¹ Using $\sigma_{tot}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

² Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

 $\Gamma(\pi^+\pi^-2\pi^0)/\Gamma_{\text{total}}$ Γ_{33}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<8.9	90	218	ABLIKIM	08N BES2	$e^+e^- \rightarrow \psi(3770)$

 $\Gamma(\rho^0\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{34}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<6.9	90	¹ ABLIKIM	07F BES2	$e^+e^- \rightarrow \psi(3770)$

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

 $\Gamma(\eta 3\pi)/\Gamma_{\text{total}}$ Γ_{35}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<13.4	90	¹ HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$

¹ Using $\sigma_{tot}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

 $\Gamma(\eta 2(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{36}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<243	90	¹ ABLIKIM	07B BES2	$e^+e^- \rightarrow \psi(3770)$

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

 $\Gamma(\eta\rho^0\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{37}/Γ

<u>VALUE (units 10^{-2})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.45	90	¹ ABLIKIM	10D BES2	$e^+e^- \rightarrow \psi(3770)$

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

 $\Gamma(\eta' 3\pi)/\Gamma_{\text{total}}$ Γ_{38}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<24.4	90	¹ HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$

¹ Using $\sigma_{tot}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

 $\Gamma(K^+K^-\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{39}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 9.0	90	¹ HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<48 90 ² ABLIKIM 07B BES2 $e^+ e^- \rightarrow \psi(3770)$

¹ Using $\sigma_{tot}(e^+ e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

² Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

$\Gamma(\phi\pi^+\pi^-)/\Gamma_{\text{total}}$

Γ_{40}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 4.1	90	¹ HUANG	06A CLEO	$e^+ e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<16 90 ² ABLIKIM 07B BES2 $e^+ e^- \rightarrow \psi(3770)$

¹ Using $\sigma_{tot}(e^+ e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

² Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

$\Gamma(K^+ K^- 2\pi^0)/\Gamma_{\text{total}}$

Γ_{41}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<4.2	90	14	ABLIKIM	08N BES2	$e^+ e^- \rightarrow \psi(3770)$

$\Gamma(4(\pi^+\pi^-))/\Gamma_{\text{total}}$

Γ_{42}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<16.7	90	¹ ABLIKIM	07F BES2	$e^+ e^- \rightarrow \psi(3770)$

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

$\Gamma(4(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}$

Γ_{43}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<30.6	90	¹ ABLIKIM	07F BES2	$e^+ e^- \rightarrow \psi(3770)$

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

$\Gamma(\phi f_0(980))/\Gamma_{\text{total}}$

Γ_{44}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<4.5	90	¹ HUANG	06A CLEO	$e^+ e^- \rightarrow \psi(3770)$

¹ Using $\sigma_{tot}(e^+ e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

$\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$

Γ_{45}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 23.6	90	¹ HUANG	06A CLEO	$e^+ e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<111 90 ² ABLIKIM 07B BES2 $e^+ e^- \rightarrow \psi(3770)$

¹ Using $\sigma_{tot}(e^+ e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

² Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

$\Gamma(K^+ K^- \rho^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{46}/Γ

<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<8	90	¹ ABLIKIM	07I	BES2 $e^+ e^-$

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

 $\Gamma(K^+ K^- \rho^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{47}/Γ

<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<146	90	¹ ABLIKIM	07I	BES2 $e^+ e^-$

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

 $\Gamma(\omega K^+ K^-)/\Gamma_{\text{total}}$ Γ_{48}/Γ

<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 3.4	90	¹ HUANG	06A	CLEO $e^+ e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<66	90	² ABLIKIM	07I	BES2 $e^+ e^-$
---------------	----	----------------------	-----	----------------

¹ Using $\sigma_{tot}(e^+ e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

² Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

 $\Gamma(\phi \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$ Γ_{49}/Γ

<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<38	90	¹ ABLIKIM	07I	BES2 $e^+ e^-$

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

 $\Gamma(K^{*0} K^- \pi^+ \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{50}/Γ

<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<162	90	¹ ABLIKIM	07I	BES2 $e^+ e^-$

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

 $\Gamma(K^{*+} K^- \pi^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{51}/Γ

<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<323	90	¹ ABLIKIM	07I	BES2 $e^+ e^-$

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

 $\Gamma(K^+ K^- \pi^+ \pi^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_{52}/Γ

<u>VALUE</u> (units 10^{-3})	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<26.7	90	24	ABLIKIM	08N	BES2 $e^+ e^- \rightarrow \psi(3770)$

$\Gamma(K^+ K^- 2(\pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{53}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<10.3	90	¹ ABLIKIM	07F BES2	$e^+ e^- \rightarrow \psi(3770)$

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

 $\Gamma(K^+ K^- 2(\pi^+ \pi^-) \pi^0)/\Gamma_{\text{total}}$ Γ_{54}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<36.0	90	¹ ABLIKIM	07F BES2	$e^+ e^- \rightarrow \psi(3770)$

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

 $\Gamma(\eta K^+ K^-)/\Gamma_{\text{total}}$ Γ_{55}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 4.1	90	¹ HUANG	06A CLEO	$e^+ e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<31	90	² ABLIKIM	10D BES2	$e^+ e^- \rightarrow \psi(3770)$
---------------	----	----------------------	----------	----------------------------------

¹ Using $\sigma_{tot}(e^+ e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

² Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

 $\Gamma(\eta K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{56}/Γ

<u>VALUE (units 10^{-2})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.24	90	¹ ABLIKIM	10D BES2	$e^+ e^- \rightarrow \psi(3770)$

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

 $\Gamma(\rho^0 K^+ K^-)/\Gamma_{\text{total}}$ Γ_{57}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<5.0	90	¹ ABLIKIM	07F BES2	$e^+ e^- \rightarrow \psi(3770)$

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

 $\Gamma(2(K^+ K^-))/\Gamma_{\text{total}}$ Γ_{58}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 6.0	90	¹ HUANG	06A CLEO	$e^+ e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<17	90	² ABLIKIM	07B BES2	$e^+ e^- \rightarrow \psi(3770)$
---------------	----	----------------------	----------	----------------------------------

¹ Using $\sigma_{tot}(e^+ e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

² Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

 $\Gamma(\phi K^+ K^-)/\Gamma_{\text{total}}$ Γ_{59}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 7.5	90	¹ HUANG	06A CLEO	$e^+ e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<24 90 ² ABLIKIM 07B BES2 $e^+ e^- \rightarrow \psi(3770)$

¹ Using $\sigma_{tot}(e^+ e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

² Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

$\Gamma(2(K^+ K^-)\pi^0)/\Gamma_{\text{total}}$

Γ_{60}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 2.9	90	1 HUANG	06A CLEO	$e^+ e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<46 90 ² ABLIKIM 07B BES2 $e^+ e^- \rightarrow \psi(3770)$

¹ Using $\sigma_{tot}(e^+ e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

² Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

$\Gamma(2(K^+ K^-)\pi^+\pi^-)/\Gamma_{\text{total}}$

Γ_{61}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
< 3.2	90	1 ABLIKIM	07F BES2	$e^+ e^- \rightarrow \psi(3770)$

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

$\Gamma(K_S^0 K^- \pi^+)/\Gamma_{\text{total}}$

Γ_{62}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 3.2	90	18	ABLIKIM	08M BES2	$e^+ e^- \rightarrow \psi(3770)$

$\Gamma(K_S^0 K^- \pi^+\pi^0)/\Gamma_{\text{total}}$

Γ_{63}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 13.3	90	40	ABLIKIM	08M BES2	$e^+ e^- \rightarrow \psi(3770)$

$\Gamma(K_S^0 K^- \rho^+)/\Gamma_{\text{total}}$

Γ_{64}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
< 6.6	90	ABLIKIM	09C BES2	$e^+ e^- \rightarrow \psi(3770)$

$\Gamma(K_S^0 K^- 2\pi^+ \pi^-)/\Gamma_{\text{total}}$

Γ_{65}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 8.7	90	39	ABLIKIM	08M BES2	$e^+ e^- \rightarrow \psi(3770)$

$\Gamma(K_S^0 K^- \pi^+ \rho^0)/\Gamma_{\text{total}}$

Γ_{66}/Γ

VALUE (units 10^{-2})	CL%	DOCUMENT ID	TECN	COMMENT
< 1.6	90	ABLIKIM	09C BES2	$e^+ e^- \rightarrow \psi(3770)$

$\Gamma(K_S^0 K^- \pi^+ \eta)/\Gamma_{\text{total}}$

Γ_{67}/Γ

VALUE (units 10^{-2})	CL%	DOCUMENT ID	TECN	COMMENT
< 1.3	90	ABLIKIM	09C BES2	$e^+ e^- \rightarrow \psi(3770)$

$\Gamma(K_S^0 K^- 2\pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	CL%	EVTS
<41.8	90	23

DOCUMENT ID	TECN	COMMENT
ABLIKIM	08M BES2	$e^+ e^- \rightarrow \psi(3770)$

 Γ_{68}/Γ $\Gamma(K_S^0 K^- 2\pi^+ \pi^- \eta)/\Gamma_{\text{total}}$

VALUE (units 10^{-2})	CL%
<4.8	90

DOCUMENT ID	TECN	COMMENT
ABLIKIM	09C BES2	$e^+ e^- \rightarrow \psi(3770)$

 Γ_{69}/Γ $\Gamma(K_S^0 K^- \pi^+ 2(\pi^+ \pi^-))/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	CL%	EVTS
<12.2	90	4

DOCUMENT ID	TECN	COMMENT
ABLIKIM	08M BES2	$e^+ e^- \rightarrow \psi(3770)$

 Γ_{70}/Γ $\Gamma(K_S^0 K^- \pi^+ 2\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	CL%	EVTS
<26.5	90	17

DOCUMENT ID	TECN	COMMENT
ABLIKIM	08M BES2	$e^+ e^- \rightarrow \psi(3770)$

 Γ_{71}/Γ $\Gamma(K_S^0 K^- K^+ K^- \pi^+)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	CL%
<4.9	90

DOCUMENT ID	TECN	COMMENT
ABLIKIM	09C BES2	$e^+ e^- \rightarrow \psi(3770)$

 Γ_{72}/Γ $\Gamma(K_S^0 K^- K^+ K^- \pi^+ \pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-2})	CL%
<3.0	90

DOCUMENT ID	TECN	COMMENT
ABLIKIM	09C BES2	$e^+ e^- \rightarrow \psi(3770)$

 Γ_{73}/Γ $\Gamma(K_S^0 K^- K^+ K^- \pi^+ \eta)/\Gamma_{\text{total}}$

VALUE (units 10^{-2})	CL%
<2.2	90

DOCUMENT ID	TECN	COMMENT
ABLIKIM	09C BES2	$e^+ e^- \rightarrow \psi(3770)$

 Γ_{74}/Γ $\Gamma(K^{*0} K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	CL%
<9.7	90

DOCUMENT ID	TECN	COMMENT
1 ABLIKIM	07F BES2	$e^+ e^- \rightarrow \psi(3770)$

 Γ_{75}/Γ $\Gamma(p\bar{p})/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	EVTS
• • • We do not use the following data for averages, fits, limits, etc. • • •	

DOCUMENT ID	TECN	COMMENT
not seen	1 AAIJ	$p\bar{p} \rightarrow B^+ X \rightarrow p\bar{p} K^+ X$

7.1 ± 8.6	684	2 ABLIKIM	$14L \text{ BES3 } e^+ e^- \rightarrow \psi(3770)$
310 ± 30	684	3 ABLIKIM	$14L \text{ BES3 } e^+ e^- \rightarrow \psi(3770)$

¹ AAIJ 17AD reports $B(B^+ \rightarrow \psi(3770) K^+ \rightarrow p\bar{p} K^+)/B(B^+ \rightarrow J/\psi K^+ \rightarrow p\bar{p} K^+) < 0.09$ (0.10) at 90% (95%) CL.

² Solution I of two equivalent solutions in a fit with a resonance interfering with continuum.

³ Solution II of two equivalent solutions in a fit with a resonance interfering with continuum.

$\Gamma(p\bar{p}\pi^0)/\Gamma_{\text{total}}$ Γ_{77}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 0.4	90	1,2 ABLIKIM	140 BES3	$e^+e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

59	$^{+3}_{-2} \pm 5$	1,3 ABLIKIM	140 BES3	$e^+e^- \rightarrow \psi(3770)$
----	--------------------	-------------	----------	---------------------------------

<12	90	4 ABLIKIM	07B BES2	$e^+e^- \rightarrow \psi(3770)$
-----	----	-----------	----------	---------------------------------

¹ Calculated by the authors using $\sigma(e^+e^- \rightarrow \psi(3770) \rightarrow \text{hadrons}) = 6.36 \pm 0.08^{+0.41}_{-0.30}$ nb from BESSON 10.

² Solution I of two equivalent solutions in a fit with a resonance interfering with continuum.

³ Solution II of two equivalent solutions in a fit with a resonance interfering with continuum.

⁴ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

 $\Gamma(p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{78}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 5.8	90	1 HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<16	90	2 ABLIKIM	07B BES2	$e^+e^- \rightarrow \psi(3770)$
-----	----	-----------	----------	---------------------------------

¹ Using $\sigma_{tot}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

² Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

 $\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$ Γ_{79}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.2 × 10 ⁻⁴	90	1 HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.8 × 10 ⁻⁴	90	2 ABLIKIM	21AS BES3	$e^+e^- \rightarrow \psi(3770)$
-------------------------	----	-----------	-----------	---------------------------------

<4 × 10 ⁻⁴	90	3 ABLIKIM	07F BES2	$e^+e^- \rightarrow \psi(3770)$
-----------------------	----	-----------	----------	---------------------------------

¹ Using $\sigma_{tot}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

² From a measurement of the $e^+e^- \rightarrow \Lambda\bar{\Lambda}$ cross section between 3.5 and 4.6 GeV. At a 90% CL the lower bound is $> 2.4 \times 10^{-6}$.

³ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

 $\Gamma(p\bar{p}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{80}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<18.5	90	1 HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<73	90	2 ABLIKIM	07B BES2	$e^+e^- \rightarrow \psi(3770)$
-----	----	-----------	----------	---------------------------------

¹ Using $\sigma_{tot}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

² Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

 $\Gamma(\omega p\bar{p})/\Gamma_{\text{total}}$ Γ_{81}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 2.9	90	1 HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<30 90 ² ABLIKIM 07I BES2 3.77 e⁺ e⁻

¹ Using $\sigma_{tot}(e^+ e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

² Using $\sigma^{obs} = 7.15 \pm 0.27 \pm 0.27$ nb and neglecting interference.

$\Gamma(\Lambda\bar{\Lambda}\pi^0)/\Gamma_{\text{total}}$

Γ_{82}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 0.7	90	¹ ABLIKIM	13Q	BES3 $e^+ e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<12 90 ² ABLIKIM 07I BES2 3.77 e⁺ e⁻

¹ Assuming that interference effects between resonance and continuum can be neglected.

² Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

$\Gamma(p\bar{p}2(\pi^+\pi^-))/\Gamma_{\text{total}}$

Γ_{83}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<2.6	90	¹ ABLIKIM	07F	BES2 $e^+ e^- \rightarrow \psi(3770)$

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

$\Gamma(\eta p\bar{p})/\Gamma_{\text{total}}$

Γ_{84}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 5.4	90	¹ HUANG	06A	CLEO $e^+ e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<11 90 ² ABLIKIM 10D BES2 3.77 e⁺ e⁻ $\rightarrow \psi(3770)$

¹ Using $\sigma_{tot}(e^+ e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

² Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

$\Gamma(\eta p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$

Γ_{85}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<3.3	90	¹ ABLIKIM	10D	BES2 $e^+ e^- \rightarrow \psi(3770)$

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

$\Gamma(\rho^0 p\bar{p})/\Gamma_{\text{total}}$

Γ_{86}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<1.7	90	¹ ABLIKIM	07F	BES2 $e^+ e^- \rightarrow \psi(3770)$

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

$\Gamma(p\bar{p}K^+K^-)/\Gamma_{\text{total}}$

Γ_{87}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 3.2	90	¹ HUANG	06A	CLEO $e^+ e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<11 90 ² ABLIKIM 07B BES2 $e^+ e^- \rightarrow \psi(3770)$

¹ Using $\sigma_{tot}(e^+ e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

² Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

$\Gamma(\eta p\bar{p}K^+ K^-)/\Gamma_{\text{total}}$

Γ_{88}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<6.9	90	¹ ABLIKIM	10D BES2	$e^+ e^- \rightarrow \psi(3770)$

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

$\Gamma(\pi^0 p\bar{p}K^+ K^-)/\Gamma_{\text{total}}$

Γ_{89}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<1.2	90	¹ ABLIKIM	10D BES2	$e^+ e^- \rightarrow \psi(3770)$

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

$\Gamma(\phi p\bar{p})/\Gamma_{\text{total}}$

Γ_{90}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.3	90	¹ HUANG	06A CLEO	$e^+ e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<9 90 ² ABLIKIM 07B BES2 $e^+ e^- \rightarrow \psi(3770)$

¹ Using $\sigma_{tot}(e^+ e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

² Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

$\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$

Γ_{91}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 2.5	90	¹ HUANG	06A CLEO	$e^+ e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 4.7 90 ² ABLIKIM 13Q BES3 $e^+ e^- \rightarrow \psi(3770)$

<39 90 ³ ABLIKIM 07F BES2 $e^+ e^- \rightarrow \psi(3770)$

¹ Using $\sigma_{tot}(e^+ e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

² Assuming that interference effects between resonance and continuum can be neglected.

³ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

$\Gamma(\Lambda\bar{p}K^+)/\Gamma_{\text{total}}$

Γ_{92}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<2.8	90	¹ HUANG	06A CLEO	$e^+ e^- \rightarrow \psi(3770)$

¹ Using $\sigma_{tot}(e^+ e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

$\Gamma(\Lambda\bar{p}K^+\pi^+\pi^-)/\Gamma_{\text{total}}$

Γ_{93}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<6.3	90	¹ HUANG	06A CLEO	$e^+ e^- \rightarrow \psi(3770)$

¹ Using $\sigma_{tot}(e^+ e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

$\Gamma(\Lambda\bar{\Lambda}\eta)/\Gamma_{\text{total}}$ Γ_{94}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.9	90	¹ ABLIKIM	13Q BES3	$e^+ e^- \rightarrow \psi(3770)$

¹ Assuming that interference effects between resonance and continuum can be neglected.

 $\Gamma(\Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}$ Γ_{95}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.0	90	¹ ABLIKIM	13Q BES3	$e^+ e^- \rightarrow \psi(3770)$

¹ Assuming that interference effects between resonance and continuum can be neglected.

 $\Gamma(\Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}$ Γ_{96}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.4	90	¹ ABLIKIM	13Q BES3	$e^+ e^- \rightarrow \psi(3770)$

¹ Assuming that interference effects between resonance and continuum can be neglected.

 $\Gamma(\Xi^+\bar{\Xi}^-)/\Gamma_{\text{total}}$ Γ_{97}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.5	90	¹ ABLIKIM	13Q BES3	$e^+ e^- \rightarrow \psi(3770)$

¹ Assuming that interference effects between resonance and continuum can be neglected.

 $\Gamma(\Xi^0\bar{\Xi}^0)/\Gamma_{\text{total}}$ Γ_{98}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.4	90	¹ ABLIKIM	13Q BES3	$e^+ e^- \rightarrow \psi(3770)$

¹ Assuming that interference effects between resonance and continuum can be neglected.

RADIATIVE DECAYS

 $\Gamma(\gamma\chi_{c2})/\Gamma_{\text{total}}$ Γ_{99}/Γ

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.64	90	¹ ABLIKIM	15J BES3	$e^+ e^- \rightarrow \psi(3770) \rightarrow \gamma\gamma J/\psi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.0 90 ² BRIERE 06 CLEO $e^+ e^- \rightarrow \psi(3770) \rightarrow \gamma + \text{hadrons}$

<0.9 90 ³ COAN 06A CLEO $e^+ e^- \rightarrow \psi(3770) \rightarrow \gamma\gamma J/\psi$

¹ This limit is equivalent to $(0.25 \pm 0.21 \pm 0.18) \times 10^{-3}$ branching fraction value.

² Uses $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = 9.22 \pm 0.11 \pm 0.46\%$ from ATHAR 04, $\psi(2S)$ mass and width from PDG 04, and $\Gamma_{ee}(\psi(2S)) = 2.54 \pm 0.03 \pm 0.11$ keV from ADAM 06.

³ Using $\Gamma_{ee}(\psi(2S)) = (2.54 \pm 0.03 \pm 0.11)$ keV from ADAM 06 and taking $\sigma(e^+ e^- \rightarrow D\bar{D})$ from HE 05 for $\sigma(e^+ e^- \rightarrow \psi(3770))$.

 $\Gamma(\gamma\chi_{c1})/\Gamma_{\text{total}}$ Γ_{100}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.49 ± 0.23 OUR AVERAGE				
1.98 $\pm 0.78 \pm 0.05$	202	¹ ABLIKIM	16B BES3	$e^+ e^- \rightarrow \psi(3770) \rightarrow \gamma + \text{hadrons}$
2.48 $\pm 0.15 \pm 0.23$	0.6k	ABLIKIM	15J BES3	$e^+ e^- \rightarrow \psi(3770) \rightarrow \gamma\gamma J/\psi$

$2.4 \pm 0.8 \pm 0.2$	² ABLIKIM	14H BES3	$e^+ e^- \rightarrow \psi(3770) \rightarrow K_S^0 K^\pm \pi^\mp$
$2.9 \pm 0.5 \pm 0.4$	³ BRIERE	06 CLEO	$e^+ e^- \rightarrow \psi(3770) \rightarrow \gamma + \text{hadrons}, \gamma\gamma J/\psi$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$3.9 \pm 1.4 \pm 0.6$	54	⁴ BRIERE	$e^+ e^- \rightarrow \psi(3770) \rightarrow \gamma + \text{hadrons}$
$2.8 \pm 0.5 \pm 0.4$	53	⁵ COAN	$e^+ e^- \rightarrow \psi(3770) \rightarrow \gamma\gamma J/\psi$

¹ ABLIKIM 16B reports $(1.94 \pm 0.42 \pm 0.64) \times 10^{-3}$ from a measurement of $[\Gamma(\psi(3770) \rightarrow \gamma\chi_{c1})/\Gamma_{\text{total}}] / [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ABLIKIM 14H reports $[\Gamma(\psi(3770) \rightarrow \gamma\chi_{c1})/\Gamma_{\text{total}}] \times [B(\chi_{c1}(1P) \rightarrow K_S^0 K^\pm \pi^\mp)] = (8.51 \pm 2.39 \pm 1.42) \times 10^{-6}$ which we divide by our best value $B(\chi_{c1}(1P) \rightarrow K_S^0 K^\pm \pi^\mp) = 0.00349 \pm 0.00029$. Our first error is their experiment's error and our second error is the systematic error from using our best value. We have calculated the best value of $B(\chi_{c1}(1P) \rightarrow K_S^0 K^\pm \pi^\mp)$ as 1/2 of $B(\chi_{c1}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.}) = (7.0 \pm 0.6) \times 10^{-3}$.

³ Averages the two measurements from COAN 06A and BRIERE 06.

⁴ Uses $B(\psi(2S) \rightarrow \gamma\chi_{c1}) = 9.07 \pm 0.11 \pm 0.54\%$ from ATHAR 04, $\psi(2S)$ mass and width from PDG 04, and $\Gamma_{ee}(\psi(2S)) = 2.54 \pm 0.03 \pm 0.11$ keV from ADAM 06.

⁵ Using $\Gamma_{ee}(\psi(2S)) = (2.54 \pm 0.03 \pm 0.11)$ keV from ADAM 06 and taking $\sigma(e^+ e^- \rightarrow D\bar{D})$ from HE 05 for $\sigma(e^+ e^- \rightarrow \psi(3770))$.

$\Gamma(\gamma\chi_{c1})/\Gamma(J/\psi\pi^+\pi^-)$		Γ_{100}/Γ_5		
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$1.49 \pm 0.31 \pm 0.26$	53 ± 10	¹ COAN	06A CLEO	$e^+ e^- \rightarrow \psi(3770) \rightarrow \gamma\gamma J/\psi$

¹ Using $B(\psi(3770) \rightarrow J/\psi\pi^+\pi^-) = (1.89 \pm 0.20 \pm 0.20) \times 10^{-3}$ from ADAM 06.

$\Gamma(\gamma\chi_{c0})/\Gamma_{\text{total}}$		Γ_{101}/Γ			
VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
6.9 ± 0.6 OUR AVERAGE					
6.7 $\pm 0.7 \pm 0.1$	2.2k	¹ ABLIKIM	16B BES3	$e^+ e^- \rightarrow \psi(3770) \rightarrow \gamma + \text{hadrons}$	
7.3 $\pm 0.7 \pm 0.6$	274	BRIERE	06 CLEO	$e^+ e^- \rightarrow \psi(3770) \rightarrow \gamma + \text{hadrons}$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 44	90	² COAN	06A CLEO	$e^+ e^- \rightarrow \psi(3770) \rightarrow \gamma\gamma J/\psi$
--------	----	-------------------	----------	--

¹ ABLIKIM 16B reports $(6.88 \pm 0.28 \pm 0.67) \times 10^{-3}$ from a measurement of $[\Gamma(\psi(3770) \rightarrow \gamma\chi_{c0})/\Gamma_{\text{total}}] / [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Using $\Gamma_{ee}(\psi(2S)) = (2.54 \pm 0.03 \pm 0.11)$ keV from ADAM 06 and taking $\sigma(e^+ e^- \rightarrow D\bar{D})$ from HE 05 for $\sigma(e^+ e^- \rightarrow \psi(3770))$.

$\Gamma(\gamma\chi_{c0})/\Gamma(\gamma\chi_{c2})$				Γ_{101}/Γ_{99}
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
>8	90	¹ BRIERE	06	CLEO $e^+ e^- \rightarrow \psi(3770)$

¹ Not independent of other results in BRIERE 06.

$\Gamma(\gamma\chi_{c0})/\Gamma(\gamma\chi_{c1})$				$\Gamma_{101}/\Gamma_{100}$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
2.5 ± 0.6		¹ BRIERE	06	CLEO $e^+ e^- \rightarrow \psi(3770)$

¹ Not independent of other results in BRIERE 06.

$\Gamma(\gamma\eta_c)/\Gamma_{\text{total}}$				Γ_{102}/Γ
VALUE	CL%	DOCUMENT ID	TECN	
$< 7 \times 10^{-4}$	90	¹ ABLIKIM	14H	BES3
1 ABLIKIM 14H reports $[\Gamma(\psi(3770) \rightarrow \gamma\eta_c)/\Gamma_{\text{total}}] \times [B(\eta_c(1S) \rightarrow K_S^0 K^\pm \pi^\mp)]$ $< 16 \times 10^{-6}$ which we divide by our best value $B(\eta_c(1S) \rightarrow K_S^0 K^\pm \pi^\mp) = 2.42 \times 10^{-2}$. We have calculated the best value of $B(\eta_c(1S) \rightarrow K_S^0 K^\pm \pi^\mp)$ as 1/3 of $B(\eta_c(1S) \rightarrow K\bar{K}\pi) = 7.3 \times 10^{-2}$.				

$\Gamma(\gamma\eta_c(2S))/\Gamma_{\text{total}}$				Γ_{103}/Γ
VALUE	CL%	DOCUMENT ID	TECN	
$< 9 \times 10^{-4}$	90	¹ ABLIKIM	14H	BES3
1 ABLIKIM 14H reports $[\Gamma(\psi(3770) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}] \times [B(\eta_c(2S) \rightarrow K_S^0 K^\pm \pi^\mp)]$ $< 5.6 \times 10^{-6}$ which we divide by our best value $B(\eta_c(2S) \rightarrow K_S^0 K^\pm \pi^\mp) = 6 \times 10^{-3}$. We have calculated the best value of $B(\eta_c(2S) \rightarrow K_S^0 K^\pm \pi^\mp)$ as 1/3 of $B(\eta_c(2S) \rightarrow K\bar{K}\pi) = 1.9 \times 10^{-2}$.				

$\Gamma(\gamma\eta')/\Gamma_{\text{total}}$				Γ_{104}/Γ
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 1.8	90	¹ PEDLAR	09	CLE3 $\psi(2S) \rightarrow \gamma X$

¹ Assuming maximal destructive interference between $\psi(3770)$ and continuum sources.

$\Gamma(\gamma\eta)/\Gamma_{\text{total}}$				Γ_{105}/Γ
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 1.5	90	¹ PEDLAR	09	CLE3 $\psi(2S) \rightarrow \gamma X$

¹ Assuming maximal destructive interference between $\psi(3770)$ and continuum sources.

$\Gamma(\gamma\pi^0)/\Gamma_{\text{total}}$				Γ_{106}/Γ
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 2	90	PEDLAR	09	CLE3 $\psi(2S) \rightarrow \gamma X$

$\psi(3770)$ REFERENCES

ABLIKIM 21AS	PR D104 L091104	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 21Z	PRL 127 082002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
PDG 20	PTEP 2020 083C01	P.A. Zyla <i>et al.</i>	(PDG Collab.)
AAIJ 19M	JHEP 1907 035	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ 17AD	PL B769 305	R. Aaij <i>et al.</i>	(LHCb Collab.)
SHAMOV 17	PL B769 187	A.G. Shamov, K.Yu. Todyshev	
ABLIKIM 16B	PL B753 103	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 15J	PR D91 092009	M. Ablikim <i>et al.</i>	(BESIII Collab.)
DRUZHININ 15	PR D92 054024	V.P. Druzhinin	(NOVO)
ABLIKIM 14H	PR D89 112005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 14L	PL B735 101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 14O	PR D90 032007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
BONVICINI 14	PR D89 072002	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
ABLIKIM 13Q	PR D87 112011	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ANASHIN 12A	PL B711 292	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
ABLIKIM 10D	EPJ C66 11	M. Ablikim <i>et al.</i>	(BES II Collab.)
BESSON 10	PRL 104 159901 (errat.)	D. Besson <i>et al.</i>	(CLEO Collab.)
ABLIKIM 09C	EPJ C64 243	M. Ablikim <i>et al.</i>	(BES Collab.)
PEDLAR 09	PR D79 111101	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)
ABLIKIM 08B	PL B659 74	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM 08D	PL B660 315	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM 08H	PR D101 102004	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM 08M	PL B670 179	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM 08N	PL B670 184	M. Ablikim <i>et al.</i>	(BES Collab.)
AUBERT 08B	PR D77 011102	B. Aubert <i>et al.</i>	(BABAR Collab.)
BRODZICKA 08	PRL 100 092001	J. Brodzicka <i>et al.</i>	(BELLE Collab.)
PAKHLOVA 08	PR D77 011103	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
ABLIKIM 07B	PL B650 111	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM 07E	PL B652 238	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM 07F	PL B656 30	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM 07I	EPJ C52 805	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM 07K	PR D76 122002	M. Ablikim <i>et al.</i>	(BES Collab.)
AUBERT 07BE	PR D76 111105	B. Aubert <i>et al.</i>	(BABAR Collab.)
DOBBS 07	PR D76 112001	S. Dobbs <i>et al.</i>	(CLEO Collab.)
ABLIKIM 06L	PRL 97 121801	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM 06N	PL B641 145	M. Ablikim <i>et al.</i>	(BES Collab.)
ADAM 06	PRL 96 082004	N.E. Adam <i>et al.</i>	(CLEO Collab.)
ADAMS 06	PR D73 012002	G.S. Adams <i>et al.</i>	(CLEO Collab.)
BESSON 06	PRL 96 092002	D. Besson <i>et al.</i>	(CLEO Collab.)
Also	PRL 104 159901 (errat.)	D. Besson <i>et al.</i>	(CLEO Collab.)
BRIERE 06	PR D74 031106	R.A. Briere <i>et al.</i>	(CLEO Collab.)
COAN 06A	PRL 96 182002	T.E. Coan <i>et al.</i>	(CLEO Collab.)
CRONIN-HEN... 06	PR D74 012005	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)
HUANG 06A	PRL 96 032003	G.S. Huang <i>et al.</i>	(CLEO Collab.)
BAI 05	PL B605 63	J.Z. Bai <i>et al.</i>	(BES Collab.)
HE 05	PR L 95 121801	Q. He <i>et al.</i>	(CLEO Collab.)
Also	PRL 96 199903 (errat.)	Q. He <i>et al.</i>	(CLEO Collab.)
ABLIKIM 04F	PR D70 077101	M. Ablikim <i>et al.</i>	(BES Collab.)
ATHAR 04	PR D70 112002	S.B. Athar <i>et al.</i>	(CLEO Collab.)
CHISTOV 04	PRL 93 051803	R. Chistov <i>et al.</i>	(BELLE Collab.)
PDG 04	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)
BAI 02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)
ADLER 88C	PRL 60 89	J. Adler <i>et al.</i>	(Mark III Collab.)
SCHINDLER 80	PR D21 2716	R.H. Schindler <i>et al.</i>	(Mark II Collab.)
BACINO 78	PRL 40 671	W.J. Bacino <i>et al.</i>	(SLAC, UCLA, UCI)
RAPIDIS 77	PRL 39 526	P.A. Rapidis <i>et al.</i>	(LGW Collab.)