

# CHARMED BARYONS ( $C = +1$ )

$$\begin{aligned}\Lambda_c^+ &= u d c, \quad \Sigma_c^{++} = u u c, \quad \Sigma_c^+ = u d c, \quad \Sigma_c^0 = d d c, \\ \Xi_c^+ &= u s c, \quad \Xi_c^0 = d s c, \quad \Omega_c^0 = s s c\end{aligned}$$

$\Lambda_c^+$

$$I(J^P) = 0(\frac{1}{2}^+)$$

Mass  $m = 2286.46 \pm 0.14$  MeV

Mean life  $\tau = (201.5 \pm 2.7) \times 10^{-15}$  s ( $S = 1.6$ )

$$c\tau = 60.4 \mu\text{m}$$

### Decay asymmetry parameters

$$\Lambda \pi^+ \quad \alpha = -0.84 \pm 0.09$$

$$\Sigma^+ \pi^0 \quad \alpha = -0.55 \pm 0.11$$

$$\alpha \text{ FOR } \Lambda_c^+ \rightarrow \Sigma^0 \pi^+ = -0.73 \pm 0.18$$

$$\Lambda \ell^+ \nu_\ell \quad \alpha = -0.86 \pm 0.04$$

$$\alpha \text{ FOR } \Lambda_c^+ \rightarrow p K_S^0 = 0.2 \pm 0.5$$

$$(\alpha + \bar{\alpha})/(\alpha - \bar{\alpha}) \text{ in } \Lambda_c^+ \rightarrow \Lambda \pi^+, \bar{\Lambda}_c^- \rightarrow \bar{\Lambda} \pi^- = -0.07 \pm 0.31$$

$$(\alpha + \bar{\alpha})/(\alpha - \bar{\alpha}) \text{ in } \Lambda_c^+ \rightarrow \Lambda e^+ \nu_e, \bar{\Lambda}_c^- \rightarrow \bar{\Lambda} e^- \bar{\nu}_e = 0.00 \pm 0.04$$

$$A_{CP}(\Lambda X) \text{ in } \Lambda_c \rightarrow \Lambda X, \bar{\Lambda}_c \rightarrow \bar{\Lambda} X = (2 \pm 7)\%$$

$$\Delta A_{CP} = A_{CP}(\Lambda_c^+ \rightarrow p K^+ K^-) - A_{CP}(\Lambda_c^+ \rightarrow p \pi^+ \pi^-) = (0.3 \pm 1.1)\%$$

Branching fractions marked with a footnote, e.g. [a], have been corrected for decay modes not observed in the experiments. For example, the sub-mode fraction  $\Lambda_c^+ \rightarrow p \bar{K}^*(892)^0$  seen in  $\Lambda_c^+ \rightarrow p K^- \pi^+$  has been multiplied up to include  $\bar{K}^*(892)^0 \rightarrow \bar{K}^0 \pi^0$  decays.

$\Lambda_c^+$ DECAY MODES	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor/ Confidence level	$p$ (MeV/c)
<b>Hadronic modes with a <math>p</math> or <math>n</math>: <math>S = -1</math> final states</b>			
$p K_S^0$	( $1.59 \pm 0.08$ ) %	S=1.1	873
$p K^- \pi^+$	( $6.28 \pm 0.32$ ) %	S=1.4	823
$p \bar{K}^*(892)^0$	[a] ( $1.96 \pm 0.27$ ) %		685
$\Delta(1232)^{++} K^-$	( $1.08 \pm 0.25$ ) %		710
$\Lambda(1520) \pi^+$	[a] ( $2.2 \pm 0.5$ ) %		628
$p K^- \pi^+$ nonresonant	( $3.5 \pm 0.4$ ) %		823
$p K_S^0 \pi^0$	( $1.97 \pm 0.13$ ) %	S=1.1	823
$n K_S^0 \pi^+$	( $1.82 \pm 0.25$ ) %		821

$p\bar{K}^0\eta$	$(8.3 \pm 1.8) \times 10^{-3}$	568
$pK_S^0\pi^+\pi^-$	$(1.60 \pm 0.12)\%$	S=1.1 754
$pK^-\pi^+\pi^0$	$(4.46 \pm 0.30)\%$	S=1.5 759
$pK^*(892)^-\pi^+$	[a] $(1.4 \pm 0.5)\%$	580
$p(K^-\pi^+)_{\text{nonresonant}}\pi^0$	$(4.6 \pm 0.8)\%$	759
$\Delta(1232)\bar{K}^*(892)$	seen	419
$pK^-2\pi^+\pi^-$	$(1.4 \pm 0.9) \times 10^{-3}$	671
$pK^-\pi^+2\pi^0$	$(1.0 \pm 0.5)\%$	678

**Hadronic modes with a  $p$ :  $S = 0$  final states**

$p\pi^0$	$< 8 \times 10^{-5}$	CL=90% 945
$p\eta$	$(1.42 \pm 0.12) \times 10^{-3}$	856
$p\omega(782)^0$	$(8.3 \pm 1.1) \times 10^{-4}$	751
$p\pi^+\pi^-$	$(4.61 \pm 0.28) \times 10^{-3}$	927
$p f_0(980)$	[a] $(3.5 \pm 2.3) \times 10^{-3}$	614
$p2\pi^+2\pi^-$	$(2.3 \pm 1.4) \times 10^{-3}$	852
$pK^+K^-$	$(1.06 \pm 0.06) \times 10^{-3}$	616
$p\phi$	[a] $(1.06 \pm 0.14) \times 10^{-3}$	590
$pK^+K^-\text{non-}\phi$	$(5.3 \pm 1.2) \times 10^{-4}$	616
$p\phi\pi^0$	$(10 \pm 4) \times 10^{-5}$	460
$pK^+K^-\pi^0\text{nonresonant}$	$< 6.3 \times 10^{-5}$	CL=90% 494

**Hadronic modes with a hyperon:  $S = -1$  final states**

$\Lambda\pi^+$	$(1.30 \pm 0.07)\%$	S=1.1 864
$\Lambda(1670)\pi^+, \Lambda(1670) \rightarrow \eta\Lambda$	$(3.5 \pm 0.5) \times 10^{-3}$	—
$\Lambda\pi^+\pi^0$	$(7.1 \pm 0.4)\%$	S=1.1 844
$\Lambda\rho^+$	$< 6\%$	CL=95% 636
$\Lambda\pi^-2\pi^+$	$(3.64 \pm 0.29)\%$	S=1.4 807
$\Sigma(1385)^+\pi^+\pi^-, \Sigma^{*+} \rightarrow \Lambda\pi^+$	$(1.0 \pm 0.5)\%$	688
$\Sigma(1385)^-\pi^+, \Sigma^{*-} \rightarrow \Lambda\pi^-$	$(7.6 \pm 1.4) \times 10^{-3}$	688
$\Lambda\pi^+\rho^0$	$(1.5 \pm 0.6)\%$	524
$\Sigma(1385)^+\rho^0, \Sigma^{*+} \rightarrow \Lambda\pi^+$	$(5 \pm 4) \times 10^{-3}$	363
$\Lambda\pi^-2\pi^+\text{nonresonant}$	$< 1.1\%$	CL=90% 807
$\Lambda\pi^-\pi^02\pi^+\text{total}$	$(2.3 \pm 0.8)\%$	757
$\Lambda\pi^+\eta$	[a] $(1.84 \pm 0.26)\%$	691
$\Sigma(1385)^+\eta$	[a] $(9.1 \pm 2.0) \times 10^{-3}$	570
$\Lambda\pi^+\omega$	[a] $(1.5 \pm 0.5)\%$	517
$\Lambda\pi^-\pi^02\pi^+, \text{no } \eta \text{ or } \omega$	$< 8 \times 10^{-3}$	CL=90% 757
$\Lambda K^+\bar{K}^0$	$(5.7 \pm 1.1) \times 10^{-3}$	S=1.9 443
$\Xi(1690)^0K^+, \Xi^{*0} \rightarrow \Lambda\bar{K}^0$	$(1.6 \pm 0.5) \times 10^{-3}$	286
$\Sigma^0\pi^+$	$(1.29 \pm 0.07)\%$	S=1.1 825
$\Sigma^0\pi^+\eta$	$(7.5 \pm 0.8) \times 10^{-3}$	635
$\Sigma^+\pi^0$	$(1.25 \pm 0.10)\%$	827

$\Sigma^+ \eta$	( 4.4 $\pm$ 2.0 ) $\times 10^{-3}$	713
$\Sigma^+ \eta'$	( 1.5 $\pm$ 0.6 ) %	391
$\Sigma^+ \pi^+ \pi^-$	( 4.50 $\pm$ 0.25 ) %	S=1.3
$\Sigma^+ \rho^0$	< 1.7 %	CL=95%
$\Sigma^- 2\pi^+$	( 1.87 $\pm$ 0.18 ) %	799
$\Sigma^0 \pi^+ \pi^0$	( 3.5 $\pm$ 0.4 ) %	803
$\Sigma^+ \pi^0 \pi^0$	( 1.55 $\pm$ 0.15 ) %	806
$\Sigma^0 \pi^- 2\pi^+$	( 1.11 $\pm$ 0.30 ) %	763
$\Sigma^+ \pi^+ \pi^- \pi^0$	—	767
$\Sigma^+ \omega$	[a] ( 1.70 $\pm$ 0.21 ) %	569
$\Sigma^- \pi^0 2\pi^+$	( 2.1 $\pm$ 0.4 ) %	762
$\Sigma^+ K^+ K^-$	( 3.5 $\pm$ 0.4 ) $\times 10^{-3}$	S=1.1
$\Sigma^+ \phi$	[a] ( 3.9 $\pm$ 0.6 ) $\times 10^{-3}$	S=1.1
$\Xi(1690)^0 K^+, \Xi^{*0} \rightarrow$	( 1.02 $\pm$ 0.25 ) $\times 10^{-3}$	286
$\Sigma^+ K^-$	—	—
$\Sigma^+ K^+ K^-$ nonresonant	< 8 $\times 10^{-4}$	CL=90%
$\Xi^0 K^+$	( 5.5 $\pm$ 0.7 ) $\times 10^{-3}$	653
$\Xi^- K^+ \pi^+$	( 6.2 $\pm$ 0.6 ) $\times 10^{-3}$	S=1.1
$\Xi(1530)^0 K^+$	( 4.3 $\pm$ 0.9 ) $\times 10^{-3}$	S=1.1
		473

**Hadronic modes with a hyperon:  $S = 0$  final states**

$\Lambda K^+$	( 6.1 $\pm$ 1.2 ) $\times 10^{-4}$	781
$\Lambda K^+ \pi^+ \pi^-$	< 5 $\times 10^{-4}$	CL=90%
$\Sigma^0 K^+$	( 5.2 $\pm$ 0.8 ) $\times 10^{-4}$	735
$\Sigma^0 K^+ \pi^+ \pi^-$	< 2.6 $\times 10^{-4}$	CL=90%
$\Sigma^+ K^+ \pi^-$	( 2.1 $\pm$ 0.6 ) $\times 10^{-3}$	670
$\Sigma^+ K^*(892)^0$	[a] ( 3.5 $\pm$ 1.0 ) $\times 10^{-3}$	470
$\Sigma^- K^+ \pi^+$	< 1.2 $\times 10^{-3}$	CL=90%
		664

**Doubly Cabibbo-suppressed modes**

$p K^+ \pi^-$	( 1.11 $\pm$ 0.18 ) $\times 10^{-4}$	823
---------------	--------------------------------------	-----

**Semileptonic modes**

$\Lambda e^+ \nu_e$	( 3.6 $\pm$ 0.4 ) %	871
$\Lambda \mu^+ \nu_\mu$	( 3.5 $\pm$ 0.5 ) %	867

**Inclusive modes**

$e^+$ anything	( 3.95 $\pm$ 0.35 ) %	—
$p$ anything	( 50 $\pm$ 16 ) %	—
$n$ anything	( 50 $\pm$ 16 ) %	—
$\Lambda$ anything	( 38.2 $\pm$ 2.9 ) %	—
$K_S^0$ anything	( 9.9 $\pm$ 0.7 ) %	—
3prongs	( 24 $\pm$ 8 ) %	—

**$\Delta C = 1$  weak neutral current (*C1*) modes, or  
Lepton Family number (*LF*), or Lepton number (*L*), or  
Baryon number (*B*) violating modes**

$p e^+ e^-$	<i>C1</i>	< 5.5	$\times 10^{-6}$	CL=90%	951
$p \mu^+ \mu^-$ non-resonant	<i>C1</i>	< 7.7	$\times 10^{-8}$	CL=90%	937
$p e^+ \mu^-$	<i>LF</i>	< 9.9	$\times 10^{-6}$	CL=90%	947
$p e^- \mu^+$	<i>LF</i>	< 1.9	$\times 10^{-5}$	CL=90%	947
$\bar{p} 2e^+$	<i>L,B</i>	< 2.7	$\times 10^{-6}$	CL=90%	951
$\bar{p} 2\mu^+$	<i>L,B</i>	< 9.4	$\times 10^{-6}$	CL=90%	937
$\bar{p} e^+ \mu^+$	<i>L,B</i>	< 1.6	$\times 10^{-5}$	CL=90%	947
$\Sigma^- \mu^+ \mu^+$	<i>L</i>	< 7.0	$\times 10^{-4}$	CL=90%	812

 **$\Lambda_c(2595)^+$** 

$$I(J^P) = 0(\frac{1}{2}^-)$$

The spin-parity follows from the fact that  $\Sigma_c(2455)\pi$  decays, with little available phase space, are dominant. This assumes that  $J^P = 1/2^+$  for the  $\Sigma_c(2455)$ .

Mass  $m = 2592.25 \pm 0.28$  MeV

$m - m_{\Lambda_c^+} = 305.79 \pm 0.24$  MeV

Full width  $\Gamma = 2.6 \pm 0.6$  MeV

$\Lambda_c^+ \pi \pi$  and its submode  $\Sigma_c(2455)\pi$  — the latter just barely — are the only strong decays allowed to an excited  $\Lambda_c^+$  having this mass; and the submode seems to dominate.

<b><math>\Lambda_c(2595)^+</math> DECAY MODES</b>	Fraction ( $\Gamma_i/\Gamma$ )	<i>p</i> (MeV/c)
$\Lambda_c^+ \pi^+ \pi^-$	[b] —	117
$\Sigma_c(2455)^{++} \pi^-$	$24 \pm 7\%$	3
$\Sigma_c(2455)^0 \pi^+$	$24 \pm 7\%$	3
$\Lambda_c^+ \pi^+ \pi^-$ 3-body	$18 \pm 10\%$	117
$\Lambda_c^+ \pi^0$	[c] not seen	258
$\Lambda_c^+ \gamma$	not seen	288

 **$\Lambda_c(2625)^+$** 

$$I(J^P) = 0(\frac{3}{2}^-)$$

$J^P$  has not been measured;  $\frac{3}{2}^-$  is the quark-model prediction.

Mass  $m = 2628.11 \pm 0.19$  MeV (S = 1.1)

$m - m_{\Lambda_c^+} = 341.65 \pm 0.13$  MeV (S = 1.1)

Full width  $\Gamma < 0.97$  MeV, CL = 90%

$\Lambda_c^+ \pi\pi$  and its submode  $\Sigma(2455)\pi$  are the only strong decays allowed to an excited  $\Lambda_c^+$  having this mass.

$\Lambda_c(2625)^+$ DECAY MODES	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level	$p$ (MeV/c)
$\Lambda_c^+ \pi^+ \pi^-$	$\approx 67\%$		184
$\Sigma_c(2455)^{++} \pi^-$	<5	90%	103
$\Sigma_c(2455)^0 \pi^+$	<5	90%	103
$\Lambda_c^+ \pi^+ \pi^-$ 3-body	large		184
$\Lambda_c^+ \pi^0$	[c] not seen		293
$\Lambda_c^+ \gamma$	not seen		319

### $\Lambda_c(2860)^+$

$$I(J^P) = 0(\frac{3}{2}^+)$$

Mass  $m = 2856.1^{+2.3}_{-6.0}$  MeV

Full width  $\Gamma = 68^{+12}_{-22}$  MeV

$\Lambda_c(2860)^+$ DECAY MODES	Fraction ( $\Gamma_i/\Gamma$ )	$p$ (MeV/c)
$D^0 p$	seen	259

### $\Lambda_c(2880)^+$

$$I(J^P) = 0(\frac{5}{2}^+)$$

Mass  $m = 2881.63 \pm 0.24$  MeV

$m - m_{\Lambda_c^+} = 595.17 \pm 0.28$  MeV

Full width  $\Gamma = 5.6^{+0.8}_{-0.6}$  MeV

$\Lambda_c(2880)^+$ DECAY MODES	Fraction ( $\Gamma_i/\Gamma$ )	$p$ (MeV/c)
$\Lambda_c^+ \pi^+ \pi^-$	seen	471
$\Sigma_c(2455)^0,^{++} \pi^\pm$	seen	376
$\Sigma_c(2520)^0,^{++} \pi^\pm$	seen	317
$p D^0$	seen	316

### $\Lambda_c(2940)^+$

$$I(J^P) = 0(\frac{3}{2}^-)$$

$J^P = 3/2^-$  is favored, but is not certain

Mass  $m = 2939.6^{+1.3}_{-1.5}$  MeV

Full width  $\Gamma = 20^{+6}_{-5}$  MeV

<b><math>\Lambda_c(2940)^+</math> DECAY MODES</b>	Fraction ( $\Gamma_i/\Gamma$ )	$p$ (MeV/c)
$p D^0$	seen	420
$\Sigma_c(2455)^0, ++ \pi^\pm$	seen	—

 **$\Sigma_c(2455)$** 

$$I(J^P) = 1(\frac{1}{2}^+)$$

$\Sigma_c(2455)^{++}$  mass  $m = 2453.97 \pm 0.14$  MeV  
 $\Sigma_c(2455)^+$  mass  $m = 2452.65^{+0.22}_{-0.16}$  MeV  
 $\Sigma_c(2455)^0$  mass  $m = 2453.75 \pm 0.14$  MeV  
 $m_{\Sigma_c(2455)^{++}} - m_{\Lambda_c^+} = 167.510 \pm 0.017$  MeV  
 $m_{\Sigma_c(2455)^+} - m_{\Lambda_c^+} = 166.19^{+0.16}_{-0.08}$  MeV  
 $m_{\Sigma_{c2455}^0} - m_{\Lambda_c^+} = 167.290 \pm 0.017$  MeV  
 $m_{\Sigma_c(2455)^{++}} - m_{\Sigma_c(2455)^0} = 0.220 \pm 0.013$  MeV  
 $m_{\Sigma_c(2455)^+} - m_{\Sigma_c(2455)^0} = -1.10^{+0.16}_{-0.08}$  MeV  
 $\Sigma_c(2455)^{++}$  full width  $\Gamma = 1.89^{+0.09}_{-0.18}$  MeV (S = 1.1)  
 $\Sigma_c(2455)^+$  full width  $\Gamma = 2.3 \pm 0.4$  MeV  
 $\Sigma_c(2455)^0$  full width  $\Gamma = 1.83^{+0.11}_{-0.19}$  MeV (S = 1.2)

$\Lambda_c^+ \pi$  is the only strong decay allowed to a  $\Sigma_c$  having this mass.

<b><math>\Sigma_c(2455)</math> DECAY MODES</b>	Fraction ( $\Gamma_i/\Gamma$ )	$p$ (MeV/c)
$\Lambda_c^+ \pi$	$\approx 100$ %	94

 **$\Sigma_c(2520)$** 

$$I(J^P) = 1(\frac{3}{2}^+)$$

$J^P$  has not been measured;  $\frac{3}{2}^+$  is the quark-model prediction.

$\Sigma_c(2520)^{++}$  mass  $m = 2518.41^{+0.22}_{-0.18}$  MeV (S = 1.1)  
 $\Sigma_c(2520)^+$  mass  $m = 2517.4^{+0.7}_{-0.5}$  MeV  
 $\Sigma_c(2520)^0$  mass  $m = 2518.48 \pm 0.20$  MeV (S = 1.1)  
 $m_{\Sigma_c(2520)^{++}} - m_{\Lambda_c^+} = 231.95^{+0.18}_{-0.12}$  MeV (S = 1.3)  
 $m_{\Sigma_c(2520)^+} - m_{\Lambda_c^+} = 230.9^{+0.7}_{-0.5}$  MeV  
 $m_{\Sigma_c(2520)^0} - m_{\Lambda_c^+} = 232.02^{+0.16}_{-0.14}$  MeV (S = 1.3)  
 $m_{\Sigma_c(2520)^{++}} - m_{\Sigma_c(2520)^0} = 0.01 \pm 0.15$  MeV  
 $\Sigma_c(2520)^{++}$  full width  $\Gamma = 14.78^{+0.30}_{-0.40}$  MeV  
 $\Sigma_c(2520)^+$  full width  $\Gamma = 17.2^{+4.0}_{-2.2}$  MeV  
 $\Sigma_c(2520)^0$  full width  $\Gamma = 15.3^{+0.4}_{-0.5}$  MeV

$\Lambda_c^+ \pi$  is the only strong decay allowed to a  $\Sigma_c$  having this mass.

$\Sigma_c(2520)$ DECAY MODES	Fraction ( $\Gamma_i/\Gamma$ )	$p$ (MeV/c)
$\Lambda_c^+ \pi$	$\approx 100\%$	179

### $\Sigma_c(2800)$

$$I(J^P) = 1(?)$$

$\Sigma_c(2800)^{++}$  mass  $m = 2801^{+4}_{-6}$  MeV

$\Sigma_c(2800)^+$  mass  $m = 2792^{+14}_{-5}$  MeV

$\Sigma_c(2800)^0$  mass  $m = 2806^{+5}_{-7}$  MeV (S = 1.3)

$m_{\Sigma_c(2800)^{++}} - m_{\Lambda_c^+} = 514^{+4}_{-6}$  MeV

$m_{\Sigma_c(2800)^+} - m_{\Lambda_c^+} = 505^{+14}_{-5}$  MeV

$m_{\Sigma_c(2800)^0} - m_{\Lambda_c^+} = 519^{+5}_{-7}$  MeV (S = 1.3)

$\Sigma_c(2800)^{++}$  full width  $\Gamma = 75^{+22}_{-17}$  MeV

$\Sigma_c(2800)^+$  full width  $\Gamma = 62^{+60}_{-40}$  MeV

$\Sigma_c(2800)^0$  full width  $\Gamma = 72^{+22}_{-15}$  MeV

$\Sigma_c(2800)$ DECAY MODES	Fraction ( $\Gamma_i/\Gamma$ )	$p$ (MeV/c)
$\Lambda_c^+ \pi$	seen	443

### $\Xi_c^+$

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

$J^P$  has not been measured;  $\frac{1}{2}^+$  is the quark-model prediction.

Mass  $m = 2467.71 \pm 0.23$  MeV (S = 1.3)

Mean life  $\tau = (453 \pm 5) \times 10^{-15}$  s

$$c\tau = 135.8 \mu\text{m}$$

Branching fractions marked with a footnote, e.g. [a], have been corrected for decay modes not observed in the experiments. For example, the sub-mode fraction  $\Xi_c^+ \rightarrow \Sigma^+ \bar{K}^*(892)^0$  seen in  $\Xi_c^+ \rightarrow \Sigma^+ K^- \pi^+$  has been multiplied up to include  $\bar{K}^*(892)^0 \rightarrow \bar{K}^0 \pi^0$  decays.

$\Xi_c^+$ DECAY MODES	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor/ Confidence level	$p$ (MeV/c)
-----------------------	--------------------------------	-----------------------------------	----------------


### Cabibbo-favored (S = -2) decays

$p 2 K_S^0$	$(2.5 \pm 1.3) \times 10^{-3}$	766
$\Lambda \bar{K}^0 \pi^+$	—	852
$\Sigma(1385)^+ \bar{K}^0$	[a] $(2.9 \pm 2.0)\%$	746

$\Lambda K^- 2\pi^+$	$(9 \pm 4) \times 10^{-3}$		787
$\Lambda \bar{K}^*(892)^0 \pi^+$	[a] $< 5 \times 10^{-3}$	CL=90%	608
$\Sigma(1385)^+ K^- \pi^+$	[a] $< 6 \times 10^{-3}$	CL=90%	678
$\Sigma^+ K^- \pi^+$	$(2.7 \pm 1.2) \%$		810
$\Sigma^+ \bar{K}^*(892)^0$	[a] $(2.3 \pm 1.1) \%$		658
$\Sigma^0 K^- 2\pi^+$	$(8 \pm 5) \times 10^{-3}$		735
$\Xi^0 \pi^+$	$(1.6 \pm 0.8) \%$		876
$\Xi^- 2\pi^+$	$(2.9 \pm 1.3) \%$		851
$\Xi(1530)^0 \pi^+$	[a] $< 2.9 \times 10^{-3}$	CL=90%	749
$\Xi(1620)^0 \pi^+$	seen		—
$\Xi(1690)^0 \pi^+$	seen		644
$\Xi^0 \pi^+ \pi^0$	$(6.7 \pm 3.5) \%$		856
$\Xi^0 \pi^- 2\pi^+$	$(5.0 \pm 2.6) \%$		818
$\Xi^0 e^+ \nu_e$	$(7 \pm 4) \%$		884
$\Omega^- K^+ \pi^+$	$(2.0 \pm 1.5) \times 10^{-3}$		399

**Cabibbo-suppressed decays**

$p K^- \pi^+$	$(6.2 \pm 3.0) \times 10^{-3}$	S=1.5	944
$p \bar{K}^*(892)^0$	[a] $(3.3 \pm 1.7) \times 10^{-3}$		828
$\Sigma^+ \pi^+ \pi^-$	$(1.4 \pm 0.8) \%$		922
$\Sigma^- 2\pi^+$	$(5.1 \pm 3.4) \times 10^{-3}$		918
$\Sigma^+ K^+ K^-$	$(4.3 \pm 2.5) \times 10^{-3}$		579
$\Sigma^+ \phi$	[a] $< 3.2 \times 10^{-3}$	CL=90%	549
$\Xi(1690)^0 K^+, \Xi^0 \rightarrow$	$< 1.3 \times 10^{-3}$	CL=90%	501
$\Sigma^+ K^-$			
$p \phi(1020)$	$(1.2 \pm 0.6) \times 10^{-4}$		751



$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

$J^P$  has not been measured;  $\frac{1}{2}^+$  is the quark-model prediction.

Mass  $m = 2470.44 \pm 0.28$  MeV (S = 1.2)

$m_{\Xi_c^0} - m_{\Xi_c^+} = 2.72 \pm 0.23$  MeV (S = 1.1)

Mean life  $\tau = (151.9 \pm 2.4) \times 10^{-15}$  s

$c\tau = 45.5 \mu\text{m}$

**Decay asymmetry parameters**

$\Xi^- \pi^+$   $\alpha = -0.64 \pm 0.05$

$\alpha$  FOR  $\Xi_c^0 \rightarrow \Xi^+ \pi^- = 0.61 \pm 0.05$

$\alpha$  FOR  $\Xi_c^0 \rightarrow \Lambda \bar{K}^*(892)^0 = 0.15 \pm 0.22$

$\alpha$  FOR  $\Xi_c^0 \rightarrow \Sigma^+ K^*(892)^- = -0.52 \pm 0.30$

$\Xi_c^0$ DECAY MODES	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor	$p$ (MeV/c)
<b>Cabibbo-favored decays</b>			
$p K^- K^- \pi^+$	$(4.8 \pm 1.2) \times 10^{-3}$	1.1	676
$p K^- \bar{K}^*(892)^0, \bar{K}^{*0} \rightarrow K^- \pi^+$	$(2.0 \pm 0.6) \times 10^{-3}$		413
$p K^- K^- \pi^+ (\text{no } \bar{K}^{*0})$	$(3.0 \pm 0.9) \times 10^{-3}$		676
$\Lambda K_S^0$	$(3.2 \pm 0.7) \times 10^{-3}$		906
$\Lambda K^- \pi^+$	$(1.45 \pm 0.33) \%$	1.1	856
$\Lambda \bar{K}^*(892)^0$	$(2.6 \pm 0.7) \times 10^{-3}$		717
$\Lambda \bar{K}^0 \pi^+ \pi^-$	seen		786
$\Lambda K^- \pi^+ \pi^+ \pi^-$	seen		703
$\Sigma^0 K_S^0$	$(5.4 \pm 1.6) \times 10^{-4}$		864
$\Sigma^+ K^-$	$(1.8 \pm 0.4) \times 10^{-3}$		868
$\Sigma^0 \bar{K}^*(892)^0$	$(9.8 \pm 2.3) \times 10^{-3}$		658
$\Sigma^+ K^*(892)^-$	$(4.9 \pm 1.4) \times 10^{-3}$		661
$\Xi^- \pi^+$	$(1.43 \pm 0.32) \%$	1.1	875
$\Xi^- \pi^+ \pi^+ \pi^-$	$(4.8 \pm 2.3) \%$		816
$\Xi^0 \phi, \phi \rightarrow K^+ K^-$	$(5.1 \pm 1.3) \times 10^{-4}$		—
$\Xi^0 K^+ K^- \text{ nonresonant}$	$(5.6 \pm 1.4) \times 10^{-4}$		444
$\Omega^- K^+$	$(4.2 \pm 1.0) \times 10^{-3}$		522
$\Xi^- e^+ \nu_e$	$(1.04 \pm 0.24) \%$		882
$\Xi^- \mu^+ \nu_\mu$	$(1.01 \pm 0.25) \%$		878
<b>Cabibbo-suppressed decays</b>			
$\Lambda_c^+ \pi^-$	$(5.5 \pm 1.8) \times 10^{-3}$		115
$\Xi^- K^+$	$(3.9 \pm 1.2) \times 10^{-4}$		789
$\Lambda K^+ K^- (\text{no } \phi)$	$(4.1 \pm 1.4) \times 10^{-4}$		648
$\Lambda \phi$	$(4.9 \pm 1.5) \times 10^{-4}$		621

 $\Xi_c'^+$ 

$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$

 $J^P$  has not been measured;  $\frac{1}{2}^+$  is the quark-model prediction.Mass  $m = 2578.2 \pm 0.5$  MeV (S = 1.1)

$m_{\Xi_c'^+} - m_{\Xi_c^+} = 110.5 \pm 0.4$  MeV

$m_{\Xi_c'^+} - m_{\Xi_c^0} = -0.5 \pm 0.6$  MeV

The  $\Xi_c'^+ - \Xi_c^+$  mass difference is too small for any strong decay to occur.

$\Xi_c'^+$ DECAY MODES	Fraction ( $\Gamma_i/\Gamma$ )	$p$ (MeV/c)
$\Xi_c^+ \gamma$	seen	108

$\Xi_c'^0$ 

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

$J^P$  has not been measured;  $\frac{1}{2}^+$  is the quark-model prediction.

Mass  $m = 2578.7 \pm 0.5$  MeV

$$m_{\Xi_c'^0} - m_{\Xi_c^0} = 108.3 \pm 0.4$$
 MeV

The  $\Xi_c'^0 - \Xi_c^0$  mass difference is too small for any strong decay to occur.

### $\Xi_c'^0$ DECAY MODES

Fraction ( $\Gamma_i/\Gamma$ )

$p$  (MeV/c)

$$\Xi_c^0 \gamma$$

seen

106

 $\Xi_c(2645)$ 

$$I(J^P) = \frac{1}{2}(\frac{3}{2}^+)$$

$J^P$  has not been measured;  $\frac{3}{2}^+$  is the quark-model prediction.

$\Xi_c(2645)^+$  mass  $m = 2645.10 \pm 0.30$  MeV ( $S = 1.2$ )

$\Xi_c(2645)^0$  mass  $m = 2646.16 \pm 0.25$  MeV ( $S = 1.3$ )

$$m_{\Xi_c(2645)^+} - m_{\Xi_c^0} = 174.67 \pm 0.09$$
 MeV

$$m_{\Xi_c(2645)^0} - m_{\Xi_c^+} = 178.45 \pm 0.10$$
 MeV

$$m_{\Xi_c(2645)^+} - m_{\Xi_c(2645)^0} = -1.06 \pm 0.27$$
 MeV ( $S = 1.1$ )

$\Xi_c(2645)^+$  full width  $\Gamma = 2.14 \pm 0.19$  MeV ( $S = 1.1$ )

$\Xi_c(2645)^0$  full width  $\Gamma = 2.35 \pm 0.22$  MeV

$\Xi_c \pi$  is the only strong decay allowed to a  $\Xi_c$  resonance having this mass.

### $\Xi_c(2645)$ DECAY MODES

Fraction ( $\Gamma_i/\Gamma$ )

$p$  (MeV/c)

$$\Xi_c^0 \pi^+$$

seen

102

$$\Xi_c^+ \pi^-$$

seen

106

 $\Xi_c(2790)$ 

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^-)$$

$J^P$  has not been measured;  $\frac{1}{2}^-$  is the quark-model prediction.

$\Xi_c(2790)^+$  mass  $= 2791.9 \pm 0.5$  MeV

$\Xi_c(2790)^0$  mass  $= 2793.9 \pm 0.5$  MeV

$$m_{\Xi_c(2790)^+} - m_{\Xi_c'^0} = 213.20 \pm 0.22$$
 MeV

$$m_{\Xi_c(2790)^0} - m_{\Xi_c'^+} = 215.70 \pm 0.22$$
 MeV

$$m_{\Xi_c(2790)^+} - m_{\Xi_c(2790)^0} = -2.0 \pm 0.7$$
 MeV

$\Xi_c(2790)^+$  width  $= 8.9 \pm 1.0$  MeV

$\Xi_c(2790)^0$  width  $= 10.0 \pm 1.1$  MeV

$\Xi_c(2790)$ DECAY MODES	Fraction ( $\Gamma_i/\Gamma$ )	$p$ (MeV/c)
$\Xi'_c \pi$	seen	159

 **$\Xi_c(2815)$** 

$$I(J^P) = \frac{1}{2}(\frac{3}{2}^-)$$

$J^P$  has not been measured;  $\frac{3}{2}^-$  is the quark-model prediction.

$$\Xi_c(2815)^+ \text{ mass } m = 2816.51 \pm 0.25 \text{ MeV } (S = 1.2)$$

$$\Xi_c(2815)^0 \text{ mass } m = 2819.79 \pm 0.30 \text{ MeV } (S = 1.1)$$

$$m_{\Xi_c(2815)^+} - m_{\Xi_c^+} = 348.80 \pm 0.10 \text{ MeV}$$

$$m_{\Xi_c(2815)^0} - m_{\Xi_c^0} = 349.35 \pm 0.11 \text{ MeV}$$

$$m_{\Xi_c(2815)^+} - m_{\Xi_c(2815)^0} = -3.27 \pm 0.27 \text{ MeV}$$

$$\Xi_c(2815)^+ \text{ full width } \Gamma = 2.43 \pm 0.26 \text{ MeV}$$

$$\Xi_c(2815)^0 \text{ full width } \Gamma = 2.54 \pm 0.25 \text{ MeV}$$

The  $\Xi_c \pi \pi$  modes are consistent with being entirely via  $\Xi_c(2645)\pi$ .

$\Xi_c(2815)$ DECAY MODES	Fraction ( $\Gamma_i/\Gamma$ )	$p$ (MeV/c)
$\Xi'_c \pi$	seen	188
$\Xi_c(2645)\pi$	seen	102
$\Xi_c^0 \gamma$	seen	325

 **$\Xi_c(2970)$** 

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

was  $\Xi_c(2980)$

$$\Xi_c(2970)^+ \text{ } m = 2964.3 \pm 1.5 \text{ MeV } (S = 3.9)$$

$$\Xi_c(2970)^0 \text{ } m = 2967.1 \pm 1.7 \text{ MeV } (S = 6.7)$$

$$m_{\Xi_c(2970)^+} - m_{\Xi_c^+} = 496.6 \pm 1.5 \text{ MeV } (S = 3.7)$$

$$m_{\Xi_c(2970)^0} - m_{\Xi_c^0} = 496.7 \pm 1.8 \text{ MeV } (S = 5.3)$$

$$m_{\Xi_c(2970)^+} - m_{\Xi_c(2970)^0} = -2.8 \pm 1.9 \text{ MeV } (S = 4.8)$$

$$\Xi_c(2970)^+ \text{ width } \Gamma = 20.9^{+2.4}_{-3.5} \text{ MeV } (S = 1.2)$$

$\Xi_c(2970)$ DECAY MODES	Fraction ( $\Gamma_i/\Gamma$ )	$p$ (MeV/c)
$\Lambda_c^+ \bar{K} \pi$	seen	223
$\Sigma_c(2455) \bar{K}$	seen	122
$\Lambda_c^+ \bar{K}$	not seen	410

$\Lambda_c^+ K^-$	seen	410
$\Xi_c^- 2\pi$	seen	381
$\Xi_c' \pi$	seen	—
$\Xi_c(2645)\pi$	seen	274

### $\Xi_c(3055)$

$$I(J^P) = ?(?)$$

Mass  $m = 3055.9 \pm 0.4$  MeV

Full width  $\Gamma = 7.8 \pm 1.9$  MeV

$\Xi_c(3055)$ DECAY MODES	Fraction ( $\Gamma_i/\Gamma$ )	$p$ (MeV/c)
$\Sigma^{++} K^-$	seen	—
$\Lambda D^+$	seen	316

### $\Xi_c(3080)$

$$I(J^P) = \frac{1}{2}(?)$$

$\Xi_c(3080)^+ m = 3077.2 \pm 0.4$  MeV

$\Xi_c(3080)^0 m = 3079.9 \pm 1.4$  MeV ( $S = 1.3$ )

$\Xi_c(3080)^+ \text{ width } \Gamma = 3.6 \pm 1.1$  MeV ( $S = 1.5$ )

$\Xi_c(3080)^0 \text{ width } \Gamma = 5.6 \pm 2.2$  MeV

$\Xi_c(3080)$ DECAY MODES	Fraction ( $\Gamma_i/\Gamma$ )	$p$ (MeV/c)
$\Lambda_c^+ \bar{K}\pi$	seen	415
$\Sigma_c(2455)\bar{K}$	seen	342
$\Sigma_c(2455)^{++} K^-$	seen	342
$\Sigma_c(2520)^{++} K^-$	seen	239
$\Sigma_c(2455)\bar{K} + \Sigma_c(2520)\bar{K}$	seen	—
$\Lambda_c^+ \bar{K}$	not seen	536
$\Lambda_c^+ \bar{K}\pi^+\pi^-$	not seen	144
$\Lambda D^+$	seen	362

### $\Omega_c^0$

$$I(J^P) = 0(\frac{1}{2}^+)$$

$J^P$  has not been measured;  $\frac{1}{2}^+$  is the quark-model prediction.

Mass  $m = 2695.2 \pm 1.7$  MeV ( $S = 1.3$ )

Mean life  $\tau = (268 \pm 26) \times 10^{-15}$  s

$c\tau = 80 \mu\text{m}$

No absolute branching fractions have been measured. The following are branching *rations* relative to  $\Omega^- \pi^+$ .

$\Omega_c^0$ DECAY MODES	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level	$p$ (MeV/c)
<b>Cabibbo-favored (<math>S = -3</math>) decays — relative to <math>\Omega^- \pi^+</math></b>			
$\Omega^- \pi^+$	DEFINED AS 1		821
$\Omega^- \pi^+ \pi^0$	$1.80 \pm 0.33$		797
$\Omega^- \rho^+$	$>1.3$	90%	532
$\Omega^- \pi^- 2\pi^+$	$0.31 \pm 0.05$		753
$\Omega^- e^+ \nu_e$	$2.4 \pm 1.2$		829
$\Xi^0 \bar{K}^0$	$1.64 \pm 0.29$		950
$\Xi^0 K^- \pi^+$	$1.20 \pm 0.18$		901
$\Xi^0 \bar{K}^{*0}, \bar{K}^{*0} \rightarrow K^- \pi^+$	$0.68 \pm 0.16$		764
$\Omega(2012)^- \pi^+, \Omega(2012)^- \rightarrow$	$0.12 \pm 0.05$		—
$\Xi^- K^-$			
$\Xi^- \bar{K}^0 \pi^+$	$2.12 \pm 0.28$		895
$\Omega(2012)^- \pi^+, \Omega(2012)^- \rightarrow$	$0.12 \pm 0.06$		—
$\Xi^- \bar{K}^0$			
$\Xi^- K^- 2\pi^+$	$0.63 \pm 0.09$		830
$\Xi(1530)^0 K^- \pi^+, \Xi^{*0} \rightarrow$	$0.21 \pm 0.06$		757
$\Xi^- \bar{K}^{*0} \pi^+$	$0.34 \pm 0.11$		653
$p K^- K^- \pi^+$	seen		864
$\Sigma^+ K^- K^- \pi^+$	$<0.32$	90%	689
$\Lambda \bar{K}^0 \bar{K}^0$	$1.72 \pm 0.35$		837

**$\Omega_c(2770)^0$**

$I(J^P) = 0(\frac{3}{2}^+)$

$J^P$  has not been measured;  $\frac{3}{2}^+$  is the quark-model prediction.

Mass  $m = 2765.9 \pm 2.0$  MeV ( $S = 1.2$ )

$$m_{\Omega_c(2770)^0} - m_{\Omega_c^0} = 70.7^{+0.8}_{-0.9} \text{ MeV}$$

The  $\Omega_c(2770)^0 - \Omega_c^0$  mass difference is too small for any strong decay to occur.

$\Omega_c(2770)^0$ DECAY MODES	Fraction ( $\Gamma_i/\Gamma$ )	$p$ (MeV/c)
$\Omega_c^0 \gamma$	presumably 100%	70

**$\Omega_c(3000)^0$**

$$I(J^P) = ?(?^?)$$

Mass  $m = 3000.41 \pm 0.22$  MeV

Full width  $\Gamma = 4.5 \pm 0.7$  MeV

**$\Omega_c(3000)^0$  DECAY MODES**

Fraction ( $\Gamma_i/\Gamma$ )

$p$  (MeV/c)

$\Xi_c^+ K^-$

seen

182

**$\Omega_c(3050)^0$**

$$I(J^P) = ?(?^?)$$

Mass  $m = 3050.19 \pm 0.13$  MeV

Full width  $\Gamma < 1.2$  MeV, CL = 95%

**$\Omega_c(3050)^0$  DECAY MODES**

Fraction ( $\Gamma_i/\Gamma$ )

$p$  (MeV/c)

$\Xi_c^+ K^-$

seen

278

**$\Omega_c(3065)^0$**

$$I(J^P) = ?(?^?)$$

Mass  $m = 3065.54 \pm 0.26$  MeV

Full width  $\Gamma = 3.3 \pm 0.6$  MeV (S = 1.5)

**$\Omega_c(3065)^0$  DECAY MODES**

Fraction ( $\Gamma_i/\Gamma$ )

$p$  (MeV/c)

$\Xi_c^+ K^-$

seen

303

**$\Omega_c(3090)^0$**

$$I(J^P) = ?(?^?)$$

Mass  $m = 3090.1 \pm 0.5$  MeV

Full width  $\Gamma = 8.7 \pm 1.3$  MeV

**$\Omega_c(3090)^0$  DECAY MODES**

Fraction ( $\Gamma_i/\Gamma$ )

$p$  (MeV/c)

$\Xi_c^+ K^-$

seen

340

**$\Omega_c(3120)^0$**

$I(J^P) = ?(?)$

Mass  $m = 3119.1 \pm 1.0$  MeV

Full width  $\Gamma < 2.6$  MeV, CL = 95%

<b><math>\Omega_c(3120)^0</math> DECAY MODES</b>	Fraction ( $\Gamma_i/\Gamma$ )	$p$ (MeV/c)
$\Xi_c^+ K^-$	seen	379

#### NOTES

[a] This branching fraction includes all the decay modes of the final-state resonance.

[b] See AALTONEN 11H, Fig. 8, for the calculated ratio of  $\Lambda_c^+ \pi^0 \pi^0$  and  $\Lambda_c^+ \pi^+ \pi^-$  partial widths as a function of the  $\Lambda_c(2595)^+ - \Lambda_c^+$  mass difference. At our value of the mass difference, the ratio is about 4.

[c] A test that the isospin is indeed 0, so that the particle is indeed a  $\Lambda_c^+$ .